

Paleodiversity of Chondrichthyan Faunas from the Neogene of Tropical America

Dissertation

zur

Erlangung der naturwissenschaftlichen Doktorwürde

(Dr. sc. Nat)

Vorgelegt der

Mathematisch-naturwissenschaftlichen Fakultät

der

Universität Zürich

Von

Jorge Domingo Carrillo Briceño

aus

Venezuela

Promotionskomitee

Prof. Dr. Marcelo R. Sánchez-Villagra

(Leitung der Dissertation und Vorsitz)

Dr. Lionel Cavin

Prof. Dr. Christian Klug

PD. Dr. Torsten M. Scheyer

Zürich, 2016

PALAEODIVERSITY OF CHONDRICHTHYAN FAUNAS FROM THE NEOGENE OF TROPICAL AMERICA



Jorge Domingo Carrillo Briceño

Universität Zürich, 2016

Title page image:

Restoration of diverse sharks and rays in coastal lagoon-estuarine environment at late Miocene times in Urumaco.

Source: Carrillo-Briceño et al. (2015b). Artwork: Jorge González, 2015.

CONTENTS

ACKNOWLEDGEMENTS	4
SUMMARY	5
ZUSAMMENFASSUNG	8
AUTHOR CONTRIBUTIONS	12
CHAPTER 1: Introduction	17
CHAPTER 2: Fossil Chondrichthyes from the central Eastern Pacific Ocean and their paleoceanographic significance	35
CHAPTER 3: A new Late Miocene chondrichthyan assemblage from the Chagres Formation, Panama	51
CHAPTER 4: Sawfishes and Other Elasmobranch Assemblages from the Mio-Pliocene of the South Caribbean (Urumaco Sequence, Northwestern Venezuela)	67
CHAPTER 5: A new early Miocene (Aquitanian) Elasmobranchii assemblage from the La Guajira Peninsula, Colombia	103
CHAPTER 6: An Early Neogene Elasmobranch fauna from the southern Caribbean (Western Venezuela)	127
CHAPTER 7: Conclusions and future perspectives	161
APPENDIX I: Late Neogene elasmobranch fauna from the Coquimbo Formation, Chile	175
APPENDIX II: The Path towards Endangered Species: Prehistoric Fisheries in Southeastern Brazil	189

ACKNOWLEDGEMENTS

A sincere thanks and appreciation to my adviser Prof. Dr. Marcelo Sánchez-Villagra, who granted me the opportunity and economic support for my Ph.D. studies and the opportunity to become a member of his research group. Thank you very much for so many years of good friendship, confidence, sincerity, cooperation, and especially for the many teachings and ethical tips that have helped me to raise my level as a good researcher.

I am thankful with Prof. Dr. Orangel Aguilera Socorro who has been an external adviser, providing important insightful and constructive comments. I would also like to thanks to Dr. Carlos Jaramillo, for his valuable academic support and opportunity for research.

For comments and useful personal and professional advice, many thanks to my friend Thodoris Argyriou. For support and help, cordial thanks to my colleagues and friends from PIMUZ: PD. Dr. Torsten Scheyer, Heike Götzmann, my office-mates Dr. James Neenan, Dr. Richard Hofmann and Dr. Ji Cheng, Dr. Gabriel Aguirre, Heinrich Walter, Prof. Dr. Christian Klug, Alexandra Wegmann, Madlen Stange, Dr. Anna Sanson Barrera, Morgane Brosse, Dr. Christian Kolb, Markus Hebeisen, Rosi Roth, Beat Scheffold, Kristof Veitschegger, Carlo Romano, Marc Leu, Stephan Spiekman, Juan David Carrillo, Ashley Latimer, Amane Tajika, among others.

I would also like to thanks my friends Dr. Analia Forasiepi, Rene Kindlimann, Rodolfo Sánchez, Dr. Eli Amson, Hiram Moreno, as well as my co-author/colleagues, for the enjoyable academic collaboration.

Finally, I would like to thanks my family and my girlfriend Linda Frey, for their invaluable understanding, patience, support, and love throughout my stay in Switzerland.

SUMMARY

The aim of this study is to investigate the Neogene chondrichthyan fauna from Tropical America, based on the documentation of fossil assemblages from different localities of Eastern Pacific and Western Atlantic. Fifteen new assemblages from the Caribbean (Colombia, Panama and Venezuela), Western Atlantic coast (Brazil, in prep.), and Eastern Pacific (Ecuador) were studied, and analyses of their associated paleoenvironments were presented. In Chapter 1, a brief introduction to the evolutionary history of chondrichthyans, a brief survey of the extant and Neogene diversity from Tropical America, and the outline of the project are presented.

The elasmobranch fauna from Eastern Pacific described in Chapter 2, included a taxonomic revision of four assemblages from the Miocene-Pleistocene of Ecuador (Angostura, Onzole, Canoa and Jama formations of Bordón and Manabí basins), resulting in the most diverse paleodiversity known from that region. This study resulted in the description of a new deep water species of frilled shark †*Chlamydoselachus landinii* and other 29 taxa with either deep or shallow water affinities, suggesting paleoenvironments associated with a short platform shelf bordering a deep margin.

Chapter 3 describes the elasmobranch fauna of two members of Chagres Formation, which is the youngest deposit of the Panama Canal Basin in the Caribbean side. A paleodiversity of 30 taxa from both the Rio Indio and Chagres Sandstone Members is reported, including 18 new records for Panama and four for Tropical America. The performed paleobathymetry analysis from the Rio Indio Member suggests a shallow water environment, whereas the assemblage from the Chagres Sandstone Member is dominated by taxa with oceanic affinities, suggesting 200-300 m water depths.

In Chapter 4, 21 taxa of sharks and rays are reported from the Socorro (Middle Miocene), Urumaco (late Miocene) and Codore (late Miocene-early Pliocene) formations located in western Venezuela. These geological units correspond to the “Urumaco sequence”, one of the most significant vertebrate fossiliferous areas from the Neogene of the Northern part of South America. The elasmobranch assemblages include the description of a new carcharhiniform species †*Carcharhinus caquetius* sp. nov., and a detailed taxonomic revision of a high number of well-preserved three-dimensional fossil sawfish rostra (*Pristis*: Rajiformes). We referred to the habitat preferences of the living representatives of fossil chondrichthyan present in the assemblages to reconstruct marine shallow waters and estuarine habitats for this assemblage.

Chapters 5 and 6 concern the taxonomic description of two elasmobranch assemblages of the early Miocene of Colombia and Venezuela, respectively. The assemblage from Colombia comes from the Aquitanian deposits of the Uitpa Formation in the La Guajira Peninsula, providing a unique glimpse into the Caribbean biodiversity at the onset of the Neogene. Thirteen taxa were recovered, and the assemblage was used to conduct a paleoenvironmental and paleobathymetric analysis. The estimation of paleobathymetry suggests that the Uitpa Formation was likely accumulated in a depth of 100 to 200 m. This suggests a rapid increase in relative sea level or basin deepening, providing new insights into the possible causes of marine biota changes in the Cocinetas Basin during the Oligocene/Miocene transition. The assemblage from Venezuela comes from Burdigalian to ?early Langhian deposits of the Cantaure Formation, located in the Paraguaná Peninsula. The fauna is characterized by 39 sharks and ray species, including 15 previously unreported taxa for Venezuela and six for Tropical America, representing one of most diverse assemblages from early Neogene of the Americas. This fauna is characterized by a predominance of

benthopelagic sharks with piscivorous feeding preferences, followed by durophagous/canctitrophic feeders. Habitat preferences, paleobathymetric analyses, and other associated invertebrate and vertebrate assemblages, support the hypothesis that the Cantaure Formation was deposited in a high productivity, insular inner-middle shelf environment.

Preliminary results reveal that throughout the Neogene the chondrichthyan paleodiversity of Tropical America has not remained constant, especially in shark genera, in both Eastern Pacific and Western Atlantic. In spite of the exponential increase in data presented in this thesis, the Neogene record of chondrichthyans is still too incomplete as to provide a solid test of the abiotic variables that affected the evolution of these animals. Our survey not offers a direct biostratigraphic inference to support any of the timings for the definitive closure of the Panama Isthmus. However, clues related with possible consequence of “post-Isthmus” environmental changes can be observed. The analysis reveal regional or complete extirpation, especially during late Miocene-Pliocene, where at least 18 shark/ray taxa (mainly neritic/epipelagic species) from the Eastern Pacific and Western Atlantic were affected.

Keywords: Tropical America, Eastern Pacific, Western Atlantic, Caribbean Sea, Brazil, Colombia, Ecuador, Panamá, Venezuela, Paleodiversity, Paleoenviroments, Central American Seaway, Fossil, Sharks, Rays.

ZUSAMMENFASSUNG

Das Ziel dieser Arbeit ist es die Neogene Chondrichthyes-Fauna aus dem tropischen Amerika, basierend auf der Dokumentierung von Fossilienvergesellschaftungen aus verschiedenen Lokalitäten vom östlichen Pazifik und dem westlichen Atlantik, zu untersuchen. Fünfzehn neue Fossilienvergesellschaftungen von der Karibik (Kolumbien, Panama und Venezuela), der westlichen Atlantikküste (Brasilien, in Vorbereitung), und des östlichen Pazifiks (Ecuador) wurden untersucht und Analysen ihrer assoziierten Palaeoumwelt wurden präsentiert. Eine kurze Einleitung in die Evolutionsgeschichte der Chondrichthyes, eine Übersicht über die heutige und die Neogene Diversität des tropischen Amerikas und ein Umriss über das gesamte Projekt werden in Kapitel 1 dieser Arbeit gegeben.

Die in Kapitel 2 beschriebene Elasmobranchier-Fauna aus dem östlichen Pazifik beinhaltete die taxonomische Revidierung von vier Fossilienvergesellschaftungen aus dem Miozän-Pleistozän von Ecuador (Angostura-, Onzole-, Canoa- und Jama-Formation aus dem Bordón-, respektive Manabí-Becken) und resultierte in der höchsten Palaeodiversität aus dieser Region. Aus dieser Arbeit resultierte die Beschreibung einer neuen Tiefseeart der Kragenhaie †*Chlamydoselachus landinii* und 29 zusätzlichen Taxa mit Affinitäten zu entweder Tiefen- oder Flachwasser. Diese Affinitäten weisen auf ein Kontinentalhang-Environment hin.

In Kapitel 3 wird die Elasmobranchier-Fauna aus zwei Members der Chagres-Formation, welche die jüngste Ablagerung des Panamkanal-Beckens auf der karibischen Seite ist, beschrieben. Die Palaeodiversität bestand aus 30 Taxa vom Rio Indio und dem Chagres Sandstone Member und beinhaltete 18 neue Taxa von Panama und vier neue Taxa aus dem tropischen Amerika. Die Analyse der Palaeobathymetrie von der Rio Indio Member deutete auf ein Flachwasser-Environment hin, während die Fossilien-gemeinschaft aus dem Chagres

Sandstone Member von Taxa mit ozeanischen Affinitäten dominiert wurde und auf eine Wassertiefe von 200 bis 300 Meter hindeutete.

In Kapitel 4 wurde von 21 Hai- und Rochen-Taxa aus der Socorro- (mittleres Miozän), Urumaco- (spätes Miozän) und Codore- (spätes Miozän-frühes Pliozän) Formation aus dem Westen Venezuelas berichtet. Diese geologische Einheit entspricht der "Urumaco Sequenz", welche eine der fossilienreichsten Gebiete des Neogens aus dem nördlichen Südamerika ist. Basierend auf der Elasmobranchier-Fauna wurde eine neue Art der Carcharhiniformes †*Carcharhinus caquetius* sp. nov. beschrieben und die Taxonomie der Sägefische (*Pristis*: Rajiformes) anhand von vielen dreidimensional erhaltenen Rostren detailliert revidiert. Anhand der Habitatspräferenzen von heute noch lebenden Vertretern, die in den Fossilienvergesellschaftungen vorkamen, konnte ein marines Flachwasser und estuarines Habitat rekonstruiert werden.

Kapitel 5 und 6 befassen sich mit der taxonomischen Beschreibung von zwei fossilen Elasmobranchier-Vergesellschaftungen aus dem frühen Miozän von Kolumbien und Venezuela. Das Material aus Kolumbien stammt aus den Aquitanium-Ablagerungen von der Uitpa Formation in der La Guajira Peninsula und gibt einen einzigartigen Einblick in die karibische Biodiversität zu Beginn des Neogens. Dreizehn Taxa wurden gefunden und das Palaeoenvironment und die Palaeobathymetrie wurden anhand der Fossilienvergesellschaftungen analysiert. Die Schätzung der Palaeobathymetrie deutet darauf hin, dass die Uitpa Formation in einer Tiefe von 100 bis 200 Metern akkumuliert wurde. Dies deutet auf einen schnellen Anstieg des relativen Meeresspiegels oder eines Absenkens des Beckens hin, was zu neuen Erkenntnissen über die möglichen Ursachen der Veränderungen in der marinen Biota im Cocinetas-Becken während des Oligozän/Miozän-Übergangs führt. Die Fossilienvergesellschaftungen aus Venezuela stammt aus den Burdigalium bis ?frühen

Langhien-Ablagerungen der Cantaure Formation, welche sich in der Paraguaná Peninsula befindet. Die Fauna ist charakterisiert durch 39 Hai- und Rochenarten, inklusive 15 zuvor unbekannten Taxa aus Venezuela, respektive sechs aus dem tropischen Amerika und repräsentieren somit eine der diversesten Fossilienvergesellschaftungen aus dem frühen Neogen von ganz Amerika. Die Fauna ist charakterisiert durch die Dominanz von benthopelagischen Haien mit piscivoren Nahrungspräferenzen, gefolgt von durophagen/cancritrophen Haiarten. Habitatspräferenzen, palaeobathymetrische Analysen und assoziierte Invertebraten- und Vertebraten-Vergesellschaftungen unterstützen die Hypothese, dass die Cantaure Formation in einem hochproduktiven, Inselähnlichem Environment abgelagert wurde.

Die vorläufigen Resultate dieser Arbeit zeigen, dass die Palaeodiversität der Chondrichthyes des tropischen Amerikas, besonders die Haigattungen im östlichen Pazifik und westlichen Atlantik, während des Neogens nicht konstant geblieben sind. Trotz des exponentiellen Anstiegs der Daten, die in dieser Arbeit präsentiert werden, ist der Fossilbericht der Neogenen Knorpelfische immer noch zu lückenhaft um einen solide Überprüfung der abiotischen Faktoren, die die Evolution dieser Tiere beeinflusst hat, zu liefern. Aus unserer Untersuchung lässt sich kein direkter biostratigraphischer Rückschluss ziehen um den Zeitpunkt der definitiven Schliessung des Panama Isthmus zu bestätigen. Jedoch konnten mögliche Konsequenzen durch die "post-Isthmus" Umweltveränderungen beobachtet werden. Die Analyse weist auf ein regionales oder vollständiges Aussterben von Taxa im tropischen Amerika hin: Besonders während des späten Miocene-Pliocene verschwanden mindestens 18 Hai- und Rochen-Taxa (hauptsächlich neritische/epipelagische Arten) aus dem östlichen Pazifik und dem westlichen Atlantik.

Schlüsselworte: Tropisches Amerika, östlicher Pazifik, westlicher Atlantik, karibisches Meer, Brasilien, Kolumbien, Ecuador, Panama, Venezuela, Palaeodiversität, Palaeoenvironment, Zentralamerkanischer Seeweg, Fossilien, Haie, Rochen.

AUTHOR CONTRIBUTIONS

Excluding chapters one (Introduction) and seven (Conclusion and Future Perspectives), the remaining chapters and appendices of this thesis have been published in peer-reviewed journals.

CHAPTER 1

Author: Carrillo-Briceño J.D.

CHAPTER 2

Fossil Chondrichthyes from the central eastern Pacific Ocean and their paleoceanographic significance.

Publication: 2014, Journal of South American Earth Sciences, 51: p. 76–70.

Authors: Carrillo-Briceño J.D., Aguilera O.A., and Rodríguez F.

Contributions: designed research (JDCB, OAA), data collection (JDCB, OAA), data analysis (JDCB, OAA), photography (OAA, FR), figures and tables (JDCB), wrote paper (JDCB, OAA).

CHAPTER 3

A new Late Miocene chondrichthyan assemblage from the Chagres Formation, Panama.

Publication: 2015, Journal of South American Earth Sciences, 60: p. 56–70.

Authors: Carrillo-Briceño J.D., De Gracia C., Pimiento C., Aguilera O.A., Kindlimann R., Santamarina P., and Jaramillo C.

Contributions: designed research (JDCB, CD, CJ), data collection (JDCB, CD, OAA), data analysis (JDCB, CD, RK, CJ), photography (JDCB, CD), figures and tables (JDCB, CD), wrote paper (JDCB, CD), editing of the manuscript (JDCB, CD, CJ, CP).

CHAPTER 4

Sawfishes and Other Elasmobranch Assemblages from the Mio-Pliocene of the South Caribbean (Urumaco Sequence, Northwestern Venezuela).

Publication: 2015, PLoS ONE, 10(10): e0139230.

Authors: Carrillo-Briceño J.D., Maxwell E., Aguilera O.A., Sánchez R., and Sánchez-Villagra M.R.

Contributions: designed research (JDCB, EM), data collection (JDCB, EM, OAA, RS), data analysis (JDCB, EM), photography (JDCB), figures and tables (JDCB), wrote paper (JDCB, EM, MRSV), editing of the manuscript (JDCB, EM, OAA, MRSV).

CHAPTER 5

A new early Miocene (Aquitania) Elasmobranchii assemblage from the La Guajira Peninsula, Colombia.

Publication: 2016, Ameghiniana 53: p. 77–99.

Authors: Carrillo-Briceño J.D., Argyriou T., Zapata V., Kindlimann R., and Jaramillo C.

Contributions: designed research (JDCB, CJ), data collection (JDCB, VZ), data analysis (JDCB, TA; CJ, RK), photography (JDCB, VZ), figures and tables (JDCB, TA), wrote paper (JDCB, TA, CJ), editing of the manuscript (JDCB, TA, CJ).

CHAPTER 6

An Early Neogene Elasmobranch fauna from the southern Caribbean (Western Venezuela).

Publication: 2016, Paleontologia Electronica.

Authors: Carrillo-Briceño J.D., Aguilera O.A., De Gracia C., Aguirre-Fernández G., Kindlimann R., and Sánchez-Villagra M.R.

Contributions: designed research (JDCB, OAA), data collection (JDCB, OAA, GAF, MESV), data analysis (JDCB, CD, RK), photography (JDCB), figures and tables (JDCB), wrote paper (JDCB), editing of the manuscript (JDCB, GAF, MRSV).

APPENDIX I

Late Neogene elasmobranch fauna from the Coquimbo Formation, Chile.

Publication: 2015, Revista Brasileira de Paleontologia, 18(2): p. 261–272.

Authors: Staig F., Hernández S., López P., Villafaña J.A., Varas C., Soto L.P., and Carrillo-Briceño, J.D.

Contributions: designed research (FS, PS, JAV), data collection (FS, PS, JAV, PLS, JDCB), data analysis (FS, JAV, PLS, JDCB), photography (FS, JAV), figures and tables (JAV), wrote paper (FS, PS, JAV, JDCB), editing of the manuscript (FS, PS, JAV, JDCB).

APPENDIX II

The Path towards Endangered Species: Prehistoric Fisheries in Southeastern Brazil.
Publication: 2016, PLoS ONE, 11(6): e0154476.

Authors: Lopes M.S., Pereira-Bertucci T.C., Rapagnã L., Tubino R.A., Monteiro-Neto C., Tomas ARG, Tenório M.C., Lima T., Souza R., Carrillo-Briceño J.D., Haimovici M., Macario K., Carvalho C., Aguilera-Socorro O.A.

Contributions: designed research (MSL, TCPB, OAAS), data collection (MSL, TCPB, OAAS), data analysis (MSL, TCPB, LR, RAT, CMN, ARG, MCT, TL, RS, JDCB, MH, KM, CC, OAAS), photography (MSL, TCPB, OAAS, JDCB), figures and tables (MSL, TCPB, OAAS, JDCB), wrote paper (MSL, TCPB, OAAS), editing of the manuscript (MSL, TCPB, LR, RAT, CMN, ARG, MCT, TL, RS, JDCB, MH, KM, CC, OAAS).

CHAPTER 1

Introduction



Tiger shark *Galeocerdo cuvier*. Artwork: Jorge González, 2015

1. Introduction

1.1 A brief introduction to the evolutionary history and extant diversity of chondrichthyans

Living chondrichthyan diversity is composed of thirteen ‘orders’ (Figure 1), separated in two big groups, the Elasmobranchii, that includes sharks and rays, and the Holocephali, represented exclusively by the chimaeras (Compagno et al., 2005a; Klimley, 2013). With a worldwide distribution, in both freshwater and marine environments, this diversity has been referred to 503 species of sharks, 699 species of rays and 49 species of chimaeras (Klimley, 2013). In the future more species are likely to be discovered as the deep waters of the world’s oceans become better explored.




Figure 1. Diversity and phylogenetic hypothesis of living chondrichthyans. Based on Naylor et al. (2012), Aschliman et al. (2012), and Klimley (2013).

The chondrichthyan evolutionary history may have undergone two major episodes of adaptive radiation (Figure 2), the first during late Silurian and Devonian (Brazeau and Friedman, 2015), resulting in sharks that broadly resemble modern genera but with a more plesiomorphic skeleton (Carroll, 1988). By the early Carboniferous, chondrichthyans underwent a second major radiation process, where many ‘families’ and major clades appeared, including holocephalans (Jong, 2011; Brazeau and Friedman, 2015). This radiation continued throughout the Mesozoic; by the end of this era most of the modern chondrichthyan ‘orders’ had appeared (e.g. Jong, 2011; Cappetta, 2012).




Figure 1. Stratigraphic range of major chondrichthyan clades (‘orders’). Based on Stahl (1999), Ginter (2010), and Cappetta (2012).

The evolution of chondrichthyans is documented paleontologically principally by fossil vertebrae, teeth, spines and dermal denticles, besides some exceptionally preserved specimens consisting of cranial elements or the whole body. The earliest fossil evidence of chondrichthyans is in the form of isolated dermal denticles, specimens found in rock deposits of the late Ordovician period (Figure 1), with an age of 455 million years ago (Ma.) (Maisey, 1996; Cuny, 2013; Klimley, 2013). According to Jong (2011), the fact these dermal denticles occur in this time without associated fossilized teeth, suggest that these early “protosharks” may not necessarily have had teeth. First shark teeth appear in the fossil record at the beginning of the Devonian period (418 Ma.), and they were clearly derived from dermal denticles, which had become concentrated by this time along the margin of the jaw (Klimley, 2013, p. 15).

1.2 The Neogene marine chondrichthyan faunas from Tropical America

The “Neogene” is a geological period which following the International Commission on Stratigraphy (ICS) began about 23 Ma. and ended 2.58 Ma.; it is sub-divided into two epochs, the Miocene and Pliocene (Figure 2). This period was a time when the world climate was warmer, producing for example sea level changes (Zachos et al., 2001). At the time, the America landmass had a similar geographic position to the recent one (Santos et al., 2004), but the northern and southern continental masses were separated by a connection between the Pacific and Atlantic Oceans called the “Central America Seaway” (Montes et al., 2012; Coates and Stallard, 2013). At the end of the Neogene, large scale geological processes, including the closure of the Central American Seaway and the rise of the Panamanian Isthmus (Figure 2), were completed (Woodburne, 2010; Montes et al., 2012, 2015; Coates and Stallard, 2013). This resulted in a massive terrestrial faunal exchange between North America

and South America, the Great American Biotic Interchange (Woodburne, 2010, Carrillo et al., 2015). The land bridge played also the role of a marine barrier, inducing environmental changes on the Pacific and Atlantic oceans, creating contrasting ecosystems through differential extinction and diversification, especially in the Tropical America region (Leigh et al., 2014).




Figure 2. Neogene chronostratigraphy (A) and the rise of the Isthmus of Panama (B); based on International Chronostratigraphic Chart 2015 (<http://www.stratigraphy.org/index.php/ics-chart-timescale>), and modified after Carrillo-Briceño (2015) and based on O'Dea et al. (2007b); respectively.

Tropical America (Neotropics) refers here to the geographical continental and marine areas of the western hemisphere located between the Tropic of Cancer (23° 27' N) and the Tropic of Capricorn (23° 27' S) (Figure 3). This vast marine region is environmentally

heterogenic and ecologically complex, supporting a high marine diversity (e.g. Miloslavich et al., 2011). Marine areas of Tropical America have a well-known chondrichthyan diversity in both Eastern Pacific and Western Atlantic zones, which represents a valuable commercial resource (Carpenter, 2002; Compagno et al., 2005b; Fowler et al., 2005). The known Tropical America's extant chondrichthyan diversity is characterized by a total of 276 species of chimaeras, sharks and rays (Eastern Pacific: 145 spp.; Western Atlantic 181 spp.), representing the highest for the group in the Americas (Figure 3). Comparing the Eastern Pacific and Western Atlantic, a clear differential pattern emerge (Figure 4). This pattern was surely influenced by the geological and oceanographic events that occurred in the region after the final closure of the Isthmus of Panama (e.g. O'Dea et al., 2007a; Coates, and Stallard, 2013; Leigh et al., 2014).




Figure 3. Tropical America and chondrichthyan diversity of the Americas. Total number of species include chimaeras, sharks and rays. From Carrillo-Briceño and Sánchez-Villagra (in prep).




Figure 4. Chondrichthyan diversity of Tropical America. A. Total number of species by ‘orders’. B. Shared/Unshared genera and species in both the Eastern Pacific and Western Atlantic. From Carrillo-Briceño and Sánchez-Villagra (in prep).

The origin and distribution pattern of this Tropical America chondrichthyan diversity is still somewhat uncertain, in part due to the fragmentary and incompletely studied Neogene fossil record. Some previous works have made contributions to document fossil chondrichthyan palaeodiversity from Tropical America, with described assemblages from: Barbados (Casier, 1958, 1966), Brazil (Santos and Travassos, 1960; Santos and Salgado, 1971; Reis, 2005; Costa et al., 2009), Costa Rica (Laurito, 1999); Cuba (Iturralde-Vinent et al., 1996; MacPhee et al., 2003), Southern Mexico (González-Rodríguez et al., 2013), Panama

(Pimiento et al., 2013a, 2013b), Peru (Kindlimann, 1990; Alván et al., 2007), Trinidad (Leriche, 1938), The Grenadines (Portell et al., 2008) and Venezuela (Leriche, 1938; Sánchez-Villagra et al., 2000; Aguilera, 2010; Aguilera and Rodrigues de Aguilera, 2004; Aguilera and Lundberg, 2010).

1.3 Chondrichthyan dental characters and the used terminology

Teeth, dermal denticles, vertebrae, fin spines, rostra, among other hard elements, are common in the fossil record of chondrichthyans. Due to the fact that selachians have a polyphyodont dentition, in which dental elements are replaced throughout life (Klimley, 2013; Pough et al., 2013), teeth are usually the most common chondrichthyan remains found in Neogene rocks (e.g. Cappetta, 2012), and their role as diagnostic elements allows for accurate taxonomic comparisons. Shark and ray teeth chemical composition is similar to that of dermal denticles, consisting mainly of mineralized tissue, dentine, interspersed with blood vessel and surrounded by highly mineralized enameloid on the crown of the tooth (Klimley, 2013). Tooth development and replacement in sharks and rays have been the subject of many studies (e.g. Cappetta, 2012, p. 10), and the study of their morphological changes in geological time provide the basis understanding of the evolution of these diverse fish groups (Underwood et al., 2015).

There are diagnostic characters for identifications of an isolated tooth based on its morphology, including size, shape and thickness of the crown and root (Cappetta, 2012). Taking under consideration dental characters in chondrichthyans can serve for generic and specific identifications. Some exceptions where taxonomic problems cannot be solved include a poor preservation of the fossil specimens and/or the biases or uncertainties resulting from

ontogenetic variation or jaw position. In the studied assemblages from Tropical America, the fossil samples are represented mainly by isolated teeth. Some other fossil elements, such as sawfish rostra (including rostral denticles), vertebrae and dermal denticles were also found in a minor proportion. The dental terminology (including measurements, heterodonty, adaptative dental types, root vascularization, jaw position and tooth histology) used to describe these assemblages from tropical America, follows Cappetta (2012).

1.4 Aims and overview

The extant Tropical America's chondrichthyan diversity (Figures 3-4) must have greatly affected by environmental changes occurring in both Eastern Pacific and Western Atlantic regions after the definitive closure of the Panama Isthmus at the end of the Neogene (e.g. O'Dea et al., 2007a; Coates and Stallard, 2013). This land-bridge played an important role as a marine barrier, probably inducing to a regional or complete extirpation and or extinctions of many chondrichthyan taxa, as well as the origin of species adapted to the new environmental conditions. To understand the current diversity, two important questions are postulated: a) what was the chondrichthyan paleodiversity in Tropical America before and after the rise of the Panama Isthmus? b) Did the new marine post-isthmus environmental differences in both oceans played an important role for chondrichthyan divergence and speciation? Despite all previous efforts in the study of the fossil chondrichthyans from Tropical America, still the lack of a solid paleontological documentation does not allow for an understanding of the impact of these regional ecological and biogeographic changes (e.g. O'Dea et al., 2007a; Coates and Stallard, 2013; Montes et al., 2015).

The aim of this study is to investigate the Neogene chondrichthyan fauna from Tropical America, based on the documentation of fossil assemblages with shallow and deep-water affinities from both sides of the Panamanian Isthmus. We studied fifteen new chondrichthyan assemblages from the Caribbean (Colombia, Panama and Venezuela), Western Atlantic coast (Brazil, in prep.), and Eastern Pacific (Ecuador), presenting a comprehensive interpretation of their associated paleoenvironments.

In chapter 2, a taxonomic revision of four elasmobranch assemblages from the Miocene-Pleistocene of Ecuador (Angostura, Onzole, Canoa and Jama formations of Bordón and Manabí basins) is provided, resulting in the most diverse paleodiversity known from the Eastern Pacific coast of Tropical America. This study reveals the presence of at least 30 taxa, including species with deep and shallow waters affinities which suggest paleoenvironments associated with a short platform shelf bordering a deep margin. The new deep water species †*Chlamydoselachus landinii* sp. nov. was described.

Chapter 3 includes the description of two new assemblages from both the Rio Indio and Chagres Sandstone Members of the Chagres Formation. This unit corresponds with youngest deposits of the Panama Canal Basin, and their paleodiversity includes 18 new fossil records for Panama and four for Tropical America. Herein it is presented a comprehensive interpretation of the paleoenvironments of assemblages of the Chagres Formation, for which a paleobathymetry analysis based on the modern water depth preference of extant chondrichthyan taxa was performed. The assemblage from the Rio Indio Member is characterized by taxa with neritic affinities, suggesting depths <100 m, whereas the assemblage from the Chagres Sandstone Member is dominated by taxa with oceanic affinities, suggesting 200-300 m water depths.

Chapter 4 is concerned with one of the most significant vertebrate fossiliferous areas from the Neogene of the Northern part of South America “The Urumaco sequence”. This stratigraphic sequence is located in western Venezuela, and preserves a wide range of fossil vertebrate associated to different paleoenvironments including terrestrial, riverine, lacustrine and marine facies. Herein, 21 taxa of sharks and rays were referred from the Socorro (Middle Miocene), Urumaco (late Miocene) and Codore (late Miocene-early Pliocene) formations. The assemblages include the description of a new carcharhiniform species †*Carcharhinus caquetius* sp. nov. The high number of well-preserved 3D fossil sawfish rostra (*Pristis*: Rajiformes), allowed a detailed taxonomic revision of these specimens, referring those in the context of the global Miocene record of *Pristis* as well as extant species. Using the habitat preference of the living representatives, fossil chondrichthyan assemblages from the Urumaco sequence are evidence for marine shallow waters and estuarine habitats.

Chapter 5 comprises the first taxonomic description for a chondrichthyan assemblage from Colombia. This elasmobranch assemblage is coming from the earliest Miocene (Aquitanian) deposits of the Uitpa Formation in the La Guajira Peninsula, providing a unique glimpse into the Caribbean biodiversity at the onset of the Neogene. Thirteen taxa were recovered, and the assemblage was used to conduct a paleoenvironmental and paleobathymetric analysis. The estimation of paleobathymetry suggests that the Uitpa Formation was likely accumulated in a depth of 100 to 200 m. This suggests a rapid increase in relative sea level or basin deepening, providing new insights into the possible causes of marine biota changes in the Cocinetas Basin during the Oligocene/Miocene transition.

Chapter 6 continues the description of assemblages from southern Caribbean, this time with a focus in the Cantaure Formation (Burdigalian to ?early Langhian), Western Venezuela. Here a taxonomic study of the elasmobranch fauna and their significance for

paleoenvironmental and paleoecological inferences is presented. The assemblage is represented by 39 shark and ray species, including 15 previously unreported taxa for Venezuela and six for Tropical America, Cantaure being thus one of most diverse assemblages from early Neogene of the Americas. The fauna is characterized by a predominance of benthopelagic sharks with piscivorous feeding preferences, followed by durophagous/canctitrophic feeders. Habitat preferences, paleobathymetric analyses, and other associated invertebrate and vertebrate assemblages, support the hypothesis that the Cantaure Formation was deposited in a high productivity, insular inner-middle shelf environment.

References

- Aguilera, O.A. 2010. Peces Fósiles del Caribe de Venezuela. Gorham Printing, Washington.
- Aguilera, O.A., and Rodrigues de Aguilera, D. 2004. Giant-toothed White Sharks and Wide-toothed Mako (Lamnidae) from the Venezuela Neogene: Their Role in the Caribbean, Shallow-water Fish Assemblage. *Caribbean Journal of Science*, 40(3):368-382.
- Aguilera, O.A., and Lundberg, J.G. 2010. Venezuelan Caribbean and Orinocoan Neogene fish, p. 129-152. In Sánchez-Villagra, M.R., Aguilera, O.A., and Carlini, F. (eds.), *Uruguay and Venezuelan Paleontology*. Indiana Press University, Bloomington.
- Alván, A., Apolín, J., and Chacaltana, C. 2006. Los dientes de seláceos (Chondrichthyes) y su aplicación estratigráfica en la lomas de Ullujaya (Ica-Perú). In Perú, S.G.P. (ed.), *XIII Congreso Peruano de Geología*, Lima, 17-20 octubre, 2006, pp. 595-598.
- Aschliman, N.C., Claeson, K.M., and McEachran, J.D. 2012. Phylogeny of Batoidea, p. 57-95. In Carrier, J.C., Musick, J.A., and Heithaus, M.R. (eds.), *Biology of Sharks and Their Relatives*. CRC Press, Taylor & Francis Group, Boca Raton, Florida.

- Brazeau, M.D., and Friedman, M. 2015. The origin and early phylogenetic history of jawed vertebrates. *Nature*, 520(7548):490-497.
- Cappetta, H. 2012. Chondrichthyes. Mesozoic and Cenozoic Elasmobranchii: Teeth. *Handbook of Paleoichthyology*, 3E. Verlag Dr. Friedrich Pfeil, Munich.
- Carpenter, K.E. 2002. The living marine resources of the Western Central Atlantic, Volume 1: Introduction, molluscs, crustaceans, hagfishes, sharks, batoid fishes and chimaeras. *FAO Species Identification Guide for Fishery Purposes and American Society of Ichthyologists and Herpetologists Special Publication No. 5*, Rome.
- Carrillo-Briceño, J.D. 2015. Bestias Prehistóricas de Venezuela “Colosos de la Edad de Hielo”. Río Verde, Caracas.
- Carrillo, J.D., Forasiepi, A., Jaramillo, C., and Sánchez-Villagra, M.R. 2015. Neotropical mammal diversity and the Great American Biotic Interchange: spatial and temporal variation in South America's fossil record. *Frontiers in Genetics*, 5.
- Carroll, R.L. 1988. *Vertebrate Paleontology and Evolution*. W.H. Freeman and Company, New York.
- Casier, E. 1958. Contribution à l'étude des poissons fossiles des Antilles. *Mémoire Suisse de Paléontologie*, 74:1-95.
- Casier, E. 1966. Sur la faune ichthyologique de la Formation de Bissex Hill et de la Série océanique, de l'Ile de la Barbade, et sur l'âge de ces formations. *Eclogae Geologicae Helvetiae*, 59:493-516.
- Coates, A.G., and Stallard, R.F. 2013. How old is the Isthmus of Panama? . *Bulletin of Marine Sciences* 89(4):801–813.

- Compagno, L.J.V., Dando, M., and Fowler, S. 2005a. *Sharks of the World*. Princeton University Press, Princeton.
- Compagno, L.J.V., Didier, D.A., and Burgess, G.H. 2005b. Classification of Chondrichthyan Fish, p. 4-11. In Fowler, S.L. et al. (eds.), *Sharks, Rays and Chimaeras: The Status of the Chondrichthyan Fishes*. IUCN/ SSC Shark Specialist Group. IUCN, Gland, Switzerland and Cambridge, UK.
- Costa, S.A.F., Richter, M., Toledo, P.M., and Moraes-Santos, H.M. 2009. Shark teeth from Pirabas formation (Lower Miocene), northeastern Amazonia, Brazil. *Boletim do Museu Paraense Emílio Goeldi, Ciencias Naturais* 4:221-230.
- Cuny, G. 2013. *Requins -De la préhistoire à nos jours*. Belin, Paris.
- Fowler, S.L., Cavanagh, R.D., Camhi, M., Burgess, G.H., Cailliet, G.M., Fordham, S.V., and A, S.C. 2005. *Sharks, Rays and Chimaeras: The Status of the Chondrichthyan Fishes*. IUCN/SSC Shark Specialist Group, Gland, Switzerland and Cambridge, UK.
- Ginter, M., Hampe, O., and Duffin, C.J. 2010. Chondrichthyes, Paleozoic Elasmobranchii: Teeth. *Handbook of Paleoichthyology*, 3D. Verlag Dr. Friedrich Pfeil, Munich.
- González-Rodríguez, K.A., Espinosa-Arrubarrena, L., and González-Barba, G. 2013. An overview of the Mexican fossil fish record, p. 9-34. In Arratia, G., Schultze, H.P., and Wilson, M.V.H. (eds.), *Mesozoic Fishes 5 – Global Diversity and Evolution*. Verlag Dr. Friedrich Pfeil, München, Germany.
- Iturralde-Vinent, M.A., Hubbell, G., and Rojas, R. 1996. Catalogue of Cuban fossil Elasmobranchii (Paleocene to Pliocene) and paleogeographic implications of their

- lower to middle Miocene occurrence. *Boletín de la Sociedad Jamaicana de Geología* 31:7-21.
- Jong, J.A. 2011. *The rise of the fishes. 500 million years of evolution*, The Johns Hopkins University Press, Baltimore.
- Kindlimann, R. 1990. Selacios del Terciario Tardío de Sacaco, Departamento de Arequipa. *Boletín de Lima* 69:91-95.
- Klimley, P.A. 2013. *The biology of sharks and rays*. The University of Chicago Press, Chicago.
- Laurito, C.A. 1999. Los seláceos fósiles de la localidad de Alto Guayacán (y otros ictiolitos asociados), Mioceno superior-Plioceno inferior de la Formación Uscari, provincia de Limón, Costa Rica. Laurito, C. (Ed.), San José.
- Leigh, E.G., O'Dea, A., and Vermeij, G.J. 2014. Historical biogeography of the Isthmus of Panama. *Biological Reviews*, 89(1):148-172.
- Leriche, M. 1938. Contribution à L'étude des Poissons fossils des pays riverains de la Méditerranée américaine, Venezuela, Trinité, Antiles, Mexique. *Mémoires de la Société paléontologique du Suisse* 61:1-52.
- MacPhee, R.D.E., Iturralde-Vinent, M.A., and Gaffney, E.S. 2003. Domo de Zaza, an Early Miocene vertebrate locality in South-Central Cuba, with notes on the tectonic evolution of Puerto Rico and the Mona Passage. *American Museum Novitates*, 3394:1-42.
- Miloslavich, P., Klein, E., Díaz, J.M., Hernández, C.E., Bigatti, G., Campos, L., Artigas, F., Castillo, J., Penchaszadeh, P.E., Neill, P.E., Carranza, A., Retana, M.V., Díaz de


- Astarloa, J.M., Lewis, M., Yorio, P., Piriz, M.L., Rodríguez, D., Yoneshigue-Valentin, Y., Gamboa, L., and Martín, A. 2011. Marine Biodiversity in the Atlantic and Pacific Coasts of South America: Knowledge and Gaps. *PLoS ONE*, 6(1):e14631.
- Montes, C., Cardona, A., McFadden, R., Morón, S.E., Silva, C.A., Restrepo-Moreno, S., Ramírez, D.A., Hoyos, N., Wilson, J., and Farris, D. 2012. Evidence for middle Eocene and younger land emergence in central Panama: implications for Isthmus closure. *Geological Society of America Bulletin*, 124(5-6):780-799.
- Montes, C., Cardona, A., Jaramillo, C., Pardo, A., Silva, J.C., Valencia, V., Ayala, C., Pérez-Angel, L.C., Rodríguez-Parra, L.A., Ramirez, V., and Niño, H. 2015. Middle Miocene closure of the Central American Seaway. *Science*, 348(6231):226-229.
- Maisey, J.G. 1996. *Discovering fossil fishes*. Henry Holt and Company, New York.
- Naylor, G.J.P., Caira, J.N., Jensen, K., Rosana, K.A.M., Straube, N., and Lakner, C. 2012. Elasmobranch Phylogeny: A Mitochondrial Estimate Based on 595 Species, p. 31-56. In Carrier, J.C., Musick, J.A., and Heithaus, M.R. (eds.), *Biology of Sharks and Their Relatives*. CRC Press, Taylor & Francis Group, Boca Raton, Florida.
- O'Dea, A., Jackson, J.B.C., Fortunato, H., Smith, J.T., D'Croz, L., Johnson, K.G., and Todd, J.A. 2007a. Environmental change preceded Caribbean extinction by 2 million years. *Proceedings of the National Academy of Sciences*, 104(13):5501-5506.
- O'Dea, A., Rodríguez, F., De Gracia, C., and Coates, A.G. 2007b. Patrimonio paleontológico: La paleontología marina en el Istmo de Panamá. *Canto Rodado: Revista especializada en patrimonio*, 2:149-179.

- Pimiento, C., González-Barba, G., Hendy, A.J.W., Jaramillo, C., MacFadden, B.J., Montes, C., Suarez, S.C., and Shippritt, M. 2013a. Early Miocene chondrichthyans from the Culebra Formation, Panama: A window into marine vertebrate faunas before closure the Central American Seaway. *Journal of South American Earth Sciences*, 42:159-170.
- Pimiento, C., González-Barba, G., Ehret, D.J., Hendy, A.J., MacFadden, B.J., and Jaramillo, C. 2013. Sharks and rays (Chondrichthyes, Elasmobranchii) from the late Miocene Gatun Formation of Panama. *Journal of Paleontology*, 87(5):755-774.
- Portell, R.W., Hubbell, G., Donovan, S.K., Green, J.L., Harper, D.A., and Pickerill, R. 2008. Miocene sharks in the Kendeace and Grand Bay formations of Carriacou, The Grenadines, Lesser Antilles. *Caribbean Journal of Science*, 44(3):279-286.
- Pough, F.H., Janis, C.M., and Heiser, J.B. 2013. *Vertebrate Life*. Pearson, Glenview.
- Reis, M.A.F.d. 2005. Chondrichthyan fauna from the Pirabas Formation, Miocene of northern Brazil, with comments on paleobiogeography. *Anuário do Instituto de Geociências*, 28(2):31-58.
- Sánchez-Villagra, M.R., Burnham, R.J., Campbell, D.C., Feldmann, R.M., Gaffney, E.S., Kay, R.F., Lozsán, R., Purdy, R., and Thewissen, J.G.M. 2000. A New Near-Shore Marine Fauna and Flora from the Early Neogene of Northwestern Venezuela. *Journal of Paleontology*, 74(5):957-968.
- Santos, R.S., and Travassos, H. 1960. Contribuição à Paleontologia do estado do Pará. Peixes fósseis da Formação Pirabas. Monografia, 16. Ministério da Agricultura, Departamento Nacional da Produção Mineral, Divisao de Geologia e Mineralogia Rio de Janeiro.

- Santos, R.S., and Salgado, M.S. 1971. Contribuição à paleontologia do estado do Pará. Novos restos de peixes da Formação Pirabas. Boletim do Museu Paraense Emílio Goeldi, 16:1-13.
- Smith, A.G., Smith, D.G., and Funnell, B.M. 2004. Atlas of Mesozoic and Cenozoic coastlines. Cambridge University Press, Cambridge.
- Stahl, B.J. 1999. Chondrichthyes III: Holocephali. Handbook of Paleoichthyology, 4. Verlag Dr. Friedrich Pfeil, Munich.
- Underwood, C.J., Johanson, Z., Welten, M., Metscher, B., Rasch, L.J., Fraser, G.J., and Smith, M.M. 2015. Development and Evolution of Dentition Pattern and Tooth Order in the Skates And Rays (Batoidea; Chondrichthyes). PLoS ONE, 10(4):e0122553.
- Woodburne, M.O. 2010. The Great American Biotic Interchange: Dispersals, Tectonics, Climate, Sea Level and Holding Pens. Journal of Mammalian Evolution, 17(4):245-264.
- Zachos, J.C., Shackleton, N.J., Revenaugh, J.S., Pälike, H., and Flower, B.P. 2001. Climate Response to Orbital Forcing Across the Oligocene-Miocene Boundary. Science, 292(5515):274-278.

CHAPTER 2

Fossil Chondrichthyes from the central Eastern Pacific Ocean and their paleoceanographic significance



Frilled shark †*Chlamydoselachus landinii*



Fossil Chondrichthyes from the central eastern Pacific Ocean and their paleoceanographic significance



Jorge D. Carrillo-Briceño^{a,*}, Orangel A. Aguilera^b, Félix Rodríguez^c

^a Paleontological Institute and Museum, University of Zurich, Karl-Schmid-Strasse 4, CH-8006 Zurich, Switzerland

^b Universidade Federal Fluminense, Instituto de Biologia, Departamento de Biologia Marinha, Programa de Pós-graduação em Biologia Marinha, Campus do Valonguinho, Outeiro São João Batista, s/n°, CEP: 24020-141 Niterói, Rio de Janeiro, Brazil

^c Smithsonian Tropical Research Institute, PO Box 0843-03092, Balboa, Panama

ARTICLE INFO

Article history:

Received 8 November 2013

Accepted 2 January 2014

Keywords:

Neogene

Tropical America

Ecuador

Sharks

Rays

Paleoenvironments

ABSTRACT

The study of the chondrichthyan fauna from the Angostura, Onzole, Canoa and Jama formations, in the Neogene of Bordon and Manabí basins, Ecuador, reveals the presence of 30 taxa, including the deep water shark *Chlamydoselachus landinii* sp. nov. The assemblages are dominated by tropical shallow and deep water chondrichthyans, suggesting paleoenvironments associated with a short platform shelf bordering a deep margin. These assemblages are the most diverse shark and ray association known from the Tropical Central Eastern Pacific Ocean in the South American coastal basin, and provide new information on the paleoecology and paleodiversity of Neogene chondrichthyans.

© 2014 Elsevier Ltd. All rights reserved.

1. Introduction

Fossil chondrichthyan assemblages from the eastern Pacific Ocean are still poorly known in spite of several publications on the subject, including: Longbottom (1979) and Aguilera et al. (2011) on Ecuador; De Muizon and Devries (1985), Kindlimann (1990), Apolin et al. (2004) and Avalán et al. (2007) on Perú; Long (1993a), Suárez and Marquardt (2003), Suárez et al. (2006) and Carrillo-Briceño et al. (2013) on Chile; Laurito (2004), and Laurito and Valerio (2008) on Central America, and Long (1993b), González-Barba and Thies (2000), and Boessenecker (2011) on North America. These previous contributions are not uniform in terms of sampling and geochronological control, preventing comparative quantitative analyses among faunas. However, the faunal assemblages along the tropical eastern Pacific coast have been the subject of studies about faunal distribution in Tropical America and the proto-Caribbean area (Iturralde-Vinent and MacPhee, 1999; Landini et al., 2002a, 2002b; Aguilera et al., 2011; Carnevale et al., 2011).

In terms of regional relevance, the central eastern Pacific Ocean represents a geographic and oceanographic complex influenced by geologic and tectonic events along the Panamanian Atlantic–Pacific

seaway (Hoernle et al., 2002; Pindell et al., 2005), related to the uplift of the Isthmus of Panama (Coates and Obando, 1996; O'Dea et al., 2007; Coates and Stallard, 2013), the Andean uplift (Garzone et al., 2008; Martinoid et al., 2010), the Ecuadorian subduction, the submarine trench system (Lonsdale, 1978; Ratzov et al., 2010) and the oceanographic dynamic (Fiedler and Talley, 2006) derived in successions of neritic to depth sea paleoenvironment changes. Longbottom (1979) described the early chondrichthyan fauna assemblage from the Miocene of the Onzole Formation and Aguilera et al. (2011) reported the presence of additional Miocene–Pleistocene taxa from the Angostura, Onzole and Jama formations from Ecuador. Teleostean records from the Neogene basin in Ecuador are known from otoliths reported from the Onzole and Canoa formations (Landini et al., 2002a, 2002b; Aguilera et al., 2011; Carnevale et al., 2011).

In this work we provide a taxonomic revision of the chondrichthyan fauna from Ecuador, resulting in an increased fossil record for the region and a better understanding of the Neogene shark and ray associations in Tropical America, especially along the eastern Pacific coast of South America.

2. Methods


The chondrichthyan assemblages (Table 1) were collected from nine localities (Fig. 1) of four Neogene geological units in Ecuador, including: Cayapas River [Telembí (0° 58' 57" N, 78° 51' 42" W)],

* Corresponding author.

E-mail addresses: jorge.carrillo@pim.uzh.ch, jorgedcb100@gmail.com (J. D. Carrillo-Briceño), orangel.aguilera@gmail.com (O.A. Aguilera), RodriguezF@si.edu (F. Rodríguez).

CHAPTER 3

A new Late Miocene chondrichthyan assemblage from the Chagres Formation, Panama



Cookiecutter shark *Isistius* sp.



Contents lists available at ScienceDirect

Journal of South American Earth Sciences

journal homepage: www.elsevier.com/locate/jsames

A new Late Miocene chondrichthyan assemblage from the Chagres Formation, Panama



Jorge D. Carrillo-Briceño^a, Carlos De Gracia^{b,*}, Catalina Pimiento^{b, c, d},
Orangel A. Aguilera^e, René Kindlimann^a, Patricio Santamarina^f, Carlos Jaramillo^b

^a Paleontological Institute and Museum, University of Zurich, Karl-Schmid-Strasse 4, CH-8006, Switzerland

^b Center for Tropical Paleocology and Archaeology, Smithsonian Tropical Research Institute, Balboa, Ancón, 0843-03092, Panama

^c Florida Museum of Natural History, University of Florida, Gainesville, FL, 32611, USA

^d University of Florida, Gainesville, FL, 32611, USA

^e Universidade Federal Fluminense, Instituto de Biologia, Departamento e Pós-graduação em Biologia Marinha e Ambientes Costeiros, Campus do Valonguinho, Outeiro São João Batista, s/n, 24020-141 Niterói, RJ, Brazil

^f Museo Argentino de Ciencias Naturales, Caballito, Buenos Aires, Argentina

ARTICLE INFO

Article history:

Received 1 December 2014

Accepted 10 February 2015

Available online 9 March 2015

Keywords:

Neogene

Tropical America

Panama Canal Basin

Sharks

Rays

Paleoecology

ABSTRACT

The Late Miocene Chagres Formation from northern Panama contains the youngest outcrops of the Panama Canal Basin. Here we report two chondrichthyan assemblages that include 30 taxa from both the Rio Indio and Chagres Sandstone Members of the Chagres Formation. We report 18 new fossil records for Panama and four for tropical America, constituting the most diverse chondrichthyan association for the Cenozoic of Panama. We performed a paleobathymetry analysis based on the modern water depth preference of extant chondrichthyan taxa. The assemblage from the Rio Indio Member is characterized by taxa with neritic affinities, suggesting depths <100 m, whereas the assemblage from the Chagres Sandstone Member is dominated by taxa with oceanic affinities, suggesting 200–300 m water depths. The Chagres Sandstone Member could have accumulated at the edge of a platform–upper slope, bordered by a deep oceanic margin.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

Panamanian sedimentary basins preserve an extensive Neogene fossil record of marine biota useful for understanding the evolution of the Isthmus of Panama. The isthmus was formed by complex tectonic processes that isolated the Atlantic from the Pacific Ocean and connected North and South America, precipitating major biogeographic, oceanographic, and environmental changes (Coates and Obando, 1996; Woodburne, 2010; Coates and Stallard, 2013; Leigh et al., 2013). The formation of the Isthmus of Panama was completed during the Pliocene, between 4.2 and 3.5 Ma (Duque-Caro, 1990; Coates et al., 1992, 2003, 2004; Coates and Obando, 1996; Haug and Tiedemann, 1998; Bartoli et al., 2005; Woodburne, 2010; Haug et al., 2001; Coates and Stallard, 2013).

Additional geological evidence suggests that the rise of the isthmus was considerably older and more complex, with an initial collision between South America and the Panama Block between 25 and 23 Ma (Farris et al., 2011; Montes et al., 2012a, 2012b). By 20 Ma, Panama would have been a peninsula connected to North America (MacFadden et al., 2010; Montes et al., 2012a; Rincon et al., 2012), with an oceanic pathway between Panama and South America, the Central American Seaway (CAS, defined here as the deep oceanic seaway along the tectonic boundary of the South American plate and the Panama microplate), connecting the Pacific and Atlantic Oceans (Montes et al., 2012a, 2012b). By 10 Ma, a full closure of CAS had occurred, ending the exchange of deep and intermediate waters between the Caribbean and the Pacific (Coates et al., 2004; Montes et al., 2012a; Sepulchre et al., 2014). However, shallow-water exchange continued between the two oceans along pathways other than CAS, allowing the migration of chondrichthyans between basins (Pimiento et al., 2013b), until the complete rise of the Isthmus at 3.5 Ma (Duque-Caro, 1990; Haug and Tiedemann, 1998; Haug et al., 2001; Coates et al., 1992, 2003, 2004; Coates and Stallard, 2013). Ultimately, the formation of the Isthmus of

* Corresponding author.


E-mail addresses: jorge.carrillo@pim.uzh.ch, jorgedcb100@gmail.com, mad_design@gmx.ch (J.D. Carrillo-Briceño), degracia@gmail.com (C. De Gracia), pimiento@ufl.edu (C. Pimiento), orangelaguilera@gmail.com (O.A. Aguilera), santamarinape@gmail.com (P. Santamarina), jaramillo@si.edu (C. Jaramillo).

<http://dx.doi.org/10.1016/j.jsames.2015.02.001>

0895-9811/© 2015 Elsevier Ltd. All rights reserved.

CHAPTER 4

Sawfishes and Other Elasmobranch Assemblages from the Mio-Pliocene of the South Caribbean (Urumaco Sequence, Northwestern Venezuela)



“Big tooth” †*Carcharocles megalodon*

RESEARCH ARTICLE

Sawfishes and Other Elasmobranch Assemblages from the Mio-Pliocene of the South Caribbean (Urumaco Sequence, Northwestern Venezuela)

Jorge D. Carrillo-Briceño¹✉, Erin Maxwell²✉, Orangel A. Aguilera³✉, Rodolfo Sánchez⁴✉, Marcelo R. Sánchez-Villagra¹✉*

1 Paleontological Institute and Museum, University of Zurich, Zürich, Switzerland, **2** Staatliches Museum für Naturkunde, Stuttgart, Germany, **3** Universidade Federal Fluminense, Instituto de Biologia, Campus do Valonguinho, Outeiro São João Batista, Niterói, Rio de Janeiro, Brasil, **4** Museo Paleontológico de Urumaco, Urumaco, estado Falcón, Venezuela

✉ These authors contributed equally to this work.

* m.sanchez@pim.uzh.ch



OPEN ACCESS

Citation: Carrillo-Briceño JD, Maxwell E, Aguilera OA, Sánchez R, Sánchez-Villagra MR (2015) Sawfishes and Other Elasmobranch Assemblages from the Mio-Pliocene of the South Caribbean (Urumaco Sequence, Northwestern Venezuela). PLoS ONE 10(10): e0139230. doi:10.1371/journal.pone.0139230

Editor: Brian Lee Beatty, New York Institute of Technology College of Osteopathic Medicine, UNITED STATES

Received: April 17, 2015

Accepted: September 9, 2015

Published: October 21, 2015

Copyright: © 2015 Carrillo-Briceño et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All data has been collected, prepared and deposited in a public repository by the authors of the article. The current data is based only in fossil specimens, to which it has been assigned a specific catalog number. All these fossil specimens used to do this publication, are completely available in a public repository. The context information for each fossil specimen, is saved in an internal data base of the institution. This public repository is: the Paleontological Collection of The

Abstract

The Urumaco stratigraphic sequence, western Venezuela, preserves a variety of paleoenvironments that include terrestrial, riverine, lacustrine and marine facies. A wide range of fossil vertebrates associated with these facies supports the hypothesis of an estuary in that geographic area connected with a hydrographic system that flowed from western Amazonia up to the Proto-Caribbean Sea during the Miocene. Here the elasmobranch assemblages of the middle Miocene to middle Pliocene section of the Urumaco sequence (Socorro, Urumaco and Codore formations) are described. Based on new findings, we document at least 21 taxa of the Lamniformes, Carcharhiniformes, Myliobatiformes and Rajiformes, and describe a new carcharhiniform species (†*Carcharhinus caquetius* sp. nov.). Moreover, the Urumaco Formation has a high number of well-preserved fossil *Pristis* rostra, for which we provide a detailed taxonomic revision, and referral in the context of the global Miocene record of *Pristis* as well as extant species. Using the habitat preference of the living representatives, we hypothesize that the fossil chondrichthyan assemblages from the Urumaco sequence are evidence for marine shallow waters and estuarine habitats.

Introduction

The Caribbean Sea today is environmentally stable and ecologically complex, and it is characterized by high fish diversity [1]. However, the origin of this regional diversity is still somewhat problematic. In particular, Neogene chondrichthyan faunas in the Caribbean region are still little known in comparison with other regions as Europe and North America (e.g. [2]), although several previous works from diverse sedimentary basins exist (e.g. [3–24]) documenting the paleodiversity of Caribbean faunas before and after the complete closure of the Central American Seaway [25].

Alcaldía Bolivariana del Municipio Urumaco and Museo Paleontológico de Urumaco, Estado Falcón, Venezuela. This paleontological collection does not have a website with a public databank. Access to this paleontological collection is permitted for national or foreign researchers with previous permission. Access to the collection is obtained by a request directly with the curator of paleontology collection and the competent authorities of the Bolivarian Mayoralty Urumaco, Falcon State, Venezuela. Other fossils referred and illustrated in our article, belong to the Museo de Ciencias de Caracas (MCNC) and to the Centro de Investigaciones Antropológicas, Arqueológicas y Paleontológicas de la Universidad Experimental Francisco de Miranda, Falcón estate, Venezuela. These institutions have paleontological collections with public access, which is obtained by a request directly with the curator of the collection and the competent authorities. All relevant data necessary to replicate the results presented in our article, are available in these paleontological collections of public access referred above.

Funding: This work was supported by Swiss National Fund SNF 31003A-149605 to MRSV (www.snf.ch). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

During the Miocene, northern Venezuela was close to the gateway between the Atlantic and Pacific Oceans prior to the definitive closure of the Panamanian Isthmus [25], and so faunal changes in this area provide an important perspective on changes in the Caribbean region as a whole. Neogene elasmobranchs from the Miocene of Venezuela have been described previously (e.g., [3, 15, 17, 20, 26]), and here we present new data on the elasmobranchs from the Urumaco sequence of northern Venezuela.

The Urumaco stratigraphic sequence is composed of seven geological units [27], represented by diverse paleoenvironment facies including marine, estuarine, riverine, lacustrine and terrestrial [27, 28]. Throughout the entire section [27], the lithology varies between more terrestrially influenced beds such as coal seams, and marine-influenced facies including sandstones, limestones and shales. The Urumaco sequence exhibits the most diverse vertebrate fauna from the Neogene of the southern Caribbean, including marine, estuarine, and freshwater fishes, freshwater and marine turtles and crocodilians, terrestrial and aquatic/semiaquatic mammals, and birds (e.g., [29–32]). The stratigraphic sequence and the associated fauna provide unequivocal evidence of a marine coastal/estuarine environment, heavily influenced by a complex hydrographic system that flowed from western Amazonia to the proto-Caribbean Sea during the Miocene (e.g. [27, 31, 33–38]). More than 20 years of paleontological expeditions to the Urumaco region made available the collection of elasmobranch assemblages reported here. We provide a taxonomic revision of the elasmobranch fauna from the Socorro, Urumaco and Codore formations and discuss the paleoenvironmental implications. Importantly, we include new shark and ray occurrence data for Tropical America, and we provide a special detailed taxonomic revision of the specimens of *Pristis* “sawfishes” (Pristidae) found in these assemblages.

Material and Methods

The fossil elasmobranch fauna from the Urumaco sequence presented here consists of 1234 specimens, and comes from 14 localities of the Socorro, Urumaco and Codore formations (Fig 1, S1 Table, S1 and S2 Appendix). Bulk samples of 10 kg each were collected on the outcrop and the sediment was screen washed using 0.5 mm open mesh. The specimens were sorted using a stereomicroscope. Large specimens were surface-collected from the outcrop. The material was collected by the authors and other collaborators since 1993 during several expeditions. The specimens are deposited in the paleontological collections of the Alcaldía Bolivariana de Urumaco (AMU-CURS), Centro de Investigaciones Antropológicas, Arqueológicas, Paleontológicas of the Universidad Experimental Francisco de Miranda (CIAAP, UNEFM-PF), and the Museo de Ciencias de Caracas (MCNC). The systematics for fossil and recent taxa follows Compagno [39] and Cappetta [2], with the exception of the extinct genus *Carcharodes* Jordan and Hannibal [40], whose assignment here has continued the discussion presented by Pimiento et al. [21]; the dental terminology follows Cappetta [2]. Terminology of the internal pristids rostral cavities follows Wueringer et al. [41]. Taxonomic identification included an extensive bibliographical review and comparative studies with fossil and extant specimens from the following collections: Fossil vertebrate section of the Museum für Naturkunde in Berlin (MB), Museo Nacional de Historia Natural de Santiago (SGO-PV) in Chile, Museu Paraense Emílio Goeldi (MPEG-V), Brazil; Natural History Museum of Basel (NMB), Switzerland; Palaeontological Institute and Museum at the University of Zurich (PIMUZ) Switzerland; René Kindlimann (private collection), Switzerland; Smithsonian Tropical Research Institute, Panama (STRI-PPP-T). We gathered habitat information of all taxa with living representatives using Compagno [42, 43], Compagno et al. [44], Musick et al. [45], Voigt and Weber [46], and the FishBase website (<http://www.fishbase.org>). The S2 Table is based on Aguilera [19], Aguilera and Lundberg [20], Lundberg et al. [47], Aguilera and Marceniuk [48], and Aguilera et al. [31].




Fig 1. Locality map. Fossiliferous localities of the Urumaco sequence.

doi:10.1371/journal.pone.0139230.g001

Measurements for *Pristis* rostra follow the recommendations and methods proposed by Whitty et al. [49]. Rostral spine counts for extant species were obtained from the literature, as well as from examination of comparative material housed at the Museum of Natural History Vienna and the Zoological Museum of the University of Zurich. Rostral cavities in the fossil pristids were visible due to pre-existing breaks.

Nomenclatural Acts

The electronic edition of this article conforms to the requirements of the amended International Code of Zoological Nomenclature, and hence the new names contained herein are

available under that Code from the electronic edition of this article. This published work and the nomenclatural acts it contains have been registered in ZooBank, the online registration system for the ICZN. The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed through any standard web browser by appending the LSID to the prefix "<http://zoobank.org/>". The LSID for this publication is: [urn:lsid:zoobank.org:pub:urn:lsid:zoobank.org:pub:8A290D4C-7169-4300-A15C-70942B19D153](http://zoobank.org/pub:urn:lsid:zoobank.org:pub:8A290D4C-7169-4300-A15C-70942B19D153). The electronic edition of this work was published in a journal with an ISSN, and has been archived and is available from the following digital repositories: PubMed Central, LOCKSS.

Geological Context

Socorro Formation (Middle Miocene)

The lower and upper contacts of the Socorro Formation (Fig 2) are conformable with the Querales and Urumaco formations, respectively [27, 50, 51]. Hambalek et al. [51] informally divided the Socorro Formation into three members. Later, Quiroz and Jaramillo [27] conducted a detailed sedimentological stratigraphic study, identifying a thickness of 2300 m to the whole Socorro Formation along Pauji Creek, 20 km east of the town of Urumaco. The Middle Member is 880 m thick, and is characterized by complex interbedding of medium to fine-grained sandstone, organic mudstone, coal, shales and coquinoïdal limestones with abundant mollusks remains [27]. The Upper Member is 639 m thick and it is characterized by gray, massive-bedded, sandy mudstone interbedding with organic mudstones with plants and coal fragments and scarce coquinoïdal limestones [27, 51].

Urumaco Formation (middle–late Miocene)

The lower and upper contacts of the Urumaco Formation (Fig 2) are conformable with the Socorro and Codore formations, respectively [27, 51]. The unit has a thickness of approximately 1700–2060 m [27, 50], and toward the east it is correlative with the Caujarao Formation that crops out in the Coro-La Vela region [27, 50]. The unit has been divided in three members: the Lower Member, 780 m thick, is dominated by dark-gray laminated mudstones and shales with interbedding fine-grained sandstone and some coquinoïdal limestone layers; the Middle Member is characterized by interbedding of medium-to fine-grained sandstone, organic-rich mudstone, shales and coquinoïdal limestones; the Upper Member is composed of laminated organic-rich mudstone and shales intercalated with very fine grained sandstones [27, 50].

Codore Formation (late Miocene–Pliocene)

The lower and upper contacts of the Codore Formation (Fig 2) are conformable with the Upper Member of the Urumaco Formation and the basal conglomeratic level of the San Gregorio formations, respectively [27, 52]. The Codore Formation has been divided into three formal members: El Jebe Member with continental deposits 475 m thick, the Chiguaje Member with marine deposits 65 m thick, and the Algodones Member with continental deposits 320 m thick [27, 52, 53]. The marine Chiguaje Member is characterized by dark-gray laminated mudstones interbedding with fine-grained muddy sandstones and coquinoïdal limestones with oysters [27]. Rey [53] assigned an early-late Pliocene age to the Chiguaje Member, based on planktonic foraminifera. Hambalek et al. [51] suggested a late Miocene–Pliocene age for the same member based on palynomorphs, and recently Smith et al. [54] suggested a late Miocene–middle Pliocene age, based on planktonic foraminifera.




Fig 2. Stratigraphic section of the Socorro, Urumaco and Codore formations. Modified from Quiroz and Jaramillo [27].

doi:10.1371/journal.pone.0139230.g002

Results

Elasmobranch—systematics

The elasmobranch assemblages from the section of the Urumaco sequence studied here include at least 21 taxa attributed to 13 genera, 9 families and 4 orders (S1 Table).

Neoselachii Compagno, 1977 [55]

Galeomorphii Compagno, 1973 [56]

Lamniformes Berg, 1937 [57]

†Otodontidae Glikman, 1964 [58]

†*Carcharocles* Jordan and Hannibal, 1923 [40]

†*Carcharocles megalodon* (Agassiz, 1843) [59] (Fig 3A–3H)

Description. Fifteen isolated teeth (five upper, seven lower and three of indeterminate jaw position) from the Urumaco (middle–late Miocene) and Codore (late Miocene–Pliocene) formations (S2 Appendix). The crown is triangular in shape, with a flat and convex labial and lingual face, respectively; the lingual face has a typical large neck (inverse “V”-shaped) between




Fig 3. Lamniform and carcharhiniform sharks of the Urumaco sequence. (A-H)

doi:10.1371/journal.pone.0139230.g003

the crown and the root. Cutting edges are finely serrated. In all teeth, the root is high, its root lobes are well developed and a weak lingual protuberance is present. The teeth range in height from 23 to 99 mm and width from 35 to 69 mm.

†*Carcharocles megalodon*: Urumaco Fm. [A-B: UNEFM-CIAAP-1292, C: AMU-CURS-455, E: UNEFM-PF-388, F-G: AMU-CURS-605, and H: AMU-CURS-338] and Codore Fm. [D: UNEFM-PF-351]. (I-L) †*Hemipristis serra*: Urumaco Fm. [I: AMU-CURS-332, and J: AMU-CURS-331] and Codore Fm. [K-L: AMU-CURS-331]. (M-N) *Paragaleus* sp.: Urumaco Fm. [AMU-CURS-640]. (O-R) *Galeocерdo cuvier*: Urumaco Fm. [O-P: UNEFM-PF-408] and Codore Fm. [Q-R: AMU-CURS-625]. (S-Y) *Rhizoprionodon* sp.: Urumaco Fm. [T-U: AMU-CURS-497, V-W: AMU-CURS-478, and X-Y: AMU-CURS-408] and Codore Fm. [AMU-CURS-635]. View: labial (B, K, N-O, R, U-W, Y), lingual (A-J, M, P-Q, S-T, V, X), profile (L). Abbreviations: Fm., Formation.

Carcharhiniformes Compagno, 1973 [56]

Hemigaleidae Hasse, 1879 [60]

Hemipristis Agassiz, 1835 [59]

†*Hemipristis serra* (Agassiz, 1835) [59] (Fig 3J–3L)

Description. Nineteen isolated teeth (nine upper, eight lower and two of indeterminate jaw position) from the Urumaco (middle–late Miocene) and Codore formations (late Miocene–Pliocene) (S2 Appendix). Upper teeth are labio-lingually compressed, with a triangular crown curved distally. The mesial cutting edge is strongly convex and bears fine serrations that end shortly before reaching the apex; distal edge is concave and coarsely serrated with the serrated part also terminating before reaching the apex. The root is high and compressed, with a strong lingual protuberance. Lower teeth with a long and lingually inclined unserrated crown; there are small cusplets near the crown base and the root is bilobate with a strong lingual protuberance. The teeth range in height from 10 to 47 mm and width from 9 to 46 mm.

Paragaleus Budker, 1935 [61]

Paragaleus sp. (Fig 3M and 3N)

Description. One lower antero-lateral tooth from the Urumaco Formation (middle–late Miocene) (S2 Appendix). The crown is prominent and straight, with smooth concave and convex mesial and distal edges, respectively. The distal heel is shorter than the mesial one, and both are low, smooth, without cusps and slightly recurved to the lingual face. The root is low with a flat base, and the lingual protuberance has a well-defined groove. The tooth height is 4 mm and the width is 4.9 mm.

Remarks. Four extant *Paragaleus* species are known from the Eastern Atlantic and Indo-Pacific regions [44], and two extinct species have been described to the Miocene of Europe [2]. Extant dentitions of *Paragaleus* spp. show strong dignathic and gradient heterodonty; for this reason a single tooth is unlikely to be diagnostic for species. The tooth AMU-CURS-640 resembles the specimen referred to *Paragaleus* sp. by Carrillo-Briceño et al. [62] from the early Pleistocene of Ecuador.

Carcharhinidae Jordan and Evermann, 1896 [63]

Galeocерdo Müller and Henle, 1837 [64]

Galeocерdo cuvier (Perón and Lesueur, 1822) [65] (Fig 3O–3R)

Description. Four isolated teeth with an unknown jaw position, from the Urumaco (middle–late Miocene) and Codore (late Miocene–Pliocene) formations (S2 Appendix). The crown is triangular and curved distally, the labial face is flat and the lingual one is more convex. The mesial cutting edge is convex and slightly sigmoidal with minor serrations; distal cutting edge is shorter and slightly straight with small serrations. The distal heel is long and strongly serrated. The root is low and slightly curved with a weak lingual protuberance. Complete teeth the range in height from 9 to 13 mm, and in width from 15 to 26 mm.

Rhizoprionodon Whitley, 1929 [66]

Rhizoprionodon sp. (Fig 3T–3Y)

Description. Fourteen isolated teeth (four upper, six lower and four of indeterminate jaw position) from the Urumaco (middle–late Miocene) and Codore (late Miocene–Pliocene) formations (S2 Appendix); most of them are broken and eroded. In upper teeth, the crown is triangular and asymmetric, with a high cusp inclined distally. The mesial cutting edge is slightly concave and the distal cutting edge is slightly straight or sigmoidal; both cutting edges are smooth. The distal heel is high, clearly separated from the cusp by a notch, and this is rounded or finely serrated, depending of the specimen. The root is low with a flat basal face; the labial face is rectilinear and the lingual face has a slight lingual protuberance with a deep medial groove. Complete teeth range in height from 2.5 to 4.4 mm, and in width from 3.5 to 6.3 mm. Due to few diagnostic characters and fragmentary preservation, specimens are not diagnostic to the species level.

Carcharhinus Blainville, 1816 [67]

†*Carcharhinus caquetius* sp. nov. (Fig 4A–4)

ZooBank life science identifier (LSID) for species. [urn:lsid:zoobank.org:act:BE13DF0A-E6F5-44D5-830A-531EE792183D](https://zoobank.org/urn:lsid:zoobank.org:act:BE13DF0A-E6F5-44D5-830A-531EE792183D)

Etymology. The species is named in honor of the Caquetío pre-hispanic tribe who lived in the Falcon state and other parts of western Venezuela.

Type locality. Puente Río Urumaco, Urumaco Formation (middle–late Miocene), Middle Member, on the right bank of the Urumaco River (near the Urumaco River bridge), northwestern Venezuela (Figs 1 and 2).

Diagnosis. Carcharhinid shark teeth that differ from all other living and extinct *Carcharhinus* species by the following unique combination of dental characteristics: 1) crown with a triangular, strongly elongated and sigmoidal cusp; 2) cusp serrated approximately to its midpoint, with a smooth apex; 3) convex distal heel clearly separated by a deep notch; 4) heel characterized by irregular, small cusps.

Referred material. Holotype: upper lateral tooth (AMU-CURS-499) (Fig 4A–4C); Paratypes: two upper lateral teeth (AMU-CURS-477) (Fig 4D–4F and 4I and 4J), all specimens come from the Middle Member of the Urumaco Formation (Puente Río Urumaco locality).

Description. The teeth range in height from 6 to 7 mm and width from 6 to 7.2 mm, including one upper tooth (3406-T-2) from the Angostura Formation (late Miocene), Ecuador (Fig 4G–4H). Teeth are characterized by a triangular shape with an elongate-sigmoidal, narrow and distally deflected cusp. The lingual face is convex and the labial face slightly convex. The mesial cutting edge is slightly convex and sigmoid, without a clear differentiation between it and the mesial heel; the distal edge is concave in the lower part and sigmoid in the upper part. The mesial crown edge has a strong serration that decreases as it approaches the cusp, giving way to a smooth edge at the apex on both edges of the cusp. The distal heel consists of irregular, small cusps, and is well-separated from the distal edge by a deep notch. The root is thick and slightly concave in the base with slightly rounded lobes; the labial surface is flat and the lingual surface has a shallow lingual protuberance with a shallow lingual groove. To date, we have not found any lower tooth with diagnostic features that justify referral to this new taxon.

Remarks. Some extant species of *Carcharhinus*, such as *C. borneensis* (Bleeker, 1859) [68], *C. dussumieri* (Valenciennes, 1839) [69], *C. fitzroyensis* (Whitley, 1943) [70], *C. hemiodon* (Valenciennes, 1839) [69], *C. porosus* (Ranzani, 1839) [71], *C. sealei* (Pietschmann, 1913) [72], *C. signatus* (Poey, 1868) [73], and *C. sorrah* (Valenciennes, 1839) [69], have upper teeth with a generalized morphological pattern characterized by triangular and distally inclined crowns with coarse serration of the distal heel (e.g., [46]). However, diagnostic elements in the morphology of the cups and the regular serration of the distal heel in the extant taxa previously mentioned permit a clear differentiation between these and the upper teeth of *Carcharhinus caquetius* sp. nov., the teeth of which have a diagnostic elongated, sigmoidal cusp, with a




Fig 4. Carcharhiniform sharks of the Urumaco sequence. (A–J) †*Carcharhinus caquetius* sp. nov. Holotype: A–C [AMU-CURS-499], paratypes: D–F, I–J [AMU-CURS-477]. (G–H) specimen of †*Carcharhinus caquetius* sp. nov. [3406-T-2, 3457-T-(1)] from the Angostura Formation, Ecuador. (K–Q) *Carcharhinus leucas*: Urumaco Fm. [K–L: AMU-CURS-360, M–N, Q: AMU-CURS-368], and Codore Fm. [O–P: AMU-CURS-622]. (R–S) *Carcharhinus limbatus*: Urumaco Fm. [AMU-CURS-456]. (T–X) *Carcharhinus obscurus*: Urumaco Fm. [T: AMU-CURS-464, U–V: AMU-CURS-463, W–X: AMU-CURS-466]. View: labial (A, F, H–I, L–M, R, V–W), lingual (C, D, G, J, K, N–Q, S–U, X), profile (B, E). Abbreviations: Fm., Formation.

doi:10.1371/journal.pone.0139230.g004

characteristic distal heel conformed by irregular cusps. The extant species most closely related to *C. caquetius* sp. nov. could be *C. signatus*; nevertheless, in the upper teeth of this taxon, such characters as a strong serration in the mesial heel, a strong and regular serration in the distal heel, a flattened labial face, and a smooth or slightly serrated, wider, shorter and straighter cusp, allow differentiation from *C. caquetius* sp. nov. The oldest record of *C. signatus* is from the late Miocene of Panama, where a few specimens have been collected from the Chagres Formation [24]. Some species of *Carcharhinus* have ontogenetic and sexual dental variation [2, 43, 46], which makes the taxonomic determination of isolated teeth difficult, especially when only

few specimens are available. However, the teeth described as *C. caquetius* sp. nov. from the Urumaco Formation, and the specimens from the Angostura Formation (Ecuador), are clearly different from the teeth of other *Carcharhinus* species.

Carcharhinus leucas (Müller and Henle, 1839) [74] (Fig 4K–4Q)

Description. Twenty-eight isolated teeth (twenty upper and eight lower) from the Urumaco (middle–late Miocene) and Codore (late Miocene–Pliocene) formations (S2 Appendix). Upper teeth are characterized by crowns with a triangular shape; the lingual face is convex and the labial one is flat. Cutting edges are regularly serrated; the serrations become progressively coarser near the base of the crown. The mesial cutting edge is straight or slightly convex, and the distal one is slightly concave or straight; the distal heel has a typical serrated blade shape. The root is thick and concave along the basal margin. The labial face of the root is flat and the lingual face is convex and is characterized by a weak lingual groove. Lower teeth have a thick and finely serrated crown with the lateral heel finely serrated; the base of the root is slightly concave. The teeth range in height from 12 to 21 mm and width from 10 to 21 mm.

Carcharhinus limbatus (Müller and Henle, 1839) [74] (Fig 4R and 4S)

Description. Two upper lateral isolated teeth from the Urumaco Formation (middle–late Miocene) (S2 Appendix). The crown is high, triangular and slightly convex with the apex inclined distally. The lingual face is convex and the labial one is flat. The mesial cutting edge is slightly convex and the distal cutting edge concave; both are serrated. The serrations on the mesial and distal heel are more developed and well-differentiated than the cutting edges. The root is wider than the crown and its base slightly concave; the labial face is flat and the lingual protuberance is slightly developed and has a shallow groove. The teeth range in height from 7 to 12 mm and width from 8 to 10 mm.

Carcharhinus obscurus (Lesueur, 1818) [75] (Fig 4T–4X)

Description. Twenty isolated teeth (sixteen upper and four lower) from the Urumaco Formation (middle–late Miocene) (S2 Appendix). Upper teeth are characterized by a triangular shape and lack of an elongate crown with regular serration. The mesial cutting edge is apically convex, with the apex deflected distally; the distal cutting edge is concave. The root is thick, low and slightly concave in the basal margin with slightly rounded lobes; the lingual surface of the root is convex with a shallow lingual groove. Lower teeth have erect crowns finely serrated and moderately arched root lobes. The teeth range in height from 12 to 21 mm and width from 10 to 21 mm.

Carcharhinus plumbeus (Nardo, 1827) [76] (Fig 5A–5F)

Description. Four upper isolated teeth from the Codore Formation (late Miocene–Pliocene) (S2 Appendix). The crown is triangular in shape and strongly elongated with the apex deflected slightly distally; the labial face is flat and the lingual one is convex. The convex mesial crown edge is continuous with the mesial heel. The distal edge is convex or slightly straight and can be differentiated from the distal heel, but a notch is lacking. The cutting edges and heels are uniformly and finely serrated. The root base is slightly concave; the lingual face is convex and it is characterized by a lingual groove. The teeth range in height from 15.5 to 18 mm and width from 7 to 22 mm.

Carcharhinus porosus (Ranzani, 1839) [71] (Fig 5G–5J)

Description. Eight isolated teeth (five upper and three lower) from the Urumaco (middle–late Miocene) and Codore (late Miocene–Pliocene) formations (S2 Appendix). In the upper teeth, the crown is triangular and asymmetric, with a high cusp inclined distally. The mesial edge is straight or slightly concave, and in most of these there is no differentiation between this and the mesial heel; the distal edge is straight. Both cutting edges are strongly serrated, and the serrations decrease in size from the base to the apex. The distal heel is long, strongly serrated and clearly separated from the cusp by a deep notch. The root is low with a slightly concave




Fig 5. Carcharhiniform sharks of the Urumaco sequence. (A–F) *Carcharhinus plumbeus*: Codore Fm. [A–F: AMU-CURS-624]. (G–J) *Carcharhinus porosus*: Urumaco Fm. [G–H: AMU-CURS-472, and I: AMU-CURS-590], and Codore Fm. [J: AMU-CURS-631]. (K–L) *Carcharhinus* spp.: Socorro Fm. [K–L: AMU-CURS-630]. (M–R) *Negaprion brevirostris*: Urumaco Fm. [M–N: AMU-CURS-468, O–P: AMU-CURS-469], and Codore Fm. [Q–R: AMU-CURS-626]. (S–V) *Sphyrna* cf. *zygaena*: Urumaco Fm. [AMU-CURS-474; AMU-CURS-477]. View: labial (A, C, F, G, K, M, O, R, T, V), lingual (B, D–E, H–J, L, N, P–Q, S, V). Abbreviations: Fm., Formation.

doi:10.1371/journal.pone.0139230.g005

base. Lower teeth are incomplete, and show oblique, narrow, serrated cusps with well-differentiated and serrated heels. The teeth range in height from 4.5 to 5 mm and width from 5 to 5.2 mm.

Carcharhinus spp. (Fig 5K and 5L)

Description. Eighty-two incomplete, eroded and non-diagnostic teeth (thirty upper and fifty-two lower) from the Socorro (middle Miocene) and Urumaco (middle–late Miocene) formations (S2 Appendix).

Negaprion brevirostris (Poey, 1868) [73] (Fig 5M–5R)

Description. Twenty-five isolated teeth (fifteen upper and ten lower) from the Socorro (middle Miocene), Urumaco (middle–late Miocene) and Codore (late Miocene–Pliocene) formations (S2 Appendix). Upper teeth show a high and triangular cusp slightly inclined distally. The cutting edges are completely smooth, except the low and weakly serrated lateral heels. The root branches are extended and the basal face is rather broad and flat, with a clear medial

lingual groove. Lower teeth are narrower than the upper ones and their heels are generally not serrated. The teeth range in height from 12 to 23 mm and in width from 16 to 22 mm.

Sphyrnidae Gill, 1872 [77]

Sphyrna Rafinesque, 1810 [78]

Sphyrna cf. *zygaena* (Linnaeus, 1758) [79] (Fig 5S–5V)

Description. One upper lateral tooth and one lower lateral tooth from the Urumaco Formation (middle–late Miocene) (S2 Appendix). The upper lateral tooth (height: 12.3 mm) has a high, triangular, asymmetric, broad and distally inclined crown (Fig 5S and 5T); the lingual face is strongly convex and the labial one is flat. The mesial cutting edge is slightly convex and slightly sigmoid and the distal cutting edge is straight; both cutting edges are completely smooth. The distal heel is low and well separated from the cusp by a shallow notch. The root is low with the basal surface slight concave, and is missing the mesial lobe; the lingual protuberance is well developed, bearing a groove. The lower lateral tooth has a stout triangular, slightly serrated crown (Fig 5U and 5V). The lingual face is convex and the labial face is flat. The cusp is inclined distally, being broken and incomplete in the apex. The mesial cutting edge is straight and there is not a clear differentiation between this and the mesial heel; the distal cutting edge is straight. The distal heel is large without serrations and is well separated from the cusp by a shallow notch. The root has a slightly concave base and the lingual protuberance is well developed, bearing a deep groove.

Remarks. Juveniles and young adults of extant *Sphyrna zygaena* have teeth with smooth cutting edges, but larger individuals may have slightly serrated teeth [80, 81].

Batomorphii Cappetta, 1980 [82]

Myliobatiformes Compagno, 1973 [56]

Dasyatidae Jordan, 1888 [83]

cf. *Dasyatis* Rafinesque, 1810 [78] (Fig 6A–6V)

Description. Sixteen isolated teeth of indeterminate jaw position from the Urumaco (middle–late Miocene) and Codore (late Miocene–Pliocene) formations (S2 Appendix). The teeth are small, only a few millimeters wide. The crown shows a middle transverse crest that separates the labial and lingual faces; the crest is lingually elongate and forms a distinctive cusp in male teeth (e.g. Fig 6G–6J, 6U and 6V). The labial face is sub-triangular in shape, with a distinctive alveolate ornamentation, and a labio-lingually extended depression; the lingual face is smooth and is divided into two lingual marginal areas by a lingual ridge. The root has two lobes, which are lingually arched with a flat or slightly bulging base. There is a foramen in the middle of the sulcus.

Remarks. *Dasyatis* shows a dental morphological diversity with distinct gynandric heterodont patterns in most species, which may be more complex at the generic level than currently accepted [2]. Knowledge of dental patterns in extant and fossil *Dasyatis* and others Dasyatidae is still scarce, making any taxonomic assignment of fossil specimens to species difficult.

Dasyatidae indet. (Fig 6W–6X)

Description. One small tooth (less than 2 mm length) with a rhomboid and labial-lingually elongated crown from the Urumaco Formation (middle–late Miocene). The occlusal surface is flat and appears to be eroded; however, this has a light alveolate ornamentation. The root is massive and presents two distinct lobes with a flat basal faces that becomes narrower lingually; between these there is a deep sulcus with two foramina. Due similarities in dental morphology among dasyatid genera and the few fossil specimens referred herein, we maintain this specimen in open nomenclature.

Myliobatidae Bonaparte, 1838 [84]

Aetobatus Blainville, 1816 [67]

Aetobatus cf. *narinari* (Euphrasen, 1790) [85] (Fig 7A–7G)




Fig 6. Rays of the Urumaco sequence. (A–V) cf. *Dasyatis*: Urumaco Fm. [A–D: AMU-CURS-589, E–F, I–J: AMU-CURS-495, G–H: 496], and Codore Fm. [M–V: AMU-CURS-636]. (W–X) *Dasyatidae* indet.: Urumaco Fm. [AMU-CURS-591]. View: labial (N, P, U), profile (B, F, H, J, L, R, T, V, X), lingual (W), occlusal (A, C–D, E, G, I, K, M, O, Q, S). Abbreviations: Fm., Formation.

doi:10.1371/journal.pone.0139230.g006

Description. Fifty-seven incomplete, isolated teeth (eight upper and forty-nine lower) and two fragmented upper dental plates, from the Socorro (middle Miocene), Urumaco (middle-late Miocene) and Codore (late Miocene–Pliocene) formations (S2 Appendix). Upper teeth are almost straight and the lower ones are mesio-distally V-shaped. The crown is labio-lingually thicker in the central than in the lateral region. In occlusal view, the crown surface is smooth and the labial and lingual margins are vertical and strongly ornamented; on the lingual margin there are well marked alternating furrows and laminae, whereas these structures are not so evident on the labial one. The root is polyaulacorhize, apico-basally flattened and decreases in height to the lateral edges of the tooth, and is displaced lingually with respect to the crown.

Remarks. Though the teeth from the Urumaco sequence described here are not adequately preserved, their tooth morphology has strong resemblance to those of the recent species *Aetobatus narinari* [86]. At least two nominal species of *Aetobatus* were described from early to late Miocene deposits of Europe: *Aetobatus arcuatus* (Agassiz, 1843) [59] and *Aetobatus cappelletti* Antunes & Balbino, 2006 [87]. *Aetobatus arcuatus* is possibly restricted to the early-middle Miocene, and *A. cappelletti* has only been reported from the late Miocene of Portugal [88]. For a recent discussion of Miocene *Aetobatus* teeth, see Bor et al. [89].

Myliobatis Cuvier, 1816 [90]

Myliobatis sp. (Fig 7H–7J)




Fig 7. Rays of the Urumaco sequence. (A–G) *Aetobatus* cf. *narinari*: Urumaco Fm. [E–G: AMU-CURS-308] and Codore Fm. [A–B: AMU-CURS-598, and C–D: AMU-CURS-614]. (H–J) *Myliobatis* sp.: Urumaco Fm. [AMU-CURS-487]. (K–X) *Rhinoptera* sp.: Urumaco Fm. [K–P: AMU-CURS-288, and Q–S: AMU-CURS-489], and Codore Fm. [T–X: AMU-CURS-621]. (Y–Z) *Myliobatiformes* indet.: Urumaco Fm. [Y: AMU-CURS-492], and Codore Fm. [Z: AMU-CURS-634]. View: basal (A, C, G, M, O, U), dorsal (Z), labial (J, L, S, X), profile (T), lingual (F, H, P, R, V), occlusal (B, D–E, I, K, N, Q,), oblique occlusal (W), ventral (Y). Abbreviations: Fm., Formation.

doi:10.1371/journal.pone.0139230.g007

Description. With 798 teeth (indeterminate upper-lower jaw position), this is the most abundant taxon sampled from the Urumaco (middle–late Miocene) and Codore (late Miocene–Pliocene) formations (S2 Appendix). The specimens consist of complete and fragmented isolated teeth, mainly from the medial (780 specimens) and lateral tooth row (18 specimens). The medial teeth are broader (mesio-distally) than long (labio-lingually), with a hexagonal contour, and in most cases are rectilinear, but some are arched. Teeth of lateral and posterior files are longer than broad, with a hexagonal, pentagonal or triangular contour depending on their position in the files. In all teeth, the occlusal crown face is practically flat and the labial and lingual margins are slightly ornamented; the root has a polyaulacorhize vascularization type. Complete teeth range in width from 8 to 57 mm, and height from 3 to 10 mm.

Remarks. Taxonomic identification based on isolated teeth of *Myliobatis* is difficult due to the large dental variation within the group [86, 91].

Rhinopteridae Jordan and Evermann, 1896 [63]

Rhinoptera Cuvier, 1829 [92]

Rhinoptera sp. (Fig 7K–7X)

Description. Fifty complete and incomplete isolated teeth (14 mesial and 36 laterals of indeterminate upper-lower jaw position) recovered from the Urumaco (middle–late Miocene) and Codore (late Miocene–Pliocene) formations (S2 Appendix). The crown is variably high, hexagonal, straight or slightly convex labially. The occlusal surface is flat and there is a thick basal ledge on the lingual face of the crown. Lateral teeth are less enlarged than the medial ones, showing a regular hexagonal shape; the distal crown height is typically lower than mesial height. The root has polyaulacorhize vascularization type, with a quadrangular shape, and does not project beyond the level of the crown. Complete teeth range in width from 13 to 38 mm, and height from 3 to 9 mm.




Fig 8. Guitarfish and sawfish of the Urumaco sequence. (A–J) *Rhynchobatus* spp.: Urumaco Fm. [A–D: AMU-CURS-482 and E–F: AMU-CURS-628], Codore Fm. [G–J: AMU-CURS-638]. (K–V) *Pristis* sp. Buccal teeth: Urumaco Fm. [K–R: AMU-CURS-484], and rostral spines: Urumaco Fm. [S–T: AMU-CURS-243 and U: AMU-CURS-244], and Codore Fm. [V: AMU-CURS-620]. View: basal (B, H, L), dorsal (S), labial (R), profile (D, F, J, N, P), lingual (C), occlusal (A, I, K, M, O, Q), oblique occlusal (E, G), ventral (U–V), posterior (T). Abbreviations: Fm., Formation.

doi:10.1371/journal.pone.0139230.g008

Remarks. As in *Myliobatis*, the taxonomic identification of isolated teeth of *Rhinoptera* is extremely difficult due to extensive dental variation in the group (e.g., [86]).

Myliobatiformes indet. (Fig 7Y and 7Z)

Description. Six fragmented caudal spines from the Urumaco (middle–late Miocene) and Codore (late Miocene–Pliocene) formations (S2 Appendix). The specimens preserved an evident dorsal groove, a ventral central ridge and strong serrations along both sides of the spine.

Remarks. The majority of myliobatiforms have caudal spines and some characters can be used to identify fossil taxa [86]; however due the fragmentary condition of the specimens referred herein, these characters cannot be usually observed.

Rajiformes Berg 1937 [57]

Rhynchobatidae Garman, 1913 [93]

Rhynchobatus Müller and Henle 1837 [64]

Rhynchobatus spp. (Fig 8A–8J)

Description. Three isolated teeth of indeterminate jaw position from the Urumaco Formation (middle–late Miocene), and two from the Codore Formation (late Miocene–Pliocene) (S2 Appendix). Teeth from the Urumaco Formation have a globular and lozenge-shaped

crown with rounded edges and with a fine granular ornamentation. The labial face is strongly convex and the lingual face presents an uvula that extends beyond the root sulcus. The uvula is flanked on both sides by a slight depression. The occlusal part of the crown in one of the specimens is flattened due to wear (Fig 8A–8D). The root is massive and presents two distinct lobes with a flat basal face that becomes narrower lingually; between these there is a root sulcus with two foramina. The teeth range in width is from 4.9 to 5 mm, and height from 4.7 to 5 mm. The two specimens from the Codore Formation have a globular and lozenge-shaped crown with rounded edges. The labial face is strongly convex and the lingual face presents an uvula that extends beyond the root sulcus. Both specimens have a very prominent arched transverse keel on the crown. The specimen AMU-CURS-638-a (Fig 8I and 8J) shows ridges and a broad, shallow depression on the labial face of the crown. The root is massive and presents two distinct lobes with flat basal faces that become narrower lingually; between these there is a root sulcus with two foramina. The teeth range in width from 2 to 2.1 mm, and height ranges from 1.9 to 2 mm.

Remarks. The specimen AMU-CURS-477-a has morphology similar to an isolated tooth from the late Miocene of Panama referred to *Rhynchobatus* sp. by Pimiento et al. [23]. Because of the range of dental variation in extant species is unknown, and recovered specimens are rare and poor preserved, we refrain from taxonomic identification at species level.

Taxonomic and morphological features of sawfish material in the sequence

The sawfish specimens (Pristidae Bonaparte, 1838 [84]) are represented by two partial rostra from the Socorro Formation (middle Miocene); four buccal teeth, one complete rostrum, 21 partial rostra, 42 rostral spines from de Urumaco Formation (middle–late Miocene), and one rostral spine from the Codore Formation (late Miocene–Pliocene) (S2 Appendix). We examined multiple morphological features, including buccal teeth (Fig 8K–8R), rostral spine shape (Fig 8S–8V), alveolar shape, and the ratio between the total rostral length and the standard rostral length. All specimens recovered from the Miocene of the Urumaco sequence are referable to *Pristis* Linck, 1790 [94] (Figs 8K–8R, 8S–8V and 9A–9L) rather than to *Anoxypristis* White and Moy-Thomas, 1940 [95], based on the internal morphology of the rostrum, as well as the shape of the spines [41]. The most complete specimen from the Urumaco sequence is AMU-CURS-023 (Urumaco Formation, Lower Member) (Fig 9A), previously referred by Aguilera [19] to the extant species *Pristis pectinata* Latham, 1794 [96]. Based on our study of numerous aspects of morphology, the pristids from the Urumaco sequence fall within the rostral morphospace of modern *Pristis* species, however are not easily referable to any one fossil or extant species.

Buccal tooth morphology. The four buccal teeth of indeterminate jaw position (Urumaco Formation, Middle Member) (Fig 8K–8R) are less than 1 mm long. The crown is semiglobular and rounded, with a transverse crest and a long uvula that does not extend beyond the root. The root is much broader than the crown; the furrow is broad and has a large foramen. Oral teeth are somewhat difficult to identify because those of extant species are poorly known and are morphologically variable [2]. According to Cappetta [2], teeth of *Pristis pristis* (Linnaeus, 1758) [79], are longer than broad; however, the teeth of *Pristis pectinata* illustrated by Herman et al. [97], show variability in length and shapes. The specimen from the Urumaco Formation also differs from the buccal teeth of *Pristis aquitanicus* Delfortrie, 1872 [98], described from the early-middle Miocene of France, and illustrated by Cappetta [99].

Rostral spine morphology. Although often referred to as teeth, the rostral spines are not homologous with the buccal teeth of other vertebrates and represent an independent derivation




Fig 9. Sawfish rostra of the Urumaco sequence. (A-L) *Pristis* sp. rostra: Socorro Fm. [E: AMU-CURS-639 and F: AMU-CURS-241] and Urumaco Fm. [A: AMU-CURS-023, B: AMU-CURS-043, C: AMU-CURS-235, D: AMU-CURS-237, G: AMU-CURS-236, H: AMU-CURS-238 and L: AMU-CURS-251]. (I-J) transverse section: [I: AMU-CURS-043 and AMU-CURS-376, Urumaco Fm.]. (K) mineralized cartilage: [AMU-CURS-235]. (L) fragment of rostrum showing the depressions and foramina interpreted as ampullae of Lorenzini. View: dorsal (A-H), posterior (I-J). Abbreviations: Fm., Formation; ad, axial duct; nd: neural ducts.

doi:10.1371/journal.pone.0139230.g009

Table 1. Measurements and ratios for extant *Pristis* spp. and rostra from the Urumaco sequence.

	Rostral spine count	SRL	SRW (TW)	DS	DPS	SRW/SRL	TW/SRL	SRL/TRL
<i>Anoxypristis</i>	16–33			1.1–2.0	0.16–0.44	0.08–0.15	0.05–0.08	0.59–0.81
<i>P. pristis</i>	14–24			2.0–2.9	0.53–1.05	0.15–0.25	0.06–0.1	0.91–0.98
<i>P. clavata</i>	18–27			1.2–2.0	0.25–0.7	0.16–0.22	0.07–0.11	0.89–0.97
<i>P. zijsron</i>	23–37			1.1–2.0	0.1–0.33	0.09–0.17	0.04–0.09	0.85–0.97
<i>P. pectinata</i>	20–30			2.1	0.21–0.41	0.13–0.15	0.07–0.08	0.86–0.99
<i>P. atlanticus</i> [†]	~20	116.3	14.5 (7.0)	—	0.29	0.12	0.06	—
AMU-CURS-023	23/24	70.9	10.7 (5.5)	2.0	0.57	0.15	0.08	0.88
AMU-CURS-235	23	57.7	7.6* (3.4)	—	0.25	0.13	0.06	—
AMU-CURS-237	20+	75.0	8.1 (4.5)	1.65	0.52	0.11	0.06	—

Data from Whitty et al. [49], except *P. pectinata* ratios which are from 14 individuals listed in Robillard and Séret [116], and *P. atlanticus* (rostral spine estimate from Casier [108]; measurement data from Zbyszewski [117]).

*Estimate based on doubling the width of the preserved half.

DS = sum of the distal and next most distal interalveolar ratios from the right and left sides; DPS score = ratio between the spacing of the most anterior two and most posterior two rostral spines; SRL = standard rostral length; SRW = standard rostral width; TRL = total rostral length; TW = tip width

† = extinct taxon.

doi:10.1371/journal.pone.0139230.t001

from highly modified scales [100]. Isolated rostral spines (Fig 8S–8V) have a length up to 95 mm; these are generally long, pointed and slightly curved distally. The anterior edge is sharp and the posterior edge is concave. These rostral spines are rarely preserved in association with rostra from the Urumaco sequence. An exception to this is the specimen AMU-CURS-251 (Urumaco Fm., Upper Member), which preserves a single associated spine, and AMU-CURS-237 and AMU-CURS-639 (Socorro Formation, Upper Member), in which multiple spines are preserved, but are heavily damaged and all lack distal tips. The spines of AMU-CURS-639 are robust, and both the rostral spines of this specimen and the isolated rostral spine of AMU-CURS-376 have a rounded anterior margin and a grooved posterior margin. These features are consistent with *Pristis*, as noted by Cappetta [2] and inconsistent with *Anoxypristis*. Alveoli are anteroposteriorly asymmetrical, with convex anterior edges and weakly concave posterior edges.

Rostral spine count. An accurate rostral spine count was available only for the rostrum AMU-CURS-023 (Fig 9A). In this specimen, 24 alveoli were found on the right-hand side of the rostrum, and 23 on the left. This count is notably lower than the count of 26 recorded by Aguilera [19] for the same specimen. Counts of 23 and 24 are consistent with every extant pristid [49, 101] (Table 1) although at the lower limit of the range for *Pristis zijsron* Bleeker, 1851 [102], and the high end of the range of *P. pristis*. Rostral spine counts appear to vary on a regional basis in pristids, and the Recent Western Atlantic population of *P. pristis* has a lower count than other populations. In contrast, the Western Atlantic population of *P. pectinata* has a higher count [103]. The specimen AMU-CURS-043 (Urumaco Formation, no collections data about member) (Fig 9B), although incomplete posteriorly and with erosion of the posterior penultimate segment preventing any count from this certainly spine-bearing region, shows a count of 22 alveoli on the best preserved side. The specimens AMU-CURS-235 and AMU-CURS-237 (both from the Urumaco Fm., Upper Member) (Fig 9C and 9D), although less well-preserved than AMU-CURS-023 and with some uncertainty regarding posterior completeness, have minimum counts of 23 and 20 alveoli on the right side, respectively (best preserved side).

Internal duct morphology. Several of the *Pristis* rostra from the Urumaco sequence preserve the internal rostral cavities, often filled with sedimentary material that increases contrast

with the rostral cartilages (Fig 9I and 9J). Anteriorly, only the ducts for the ophthalmic and buccal nerves (following Wueringer et al. [41]) are present; the precranial cavity is absent. The neural ducts are roughly ovate, tapering slightly laterally. Posterior to this, the precranial cavity first appears as a dorsal and a ventral opening. Preservation does not permit assessment as to whether these are connected by a medial constriction, or separated by mineralized cartilage (AMU-CURS-241 and AMU-CURS-376); however slightly more posteriorly they are joined by a medial constriction, and the dorsal and ventral portions are expanded parallel to the rostral margins (AMU-CURS-376) (Fig 9J). The precranial cavity becomes square in outline, and increases in size. The neural ducts develop flattened medial edges, and more tapered lateral edges. The dorsal surface of the axial duct is embayed to contain a small foramen located dorsal to the precranial cavity. This secondary duct in the precranial cavity is found in all adequately preserved rostra from the Socorro and Urumaco formations, and has also been figured in some extant *Pristis* material (e.g. [104]). Its presence has been explained as an artifact relating to shrinking of the hyaline cartilage during the preparation of dried specimens [105], and a similar explanation cannot be ruled out in the fossil material. The structure of the internal rostral cavities is broadly similar to those described and illustrated for *Pristis* by Duméril [104], Hoffmann [105], Kirkland and Aguillón-Martínez [106], Cicimurri [107], and Wueringer et al. [41]. The presence of a single, oval to tear-drop shaped neural duct as opposed to two circular ducts differentiates the genera *Pristis* and *Anoxypristis* [41].

Mineralized cartilage organization. The *Pristis* rostra from Socorro and Urumaco formations share a complex pattern of rostral mineralization. There are two thick layers of prismatic calcified cartilage surrounding the external surface of the rostrum, forming embayments for the alveoli and surrounding the neural canals. On the inner surface of the neural canals, a third layer is present, thinner than the two thus described and not as clearly composed of discrete prisms. There also appears to be an additional, very thin layer of mineralized tissue on the external surface of the rostrum. As many of the rostra are quite weathered, this feature can be seen in only a few specimens. It has a fibrous appearance in macroscopic view. Recent sawfish rostra have been described as consisting of three mineralized cartilage layers [105, 108]: an external prismatic layer and prismatic layers encircling the neural ducts, and a layer lining the neural ducts. Hoffmann [105] mentioned the fibrous outer layer of connective tissue, but did not suggest that it was normally mineralized; however this outer mineralized layer was preserved in *Pristis propinquidens* Casier, 1949 [108]. Many authors failed to note the mineralized layer lining the neural ducts [41, 106], but the studies of Hoffmann [105] and Casier [108], demonstrated that this mineralized layer is present in both Recent and fossil *Pristis* species.

Ampullae of Lorenzini. Depressions and foramina are visible in the mineralized fibrous layer on the ventral surface and immediately lateral to the neural duct of one of the more exceptionally preserved rostra AMU-CURS-251 (Fig 9L). These are distributed in approximately two closely spaced rows parallel to the edge of the neural duct, and are interpreted as housing the ampullae of Lorenzini, with the foramina transmitting the sensory nerves from the lateral line and ampullae of Lorenzini. These depressions were interpreted as related to electroreception rather than to the lateral line system, as the latter, while occasionally associated with subdermal tissues [109] is described as being composed of tubules rather than clusters of receptors. In contrast, the ampullary clusters are not part of a continuous tubular system and are described specifically as being embedded in connective tissue [110]. Differences exist in the distribution pattern of the ampullae of Lorenzini between *Anoxypristis*, *Pristis pristis* and *P. clavata* Garman, 1906 [111] (e.g. [110]). There is no information available as to whether the distribution of ampullae in the intact rostrum mirrors the pattern of fossae in the fibrous outer mineralized layer of the ventral surface of the rostrum. Assuming that it does, the distribution

of ampullae seen in the *Pristis* material from the Socorro and Urumaco formations is closer to that of smalltooth sawfish (*P. zizsron*, *P. pectinata*, and *P. clavata*) than to *P. pristis*.

Discussion

Taxonomic composition

The elasmobranch faunal assemblages described here from the Urumaco sequence include at least 21 taxa (S1 Table), of which only three are extinct (*Carcharocles megalodon*, *Hemipristis serra* and *Carcharhinus caquetius* sp. nov.). From the complete assemblages, two taxa were found to be new fossil records for Venezuela and the Caribbean region (*Carcharhinus caquetius* sp. nov., and *Carcharhinus porosus*). The remaining taxa have been recorded from other Neogene marine deposits of Venezuela [20], Central, North, and South America [18, 24, 62]. Many of the taxa from the Urumaco sequence have also been found in Neogene rocks around the world [2], confirming the cosmopolitan distribution of many of these species during the Miocene–Pliocene.

The elasmobranch fauna of the Urumaco sequence shows a clear differentiation in paleodiversity between geologic units. The Urumaco Formation, with 11 localities, has a paleodiversity of 20 taxa, while the Codore Formation, with only one studied locality, is characterized by 16 taxa and the Socorro Formation only by six taxa (S1 Table). The low paleodiversity of the Socorro Formation in comparison with the other units could be attributable to less intensive sampling. In spite of their absence from the Urumaco sequence, taxa such as *Carcharocles megalodon* and *H. serra* have been found in the Socorro Formation at outcrops located closer to Coro City [3].

Fossil taxa such as *Galeocerdo cuvier*, *Rhizoprionodon* sp., *Carcharhinus leucas*, *C. limbatus*, *C. obscurus*, *C. plumbeus*, *C. porosus*, *Negaprion brevirostris*, *Sphyrna* cf. *zygaena*, *Sphyrna* sp., cf. *Dasyatis*, *Aetobatus* cf. *narinari*, *Myliobatis* sp., *Rhinoptera* sp., and *Pristis* sp. have extant counterparts in tropical America and adjacent regions; other taxa such as the genus *Rhynchobatus* live exclusively in the eastern Atlantic (off the African coast) and Indo-West Pacific [112]. The occurrence of *Rhynchobatus* in the marine facies of the Urumaco and Codore formations and in other geologic units of Venezuela [20], Costa Rica [10], and Panamá [23] confirms the presence of this taxon in the Caribbean region during the late Miocene, suggesting that it became extinct from the proto-Caribbean Sea and western Atlantic, possibly as a consequence of environmental changes during the final stage of closure of the Panamanian isthmus [25], or due to competition with other species. The presence of *Carcharhinus caquetius* sp. nov. in the late Miocene of Ecuador suggests a neritic distribution in the proto-Caribbean and Eastern Pacific for this species.

Taxonomic and morphological features of sawfish

There are four valid extant species of *Pristis*, *P. pristis*, *P. clavata*, *P. zizsron* and *P. pectinata*, supported by morphological and molecular data [103]. *P. pristis*, with a circumtropical distribution, is the sister taxon to the others (“smalltooth sawfish”), all of which show more localized distributions either in the Atlantic-Caribbean (*P. pectinata*) or the Indo-West Pacific (other species). The number and spacing of rostral spines has some discriminatory power in identifying the extant species of *Pristis*, although extensive variation within and between populations has been noted [101, 103]. The number of spine positions is determined early in embryological development [113], however, the environmental and developmental factors underlying the number and position of the rostral spines are unknown.

Whether all *Pristis* material from the Urumaco sequence can be referred to a single species must be considered. This question is especially relevant since modern sawfish species have

overlapping geographic ranges [101], and in historic times, the living species *P. pristis* and *P. pectinata* were present sympatrically in the Caribbean and Gulf of Mexico [114, 115]. At present, there seems to be little evidence for two fossil species in the Urumaco Formation, although the range of DPS values is quite broad (Table 1), especially including the specimen AMU-CURS-235. However, based on other metrics this specimen is generally consistent with other material from the locality, and due to poor preservation on the posterior right side, it is possible that rostral spine spacing is anomalous on the left and is giving a false signal.

Based on our study of numerous aspects of morphology, the Miocene pristids from the Urumaco sequence fall within the rostral morphospace of modern *Pristis* spp., nevertheless, are not easily referable to any one extant species. The *Pristis* material from the Urumaco sequence shows overlap in the number of rostral teeth with *P. pristis*, although at the upper end of the range of this species. Nonetheless, if the whole Venezuelan collection is considered as a single species, DS and DPS values lie outside of the recorded range of *P. pristis* (Table 1). In addition, the posterior rostrum is more slender (SRW/SRL), and the space between the most posterior rostral spine and the chondrocranium is greater (TRL/SRL). In general, the combination of these variables suggests that the fossil material of sawfish from the Urumaco sequence is inconsistent with *P. pristis*, and should be included within the smalltooth sawfish clade.

Within the smalltooth sawfishes, *P. clavata* has a more robust rostrum with a shorter space anterior to the chondrocranium than the Urumaco sequence material (Table 1). Generally high DPS scores and lower rostral spine counts also make referral to *P. zizsron* unlikely. While geographic distribution is most consistent with *P. pectinata*, the high DPS score and slightly narrower rostra of the best-preserved Urumaco material creates uncertainty. *P. zizsron* and *P. pectinata* are sister taxa [103], and there may have been less differentiation in rostral morphospace occupation in Miocene smalltooth sawfish of this lineage, including the Urumaco rostra and *P. atlanticus* from the middle Miocene of Portugal [117].

The total rostral length in *Pristis* is generally considered to be between 1/3 and 1/5 of the total length of the fish [118], and provides a broad size estimate of between 2.4–4 m total length for AMU-CURS-023, 2.1–3.5 m total length for AMU-CURS-235, and 2.7–4.8 m total length (AMU-CURS-237). *P. pectinata* is thought to reach maturity at 2.7–3.6 m, and has a maximum length of over 6.0 m [119]. It seems likely that AMU-CURS-023 and AMU-CURS-237 are adults, and AMU-CURS-235 is either a large juvenile or small adult, but that all are within the size range predicted by extant *Pristis* species, and specifically *P. pectinata*.

Paleoenvironment and paleoecology

The elasmobranch faunistic assemblages from the Urumaco sequence (S1 Table) are distributed across a section more than 2800 m thick (Fig 2). The marine facies in the three members of the Urumaco Formation and in the Chiguaje Member (Codore Formation) have been characterized by Smith et al. [54] as shallow water paleoenvironments based on the study of foraminifera. In addition, most of the living elasmobranch taxa closely related to taxa from the Urumaco sequence prefer coastal habitats, with the exception of some species that show broader preferences (Fig 10). Extant taxa such as *Galeocерdo cuvier*, *Carcharhinus leucas*, *C. limbatus*, *C. obscurus*, *C. plumbeus*, *C. porosus*, *Negaprion brevirostris*, *Sphyrna zygaena*, *Aetobatus narinari* and representatives of the genera *Paragaleus*, *Rhizoprionodon*, *Dasyatis*, *Myliobatis*, *Rhinoptera*, *Rhynchobatus*, and *Pristis* are common inhabitants of marginal marine environments [39, 44–46]. Using the habitat preferences of living elasmobranchs (Fig 10), paleoenvironments suggested in previous studies [27], and the wide range of fossils vertebrates such as reptiles (crocodiles and turtles), mammals (cetaceans and sea cows), and especially bony fishes (S2 Table), that have been described from the whole stratigraphic sequence




Fig 10. Habitat preferences of Urumaco sequence elasmobranch taxa, based on preferences of extant relatives. Light-gray shading indicates extinct taxa.

doi:10.1371/journal.pone.0139230.g010

(including the 14 localities included in the present study) (e.g., [19, 26, 29–32, 36, 120]), we hypothesize that the fossil elasmobranch assemblages from the Urumaco sequence are associated primarily with shallow marine and estuarine environments (Fig 11) (S1 Table).

Of the fourteen localities from the Urumaco sequence with associated elasmobranch assemblages (Fig 1 and S1 Table), we suggest that the El Hatillo Norte locality (Urumaco Formation), which is characterized by a coquinoid limestone rich in marine mollusks (currently under study), and Casa el Jebe (Codore Formation) could be indicative of very shallow marine paleoenvironments. The elasmobranch assemblages at other localities of the Urumaco Formation and those of the Socorro Formation (S1 Table) are characterized by the presence of abundant remains of brackish and freshwater fishes, freshwater and marine turtles and crocodilians, and terrestrial and aquatic/semiaquatic mammals [29–32]. The association of this diverse vertebrate fauna suggests paleoenvironments associated with coastal lagoons and estuaries,



Fig 11. Restoration of diverse sharks and rays in coastal lagoon-estuarine at late Miocene times in Urumaco. (A) Sharpnose shark *Rhizoprionodon* sp., (B) Hammerhead shark *Sphyma* cf. *zygaena*, (C) Bull shark *Carcharhinus leucas*, (D) "Big tooth" *Carcharocles megalodon*, (E) Tiger shark *Galeocerdo cuvier*, (F) Spotted eagle ray *Aetobatus* cf. *narinari*, (G) Eagle ray *Myliobatis* sp., (H) Guitarfish *Rhynchobatus* sp., (I) Sawfish *Pristis* sp., (J) Stingray cf. *Dasyatis*. Artwork by Jorge Gonzalez.

doi:10.1371/journal.pone.0139230.g011

especially the latter environment. Of this fossil aquatic fauna (S2 Table), at least, ten genera of freshwater fishes that includes characins, perciforms, and siluriforms (e.g., [47]), as well the turtle †*Chelus lewisi* Wood, 1976 [121], are phylogenetically close to living groups that occur today in the Orinoco and Amazon drainage systems [20, 30, 47].

The marginal marine paleoenvironments near the mouths of the rivers of the Socorro and Urumaco formations must have been subject to great variability in temperature, pH, sediment load and salinity. This is supported by the abundance in the Urumaco Formation of the foram *Ammonia parkinsoniana* which tolerates a wide range of salinities, and is a dominant taxon in low-salinity lagoons [54]. This suggests that many of the taxa that inhabited these coastal waters may have had broad salinity tolerance (Fig 10). *Galeocerdo cuvier*, *Carcharhinus limbatus*, *C. porosus* and *Negaprion brevirostris* are frequent in estuaries and river mouths, tolerating low levels of salinity, but do not ascend into rivers [43, 44, 46], unlike *Carcharhinus leucas* and *Pristis* species [43, 46, 47]. Batoids such as *Aetobatus narinari*, *Rhinoptera*, *Myliobatis*, and *Dasyatis* are also found in estuaries with low salinity levels [39]; *Rhinoptera bonasus* Mitchell, 1815 [122] even reproduces in the freshwater Maracaibo Lake (Venezuela) [47].

The extinct *Carcharocles megalodon* has been interpreted as a common inhabitant of tropical to warm-temperate coastal-oceanic habitats [2, 23, 24, 123], where it predated primarily cetaceans [81]. Trophic interactions between this large shark (*Carcharocles*) and marine mammals have been inferred based on remains from the early Miocene Cantaure Formation (*Carcharocles chubutensis* Ameghino, 1901 [124]) and early Pliocene Paraganá Formation (*Carcharocles megalodon*), both in Venezuela [15, 17]. However, the presence of *Carcharocles*

in different marine paleoenvironments from the Neogene of Venezuela suggests that this species was a large, transient predator that may have had enough behavioral flexibility to occupy different environments, feeding on fish, turtles, cetaceans and sirenids [15, 17]. In the Urumaco sequence, especially in the Urumaco Formation, we have found isolated teeth of *Carcharocles megalodon* in marine facies and in paleoenvironments interpreted as coastal lagoons and estuaries (S1 Table). In the marine and coastal lagoon facies of the Urumaco Formation, there is no evidence of marine cetaceans; small dolphins probably related to freshwater species that inhabit the Amazon and Orinoco basins are present [30]. The occurrence of *C. megalodon* in the same strata as bony fishes, turtles, sirenids, and crocodiles suggests that these could possibly have been prey for this shark. At present, the presence of *C. megalodon* in freshwater paleoenvironments of the Urumaco sequence cannot be confirmed. Its presence in environments with variable salinity suggests that this taxon had physiological capabilities that allowed it to withstand the variations in salinity in estuarine and possibly river mouth habitats, as do some extant carcharhiniforms [43, 46].

Conclusion

The lithologies of the Urumaco sequence are characterized by substantial variation, indicating the complexity and heterogeneity of these geologic units. Both fossils and the sedimentology document terrestrial and marine facies, including transitional paleoenvironments, and consequently a fauna tolerant to these environments. The elasmobranch fauna from the Urumaco sequence (Socorro-Urumaco-Codore formations), with almost 21 taxa, is associated principally with estuarine coastal lagoon and very shallow marine waters. The presence of elasmobranchs in association with others marine, freshwater and terrestrial vertebrates which provide seemingly contradictory signals for a palaeoenvironmental reconstruction is consistent across the larger stratigraphic sequence. This pattern is not the result of taphonomic processes, but instead proof of mixed coastal marine and fluvial-estuarine hydrographic environments during the Miocene. At the same time, the presence in the Urumaco sequence of abundant aquatic/semiaquatic vertebrates, phylogenetically close to extant groups that occur today in the Orinoco and the Amazon drainage system, support the highly debated hypothesis of a paleo-hydrographic fluvial, lacustrine or wetland complex drainage flowing along the northwestern coast of the Miocene Falcón basin into the proto-Caribbean.

Supporting Information

S1 Appendix. Geographic coordinates.
(DOC)

S2 Appendix. Referred fossil specimens.
(DOC)

S1 Table. Fossil elasmobranchs from the Urumaco sequence and paleoenvironments
(XLS)

S2 Table. Osteichthyan paleodiversity of the Urumaco sequence (Socorro, Urumaco and Codore formations) and the habitat of their living representatives.
(XLS)

Acknowledgments

This work was supported by Swiss National Science Foundation SNF 31003A-149605 to MRSV. The authors wish to especially thank René Kindlimann, Jacques Herman, Carlos

Jaramillo, Helmut Wellendorf, Barbara Oberholzer, and Jorge Ceballos for their generous and important counseling and collaboration; the Instituto del Patrimonio Cultural de Venezuela, for the authorization and permissions to collect fossil material. The Alcaldía Bolivariana de Urumaco, the Universidad Experimental Francisco de Miranda, Museo de Ciencias Naturales de Caracas in Venezuela; Museum of Natural History Vienna, Museum für Naturkunde in Berlin, Museo Nacional de Historia Natural de Santiago in Chile, Museu Paraense Emílio Goeldi, Brazil; Natural History Museum of Basel, Switzerland; Palaeontological Institute and Museum at the University of Zurich, and the Smithsonian Tropical Research Institute, Panama, for their valuable assistance and for access to comparative material. We greatly appreciate comments and suggestions from Dr. Thomas Reinecke and one anonymous reviewer.

Author Contributions

Conceived and designed the experiments: JDCB EM MRSV. Performed the experiments: JDCB EM OA MRSV. Analyzed the data: JDCB EM OA MRSV. Contributed reagents/materials/analysis tools: JDCB EM OA RS MRSV. Wrote the paper: JDCB EM OA MRSV.

References

1. Carpenter KE. The living marine resources of the Western Central Atlantic, Volume 1: Introduction, molluscs, crustaceans, hagfishes, sharks, batoid fishes and chimaeras. FAO Species Identification Guide for Fishery Purposes, Special Publication N° 5. Rome: FAO; 2002.
2. Cappetta H. Handbook of Paleichthyology, Vol. 3E: Chondrichthyes. Mesozoic and Cenozoic Elasmobranchii: Teeth. München: Verlag Dr. Friedrich Pfeil; 2012.
3. Leriche M. Contribution à l'étude des poissons fossiles des pays riverains de la Méditerranée américaine, Venezuela, Trinité, Antilles, Mexique. Mémoires de la Société paléontologique du Suisse. 1938; 61: 1–52.
4. Casier E. Contribution à l'étude des poissons fossiles des Antilles. Mémoire Suisse de Paléontologie. 1958; 74: 1–95.
5. Gillette DD. A marine ichthyofauna from the Miocene of Panama, and the Tertiary Caribbean faunal province. *Journal of Vertebrate Paleontology*. 1984; 4: 172–186. doi: [10.1080/02724634.1984.10012001](https://doi.org/10.1080/02724634.1984.10012001)
6. Kruckow T, Thies D. Die Neoselachier der Paleokaribik (Pisces: Elasmobranchii). *Courier Forschungsinstitut Senckenberg*. 1990; 119: 1–102.
7. Purdy RW, Donovan SK, Pickerill RK, Dixon HL. Fish teeth from the Pleistocene of Jamaica. *Journal of Vertebrate Paleontology*. 1996; 16: 165–167. doi: [10.1080/02724634.1996.10011294](https://doi.org/10.1080/02724634.1996.10011294)
8. Iturralde-Vinent M, Hubbell G, Rojas R. Catalogue of Cuban fossil Elasmobranchii (Paleocene to Pliocene) and paleogeographic implications of their lower to middle Miocene occurrence. *Boletín de la Sociedad Jamaicana de Geología*. 1996; 31: 7–21.
9. Iturralde-Vinent M, Laurito C, Rojas R, Gutiérrez R. Myliobatidae (Elasmobranchii: Batomorphii) del Terciario de Cuba. *Revista de la Sociedad Mexicana de Paleontología*. 1998; 8: 135–145.
10. Laurito C. Los seláceos fósiles de la localidad de Alto Guayacán (y otros ictiolitos asociados), Mioceno superior-Plioceno inferior de la Formación Uscari, provincia de Limón, Costa Rica. San José: Laurito Ceditores; 1999.
11. Laurito C. Ictiofauna de la Formación Punta Judas, Mioceno Medio, Provincia de Punta Arenas, Costa Rica. *Brenesia*. 2004; 62: 57–74.
12. Donovan SK, Gunter GC. Fossil sharks from Jamaica. *Bulletin of the Mizunami Fossil Museum*. 2001; 28: 211–215.
13. Nieves-Rivera AM, Ruizyantin M, Gottfried MD. New Record of the Lamnid Shark *Carcharodon megalodon* from the Middle Miocene of Puerto Rico. *Caribbean Journal of Science*. 2003; 39: 223–227.
14. Underwood C, Simon M. Sharks, bony fishes and endodontal borings from the Miocene Montpelier Formation (White Limestone) of Jamaica. *Cainozoic Research*. 2004; 3: 157–165.
15. Aguilera O, Rodríguez de Aguilera D. Giant-toothed white sharks and wide toothed mako (Lamnidae) from the Venezuela Neogene: their role in the Caribbean, shallow-water fish assemblage. *Caribbean Journal of Science*. 2004; 40 (3): 368–382.

16. Portell RW, Hubbell G, Donovan SK, Green JL. Miocene sharks in the Kendeace and Grand Bay formations of Carriacou, The Grenadines, Lesser Antilles. *Caribbean Journal of Science*. 2008; 44 (3): 279–286.
17. Aguilera O, García L, Cozzuol M. Giant-toothed white sharks and cetacean trophic interaction from the Pliocene Caribbean Paraguaná Formation. *Paläontologische Zeitschrift*. 2008; 82: 204–208. doi: [10.1007/BF02988410](https://doi.org/10.1007/BF02988410)
18. Aguilera O, Ramos MI, Paes E, Costa S, Sánchez-Villagra M. The Neogene Tropical America fish assemblage and the palaeobiogeography of the Caribbean region. *Swiss Journal of Palaeontology*. 2011; 130 (2): 217–240. doi: [10.1007/s13358-011-0020-9](https://doi.org/10.1007/s13358-011-0020-9)
19. Aguilera O. *Peces fósiles del Caribe de Venezuela*. Washington, DC: Gorham Printing; 2010.
20. Aguilera O, Lundberg JG. Venezuelan Caribbean and Orinocoan Neogene fish. In: Sánchez-Villagra MR, Aguilera O, Carlini F, editors. *Urumaco and Venezuelan Paleontology*. Indiana: Indiana Press University; 2010. pp. 129–152.
21. Pimiento C, Ehret DJ, MacFadden BJ, Hubbell G. Ancient Nursery Area for the Extinct Giant Shark *Megalodon* from the Miocene of Panama. *PLoS ONE*. 2010; 5: e10552. doi: [10.1371/journal.pone.0010552](https://doi.org/10.1371/journal.pone.0010552) PMID: 20479893
22. Pimiento C, Gonzales G, Hendy A, Jaramillo C, MacFadden B, Montes C, et al. Early Miocene chondrichthyes from the Culebra Formation, Panama: a window into marine vertebrate faunas before closure the Central American Seaway. *Journal of South American Earth Sciences*. 2013; 42: 159–170. doi: [10.1016/j.jsames.2012.11.005](https://doi.org/10.1016/j.jsames.2012.11.005)
23. Pimiento C, González-Barba G, Ehret DJ, Hendy AJW, MacFadden BJ, Jaramillo C. Sharks and rays (Chondrichthyes, Elasmobranchii) from the Late Miocene Gatun Formation of Panama. *Journal of Paleontology*. 2013; 87: 755–774. doi: <http://dx.doi.org/10.1666/12-117>
24. Carrillo-Briceño JD, De Gracia C, Pimiento C, Aguilera OA, Kindlimann R, Santamarina P, Jaramillo C. A New Late Miocene Chondrichthyan Assemblage from the Chagres Formation, Panama. *Journal of South American Earth Science*. 2015; 60: 56–70. doi: [10.1016/j.jsames.2015.02.001](https://doi.org/10.1016/j.jsames.2015.02.001)
25. Coates AG, Stallard RF. How old is the Isthmus of Panama? *Bulletin of Marine Sciences*. 2013; 89 (4): 801–813. doi: [10.5343/bms.2012.1076](https://doi.org/10.5343/bms.2012.1076)
26. Aguilera O. *Tesoros Paleontológicos de Venezuela: Urumaco, Patrimonio Natural y Cultural de la Humanidad*. Caracas: Editorial Arte; 2004.
27. Quiroz L, Jaramillo C. Stratigraphy and sedimentary environments of Miocene shallow to marginal marine deposits in the Urumaco Trough, Falcon Basin, western Venezuela. In: Sánchez-Villagra MR, Aguilera O, Carlini F, editors. *Urumaco and Venezuelan Paleontology*. Indiana: Indiana Press University; 2010. pp. 153–122.
28. Díaz de Gamero ML. El Mioceno Temprano y Medio de Falcón septentrional. *GEOS*. 1989; 29: 25–35.
29. Sánchez-Villagra MR, Aguilera OA. Neogene vertebrates from Urumaco, Falcón State, Venezuela: diversity and significance. *Journal of Systematic Palaeontology*. 2006; 4: 213–2. doi: [10.1017/S1477201906001829](https://doi.org/10.1017/S1477201906001829)
30. Sánchez-Villagra MR, Aguilera O, Carlini AA. *Urumaco and Venezuelan Paleontology*. The fossil record of the Northern Neotropics. Indiana: Indiana Press University; 2010.
31. Aguilera O, Lundberg J, Birindelli J, Sabaj Pérez M, Jaramillo C, Sánchez-Villagra MR. Palaeontological evidence for the last temporal occurrence of the ancient western Amazonian River outflow into the Caribbean. *PLoS ONE*. 2013; 8(9): e76202. doi: [10.1371/journal.pone.0076202](https://doi.org/10.1371/journal.pone.0076202) PMID: 24098778
32. Scheyer TM, Aguilera OA, Delfino M, Fortier DC, Carlini AA, Sánchez R, et al. Crocodylian diversity peak and extinction in the late Cenozoic of the northern Neotropics. *Nature Communications*. 2013; 4: 1907. doi: [10.1038/ncomms2940](https://doi.org/10.1038/ncomms2940) PMID: 23695701
33. Rod E. Notes on the shifting course of the ancient Rio Orinoco from late Cretaceous to Oligocene time. *GEOS*. 1981; 26: 54–56.
34. Hoon C, Guerrero J, Sarmiento GA, Lorente MA. Andean tectonics as a cause for changing drainage patterns in Miocene northern South America. *Geology*. 1995; 23: 237–240. doi: [10.1130/0091-7613\(1995\)023<0237:ATAACF>2.3.CO;2](https://doi.org/10.1130/0091-7613(1995)023<0237:ATAACF>2.3.CO;2)
35. Díaz de Gamero ML. The changing course of the Orinoco River during the Neogene: a review. *Palaeogeography, Palaeoclimatology, Palaeoecology*. 1996; 123: 385–402. doi: [10.1016/0031-0182\(96\)00115-0](https://doi.org/10.1016/0031-0182(96)00115-0)
36. Sabaj Pérez M, Aguilera O, Lundberg JG. Fossil catfishes of the families Doradidae and Pimelodidae (Teleostei: Siluriformes) from the Miocene Urumaco Formation of Venezuela. *Proceedings of the Academy of Natural Sciences of Philadelphia*. 2007; 156: 157–194. doi: [10.1635/0097-3157\(2007\)156\[157:FCOTFD\]2.0.CO;2](https://doi.org/10.1635/0097-3157(2007)156[157:FCOTFD]2.0.CO;2)

37. Johnson K, Sánchez-Villagra M, Aguilera O. The Oligocene–Miocene transition on coral reefs in the Falcon basin (NW Venezuela). *Palaios*. 2009; 24: 59–69. doi: [10.2110/palo.2008.p08-004r](https://doi.org/10.2110/palo.2008.p08-004r)
38. Lundberg JG, Aguilera O. The late Miocene *Phractocephalus* catfish (Siluriformes: Pimelodidae) from Urumaco, Venezuela: additional specimens and reinterpretation as a distinct species. *Neotropical Ichthyology*. 2003; 1: 97–109. doi: [10.1590/S1679-62252003000200004](https://doi.org/10.1590/S1679-62252003000200004)
39. Compagno LJV. Checklist of living Chondrichthyes. In: Hamlett WC editor. Reproductive biology and phylogeny of Chondrichthyes (sharks, batoids and chimaeras). Enfield New Hampshire: Science Publishers, Inc; 2005. pp. 503–548.
40. Jordan DS, Hannibal H. Fossil sharks and rays of the Pacific Slope of North America. *Bulletin of the Southern California Academy of Sciences*. 1923; 22: 27–63.
41. Wueringer BE, Squire LJ, Collin SP. The biology of extinct and extant sawfish (Batoidea: Sclerorhynchidae and Pristidae). *Reviews in Fish Biology and Fisheries*. 2009; 19: 445–464. doi: [10.1007/s11160-009-9112-7](https://doi.org/10.1007/s11160-009-9112-7)
42. Compagno LJV. FAO Species Catalogue. Vol. 4: Sharks of the World. An annotated and illustrated catalogue of shark species known to date. Part 1: Hexanchiformes to Lamniformes. Rome: Food and Agriculture Organization of the United Nations, 1984.
43. Compagno LJV. FAO Species Catalogue. Vol. 4: Sharks of the World. An annotated and illustrated catalogue of shark species known to date. Part 2: Carcharhiniformes. Rome: Food and Agriculture Organization of the United Nations; 1984.
44. Compagno LJV, Dando VM, Flower S. Sharks of the World. New Jersey: Princeton University Press; 2005.
45. Musick JA, Harbin MM, Compagno LJV. Historical zoogeography of the Selachii. In: Carrier JC, Musick JA, Heithaus MR, editors. *Biology of Sharks and their Relatives*. Washington: CRC Press; 2004. pp. 33–78.
46. Voigt M, Weber D. Field guide for sharks of the genus *Carcharhinus*. München: Verlag Dr. Friedrich Pfeil; 2011.
47. Lundberg JG, Sabaj Perez MH, Dahdul WM, Aguilera OA. The Amazonian Neogene fish fauna. In: Hoorn C, Wesselingh F, editors. *Amazonia: landscape and species evolution. A look into the past*. Chichester, UK: Wiley-Blackwell; 2010. Pp. 281–301.
48. Aguilera O, Marceniuk P. *Aspistor verumquadriscutis*, a new fossil species of sea catfishes (Siluriformes; Ariidae) from the upper Miocene of Venezuela. *Swiss Journal of Palaeontology*. 2012; 131:265–274. doi: [10.1007/s13358-012-0043-x](https://doi.org/10.1007/s13358-012-0043-x)
49. Whitty JM, Phillips NM, Thorburn DC, Simpfendorfer CA, Field I, Peverell SC, et al. Utility of rostra in the identification of Australian sawfishes (Chondrichthyes: Pristidae). *Aquatic Conservation: Marine and Freshwater Ecosystems*. 2014; 24: 791–804. doi: [10.1002/aqc.2398](https://doi.org/10.1002/aqc.2398)
50. Díaz de Gamero ML, Linares OJ. Estratigrafía y paleontología de la Formación Urumaco, del Mioceno Tardío de Falcón Noroccidental. VII Congreso Geológico Venezolano, Memorias. 1989; 1: 419–438.
51. Hambalek N, Rull V, De Digiacoimo E, Díaz de Gamero ML. Evolución paleoecológica y paleoambiental de la secuencia del Neógeno en el surco de Urumaco. Estudio palinológico y litológico. *Boletín de la Sociedad Venezolana de Geólogos*. 1994; 191: 7–19.
52. Rey OT. Miembro Chiguaje: Formación La Vela o Formación Codore. *Boletín de la Sociedad Venezolana de Geólogos*. 1994; 1–2: 50–53.
53. Rey OT. Análisis comparativo y correlación de las formaciones Codore y La Vela, estado Falcón. M. Sc. Thesis, Universidad Central de Venezuela, 1990.
54. Smith CJ, Collins LS, Jaramillo C, Quiroz L. Marine paleoenvironment of Miocene-Pliocene formations of north-central Falcón state, Venezuela. *Journal of Foraminiferal Research*. 2010; 40 (3): 266–282. doi: [10.2113/gsjfr.40.3.266](https://doi.org/10.2113/gsjfr.40.3.266)
55. Compagno LJV. Phyletic relationships of living sharks and rays. *American Zoologist*. 1977; 17: 303–322. doi: [10.1093/icb/17.2.303](https://doi.org/10.1093/icb/17.2.303)
56. Compagno LJV. Interrelationships of living elasmobranchs. In: Greenwood PH, Miles RS, Patterson C, editors. *Interrelationships of fishes*. Zoological Journal of the Linnean Society, Supplement 1; 1973. pp. 15–61.
57. Berg LS. A classification of fish-like vertebrates. *Bulletin of the Academy of Sciences of the USSR, Division of Chemical Science*. 1937; 1937: 1277–1280.
58. Glikman LS. Sharks of the Paleogene and their stratigraphic significance. Moscow: Nauka Press; 1964.
59. Agassiz L. Recherches sur les poissons fossiles. Neuchatel: Petitpierre; 1833–1843.

60. Hasse JCF. Das natürliche System der Elasmobranchier auf Grundlage des Baues und der Entwicklung ihrer Wirbelsäule. Eine morphologische und palaontologische Studie. Allgemeiner Theil. Jena: Gustav Fischer Verlag; 1879.
61. Budker P. 1935. Description d'un genre nouveau de la famille des Carcharhinides. Bulletin du Museum D'Histoire Naturelle, Paris. 1935; 7 (2): 107–112.
62. Carrillo-Briceño JD, Aguilera O, Rodríguez F. Fossil Chondrichthyes from the Central Eastern Pacific Ocean and their Paleooceanographic Significance. Journal of South American Earth Sciences. 2014; 51: 76–90. doi: [10.1016/j.jsames.2014.01.001](https://doi.org/10.1016/j.jsames.2014.01.001)
63. Jordan DS, Evermann BW. The fishes of North and Middle America: a descriptive catalogue of the species of fish-like vertebrates found in the waters of North America, north of the Isthmus of Panama. Part I. Bulletin of the United States National Museum. 1896; 47: 1–1240.
64. Müller J, Henle J. Gattungen der Haifische und Rochen nach einer von ihm mit Hrn. Henle unternommenen gemeinschaftlichen Arbeit über die Naturgeschichte der Knorpelfische. Akademie der Wissenschaften zu Berlin. 1837; 2: 111–118.
65. Peron F, Lesueur CA. Description of a *Squalus*, of a very large size, which was taken on the coast of New Jersey. Journal of the Philadelphia Academy of Natural Sciences., 1822; 2: 343–352.
66. Whitley GP. Additions to the check-list of the fishes of New South Wales. No. 2. Australian Zoologist. 1929; 5 (4): 353–357.
67. Blainville HMD. Prodrome d'une nouvelle distribution systematique du regne animal. Bulletin de la Société Philomathique de Paris. 1816; 8: 105–112, 121–124.
68. Bleeker PR. Enumeratio specierum piscium hucusque in Archipelago indico observatarum. Acta Societatis Scientiarum Indo-Neerlandae. 1859; 6: 1–276.
69. Valenciennes A. In Müller J, Henle FGJ. Systematische Beschreibung der Plagiostomen, Berlin: Veit (2); 1839. Pp. 39–102.
70. Whitley GP. Ichthyological descriptions and notes. Proceedings of the Linnean Society of New South Wales. 1943; 68: 114–144.
71. Ranzani C. De novis speciebus piscium. Dissertatio prima. Novi Commentarii Academiae Scientiarum Instituti Bononiensis. 1839; 4: 65–83.
72. Pietschmann V. Fische des Wiesbadener Museums. Jahrbücher des Nassauischen Vereins für Naturkunde. 1913; 66: 170–201.
73. Poey F. Synopsis piscium cubensium. Catálogo Razonado de los peces de la isla de Cuba. Repertorio Físico-Natural de la Isla de Cuba. 1868; 2: 279–484.
74. Müller J, Henle J. Systematische Beschreibung der Plagiostomen. Berlin: Veit; 1839.
75. Lesueur CA. Description of several new species of North American fishes. Journal of the Philadelphia Academy of Natural Sciences. 1818; 1: 222–235.
76. Nardo JD. Prodromus observationum et disquisitionum ichthyologiae Adriaticae. Oken's Isis. 1827; 20: 472–631.
77. Gill T. Arrangement of the families of fishes, or classes Pisces, Marsupiobranchii, and Leptocardii. Smithsonian Institution Miscellaneous Collection. 1872; 247: 1–49.
78. Rafinesque CS. Caratteri di alcuni nuovi generi e nuove specie di Animali e Piante della Sicilia con varie osservazioni sopra i medesimi. Palermo: Sanfilippo; 1810.
79. Linnaeus C. Systema naturae. 10th Edition. Stockholm: Larentii Salvii; 1758.
80. Gilbert CR. A revision of the hammerhead sharks (family Sphymidae). Proceedings of the United States National Museum, Smithsonian Institution. 1967; 119: 1–88.
81. Purdy R, Clellan JHM, Schneider VP, Applegate SP, Meyer R, Slaughter R. The Neogene sharks, rays and bony fishes from Lee Creek Mine, Aurora, North Carolina. Smithsonian Contributions to Paleobiology. 2001; 90: 71–202.
82. Cappetta H. Modification du statut générique de quelques espèces de sélaciens crétacés et tertiaires. Palaeovertebrata. 1980; 10: 29–42.
83. Jordan DS. Description of two new species of fishes from South America. Proceedings of the Academy of Natural Sciences of Philadelphia. 1888; 39:387–388.
84. Bonaparte CL. Selachorum tabula analytica. Nuovi Annali della Science Naturali Bologna. 1838; 1: 195–214.
85. Euphrasen BA. *Raja (Narinari)*. Kongliga Vetenskaps Akademiens nya Handlingar, Stockholm. 1790; 11: 217–219.

86. Hovestadt DC, Hovestadt-Euler M. Generic Assessment and Reallocation of Cenozoic Myliobatinae based on new information of tooth, tooth plate and caudal spine morphology of extant taxa. *Palaeontos*. 2013; 24, 1–66.
87. Antunes MT, Balbino AC. Latest Miocene Myliobatids (Batoidei, Selachii) from the Alvalade Basin, Portugal. *Cainozoic Research*. 2006; 4: 41–49.
88. Reinecke T, Louwye S, Havekost U, Moths H. The elasmobranch fauna of the late Burdigalian, Miocene, at Werder-Uesen, Lower Saxony, Germany, and its relationships with Early Miocene faunas in the North Atlantic, Central Paratethys and Mediterranean. *Palaeontos*. 2011; 20: 1–170.
89. Bor TJ, Reinecke T, Verschueren S. Miocene Chondrichthyes from Miste, The Netherlands. *Palaeontos*. 2012; 21: 1–136.
90. Cuvier GLCFD. Le règne animal distribué d'après son organisation pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée. Les reptiles, les poissons, les mollusques et les annélides. Paris: Deterville; 1816.
91. Nishida K. Phylogeny of the Suborder Myliobatoidei. *Memoirs of the Faculty of Zoology, Hokkaido University*. 1990; 37: 1–108.
92. Cuvier GLCFD. Le règne animal, distribué d'après son organisation, pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée. Edition 2. Paris: Chez Deéerville; 1829.
93. Garman S. The Plagiostomia (Sharks, Skates and Rays). *Memoirs of the Museum of Comparative Zoology at Harvard College*. 1913; 36: 1–528.
94. Linck HF. Versuch einer Eintheilung der Fische nach den Zähnen. *Magazin für das Neueste aus der Physik und Naturgeschichte*. 1790; 6 (3): 28–38.
95. White EI, Moy-Thomas JA. Notes on the nomenclature of fossil fishes. Part. III: Homonyms M-Z. *Annals and Magazine of Natural History, (Series 11)*. 1941; 7: 395–400.
96. Latham J. An essay on the various species of Sawfish. *Transactions of the Linnean Society of London*. 1794; 2: 273–282.
97. Herman J, Hovestadt-Euler M, Hovestadt DC. Contributions to the study of the comparative morphology of teeth and other relevant ichthyodorulites in living superaspecific taxa of Chondrichthyan fishes. Part B: Batomorphii No. 2: Order: Rajiformes—Suborder: Pristoidei—Family: Pristidae—Genera: *Anoxypristis* and *Pristis*; No. 3: Suborder: Rajoidei—Superfamily: Rhinobatoidea—Families: Rhinidae—Genera: *Rhina* and *Rhynchobatus* and Rhinobatidae—Genera: *Aptychotrema*, *Platyrhina*, *Platyrhinoidis*, *Rhinobatos*, *Trygonorrhina*, *Zanobatus* and *Zapteryx*. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Biologie*. 1997; 67: 107–162.
98. Delfortrie E. Les broyeur du Tertiaire aquitanien. *Actes de la Société linnéenne de Bordeaux*. 1871; 28: 213–236.
99. Cappetta H. Les Sélaciens du Miocène de la région de Montpellier. *Palaeovertebrata, Mémoire Extraordinaire*. 1970; 1–139.
100. Reif WE. Evolution of dermal skeleton and dentition in vertebrates: the odontode regulation theory. *Evolutionary Biology*. 1982; 15: 287–368. doi: [10.1007/978-1-4615-6968-8_7](https://doi.org/10.1007/978-1-4615-6968-8_7)
101. Wiley TR, Simpfendorfer CA, Faria VV, McDavitt MT. Range, sexual dimorphism and bilateral asymmetry of rostral tooth counts in the smalltooth sawfish *Pristis pectinata* Latham (Chondrichthyes: Pristidae) of the southeastern United States. *Zootaxa*. 2008; 1810: 51–59.
102. Bleeker PR. Vijfde bijdrage tot de kennis der ichthyologische fauna van Borneo, met beschrijving van eenige nieuwe soorten van zoetwatervisschen. *Natuurkundig Tijdschrift voor Nederlandsch Indië*. 1851; 2: 415–442.
103. Faria VV, McDavitt MT, Charvet P, Wiley TR, Simpfendorfer CA, Naylor GJP. Species delineation and global population structure of critically endangered sawfishes (Pristidae). *Zoological Journal of the Linnean Society*. 2013; 167: 136–164. doi: [10.1111/j.1096-3642.2012.00872.x](https://doi.org/10.1111/j.1096-3642.2012.00872.x)
104. Duméril A. Histoire naturelle des poissons ou ichthyologie générale. Tome Premier. Elasmobranchés. Plagiostomes et Holocéphales ou Chimères. Paris: Librairie Encyclopédique de Roret; 1865.
105. Hoffmann L. Zur Kenntnis des Neurocraniums der Pristiden und Pristiophoriden. *Zoologische Jahrbucher. Abteilung für Anatomie und Ontogenie der Tiere*. 1912; 33: 237–360.
106. Kirkland JJ, Aguillón-Martínez MC. *Schizorhiza*: a unique sawfish paradigm from the Difunta Group, Coahuila, Mexico. *Revista Mexicana de Ciencias Geológicas*. 2002; 19: 16–24.
107. Cicimurri DJ. A partial rostrum of the sawfish *Pristis lathamii* Galeotti, 1837, from the Eocene of South Carolina. *Journal of Paleontology*. 2007; 81: 597–601. doi: [10.1666/05086.1](https://doi.org/10.1666/05086.1)
108. Casier E. Contributions à l'étude des poissons fossiles de la Belgique. VIII. Les Pristidés éocènes. *Institut royal des Sciences naturelles de Belgique*. 1949; 25: 1–52.

109. Wueringer BE, Peverell SC, Seymour J, Squire LJ, Collin SP. Sensory systems in sawfishes. 2. The lateral line. *Brain, Behavior and Evolution*. 2011; 78: 150–161. doi: [10.1159/000329518](https://doi.org/10.1159/000329518) PMID: [21829005](https://pubmed.ncbi.nlm.nih.gov/21829005/)
110. Wueringer BE, Peverell SC, Seymour J, Squire LJ, Kajiura SM, Collin SP. Sensory systems in sawfishes. 1. The ampullae of Lorenzini. *Brain, Behavior and Evolution*. 2011; 78: 139–149. doi: [10.1159/000329515](https://doi.org/10.1159/000329515) PMID: [21829004](https://pubmed.ncbi.nlm.nih.gov/21829004/)
111. Gaman S. New Plagiostomia. *Bulletin of the Museum of Comparative Zoology at Harvard College*. 1906; 46 (11): 203–208.
112. Nelson JS. *Fishes of the World* (Fourth Edition). New York: J. Wiley; 2006.
113. Miller WA. Observations on the developing rostrum and rostral teeth of sawfish: *Pristis perotteti* and *P. cuspidatus*. *Copeia*. 1974; 74: 311–318.
114. Baughman JL. Notes on sawfish *Pristis perotteti* Müller and Henle, not previously reported from the waters of the United States. *Copeia*. 1943; 1943: 43–48.
115. Schultz LP. A further contribution to the ichthyology of Venezuela. *Proceedings of the United States National Museum*. 1949; 99: 1–211.
116. Robillard M, Séret B. Cultural importance and decline of sawfish (Pristidae) populations in West Africa. *Cybum*. 2006; 30: 23–30.
117. Zbyszewski G. Découverte d'un rostre de grand *Pristis* dans l'Helvétien de Lisbonne. *Boletim da Sociedade Geológica de Portugal*. 1947; 6: 237–242.
118. de Carvalho MR, McEachran JD. Family Pristidae (Sawfishes). In: Reis E, Ferraris CJJ, Kullander SO, editors. *Check list of the freshwater fishes of South and Central America*. Porto Alegre: Edipucrs; 2003. Pp. 17–20.
119. Simpfendorfer CA, Poulakis GR, O'Donnell PM, Wiley TR. Growth rates of juvenile smalltooth sawfish *Pristis pectinata* Latham in the western Atlantic. *Journal of Fish Biology*. 2008; 72:711–723. doi: [10.1111/j.1095-8649.2007.01764.x](https://doi.org/10.1111/j.1095-8649.2007.01764.x)
120. Aguilera O, Rodríguez de Aguilera D. Goliath grouper (Pisces: Serranidae) from the Upper Miocene Urumaco Formation, Venezuela. *Journal of Paleontology*. 2004; 78: 1202–1206. doi: [10.1666/0022-3360\(2004\)078<1202:GGPSFT>2.0.CO;2](https://doi.org/10.1666/0022-3360(2004)078<1202:GGPSFT>2.0.CO;2)
121. Wood RC. Two new species of *Chelus* (Testudines: Pleurodira) from the Late Tertiary of northern South America. *Breviora*. 1976; 435: 1–26.
122. Mitchill SL. The fishes of New York described and arranged. *Transactions of the Literary and Philosophical Society of New York*. 1815; 1: 355–492.
123. Pimiento C, Balk MA. Body-size trends of the extinct giant shark *Carcharocles megalodon*: a deep-time perspective on marine apex predators. *Paleobiology*. 2014; 41(3): 479–490.
124. Ameghino F. L'âge des formations sédimentaires de Patagonie. *Anales de la Sociedad Científica Argentina*. 1901; 51: 20–39, 65–91.

S1 Appendix. Geographic coordinates.

Socorro Formation (middle Miocene)

-Cerro Overo (11°11'14"N, 70° 8'44"W)

- Cerro Alto (11°12'30"N, 70° 8'12"W)

Urumaco Formation (late Miocene)

-Sur Quebrada Bejucal (11°11'18"N, 70°15'3"W)

- Quebrada Bejucal (11°11'22"N, 70°15'20"W)

- Puente Río Urumaco (11°12'24"N, 70°15'00"W)

-Domo de Agua Blanca (11°13'25"N, 70°14'5"W)

-El Hatillo (11°14'7"N, 70°15'3"W)

-El Hatillo Norte (11°14'40"N, 70°14'44"W)

-El Mamón (11°14'1"N, 70°16'11"W)

-El Mamón Norte (11°14'27"N, 70°16'2"W)

-El Picache (11°14'26"N, 70°13'27"W)

-Corralito (11°14'57"N, 70°16'9"W)

-Tío Gregorio (11°14'42"N, 70°18'19"W)

Codore Formation (late Miocene–Pliocene)

-Casa El Jebe (11°15'54"N, 70°16'35"W).

S2 Appendix. Referred fossil specimens.

-Socorro Formation: *Carcharhinus* spp. [5 teeth (AMU-CURS-630)]; *Negaprion brevirostris* [1 tooth (AMU-CURS-630)]; *Aetobatus* cf. *narinari* [1 teeth (AMU-CURS-323)]; *Myliobatis* sp. [11 teeth (AMU-CURS-322)]; *Rhinoptera* sp. [2 teeth (AMU-CURS-)]; *Pristis* sp. [2 rostra (AMU-CURS-241 and AMU-CURS-639)].

-Urumaco Formation: *Carcharocles megalodon* [15 teeth (AMU-CURS-117, 324, 330, 338, 339, 455 and 605; UNEFM-CIAAP-359, 360, 1225, 1226 and 1292; UNEFM-PF-338)]; *Hemipristis serra* [16 teeth (AMU-CURS-331, 332-337 and 362)]; *Paragaleus* sp. [1 tooth (AMU-CURS-640)]; *Galeocерdo cuvier* [2 teeth (AMU-CURS-341 and UNEFM-PF-408)]; *Rhizoprionodon* sp. [11 teeth (AMU-CURS-478, 479-481, 485, 497 and 498)]; *Carcharhinus caquetius* sp. nov. [5 teeth (AMU-CURS-477 and 499)]; *Carcharhinus leucas* [23 teeth (AMU-CURS-355, 360, 363, 364, 368, 458-462)]; *Carcharhinus limbatus* [2 teeth (AMU-CURS-456 and 457)]; *Carcharhinus obscurus* [20 teeth (AMU-CURS-356, 361, 367, 369-371, 463-467)]; *Carcharhinus porosus* [4 teeth (AMU-CURS-472, 473 and 590)]; *Carcharhinus* spp. [77 teeth (AMU-CURS-60, 358, 359, 365, 366, 373-375, 471, 486 and 629)]; *Negaprion brevirostris* [20 teeth (AMU-CURS-342, 343-349, 372, 468-470, 616, 617 and 1039)]; *Sphyrna* cf. *zygaena* [2 teeth (AMU-CURS-474; AMU-CURS-474)]; *Sphyrna* sp. [1 tooth (AMU-CURS-477)]; *Dasyatis* spp. [9 teeth (AMU-CURS-493, 495, 496 and 589)]; *Dasyatidae* indet. [1 tooth (AMU-CURS-591)]; *Aetobatus* cf. *narinari* [39 teeth (AMU-CURS-272, 273, 275, 276, 308, 311, 313, 318, 329, 488, and 490; UNEFM-CIAAP-1282 and 1288)]; *Myliobatis* sp. [692 teeth (AMU-CURS-59, 61, 260, 262-269, 271, 274, 277, 279, 281-283, 285, 287, 289, 303-307, 312, 314-316, 319, 326, 434, 435, 487, 615 and 618)]; *Rhinoptera* sp. [31 teeth (AMU-CURS-270, 278, 284, 286, 288, 309, 310, 317, 320, 328, 489 and 491)]; *Myliobatiformes* indet. [5 caudal spines (AMU-CURS-290 and 492)]; *Rhynchobatus* sp. [3 teeth (AMU-CURS-482, 483 and 628)]; *Pristis* sp. [22 rostra (AMU-CURS-23, 40, 41, 43, 45, 102, 107, 235-240, 242, 245, 251 and 376; MCNC-45-72-V, 93-72-V, 99-72-V, 154-72-V and 155-72-V), 41 rostral spines (AMU-CURS-242-244, 246,-251, 327 and 1065), and 4 bucal teeth (AMU-CURS-484)].

-Codore Formation: *Carcharocles megalodon* [3 teeth (AMU-CURS-599; UNEFM-PF-351)]; *Hemipristis serra* [3 teeth (AMU-CURS-623)]; *Galeocерdo cuvier* [2 teeth (AMU-CURS-625)]; *Rhizoprionodon* sp. [3 teeth (AMU-CURS-635)]; *Carcharhinus leucas* [5 teeth (AMU-CURS-622)]; *Carcharhinus plumbeus* [4 teeth (AMU-CURS-624)]; *Carcharhinus porosus* [4 teeth (AMU-CURS-632)]; *Negaprion brevirostris* [4 teeth (AMU-CURS-626)]; *Dasyatis* spp. [7 teeth (AMU-CURS-636)]; *Aetobatus* cf. *narinari* [3 fragmented dental plates (AMU-CURS-598), and 14 teeth (AMU-CURS-614)]; *Myliobatis* sp. [95 teeth (AMU-CURS-619)]; *Rhinoptera* sp. [17 teeth (AMU-CURS-621)]; *Myliobatiformes* indet. [1 caudal spine (AMU-CURS-634)]; cf. *Rhynchobatus* [2 teeth (AMU-CURS-638)]; *Pristis* sp. [1 rostral spine (AMU-CURS-620)].

S1 Table. Fossil elasmobranch from the Urumaco sequence and paleoenvironments.															
Taxon	Socorro Formation			Urumaco Formation										Codore Fm	
	Middle Mb	Upper Mb		Lower Member		Middle Member		Upper Member						Chiguaje Member	Casa El Jebe
	Cerro Overo	Cerro Alto	Quebrada Bejucal	Sur Quebrada Bejucal	Quebrada Bejucal	Puente Río Urumaco	Domo de Agua Blanca	El Hatillo	El Hatillo Norte	El Mamón	El Mamón Norte	El Picache	Corralito	Tío Gregorio	
<i>†Carcharocles megalodon</i>			X	X	X	X	X	X	X	X		X		X	
<i>†Hemipristis serra</i>			X	X				X	X	X		X	X	X	
<i>Paragaleus</i> sp.									X	X					
<i>Galeocerdo cuvier</i>									X	X				X	
<i>Rhizoprionodon</i> sp.						X		X		X				X	
<i>†Carcharhinus caquetius</i> nov. sp.						X									
<i>Carcharhinus leucas</i>			X	X				X	X	X		X	X	X	
<i>Carcharhinus limbatus</i>								X	X			X			
<i>Carcharhinus obscurus</i>						X		X	X	X		X	X		
<i>Carcharhinus plumbeus</i>														X	
<i>Carcharhinus porosus</i>						X				X				X	
<i>Carcharhinus</i> spp.	X		X	X	X	X		X		X			X	X	
<i>Negaprion brevirostris</i>	X		X	X				X	X	X		X	X	X	
<i>Sphyrna</i> cf. <i>zygaena</i>								X	X	X					
cf. <i>Dasyatis</i>						X		X	X		X			X	
<i>Dasyatidae</i> indet.											X				
<i>Aetobatus</i> cf. <i>narinari</i>		X			X	X		X	X	X		X	X	X	
<i>Myliobatis</i> sp.		X	X	X	X	X		X	X	X		X	X	X	
<i>Rhinoptera</i> sp.		X	X	X	X			X	X			X	X	X	
<i>Myliobatiformes</i> indet.										X				X	
<i>Rhynchobatus</i> spp.			X	X	X			X	X					X	
<i>Pristis</i> sp.	X	X	X	X	X	X		X			X	X	X	X	
Paleoenvironment															
Marine (very shallow waters)								X	X					X	
Coastal lagoon		X			X	X	X	X		X	X				
Estuarine	X	X	X	X	X	X		X		X	X	X	X		
Riverine															

S2 Table. Osteichthyan paleodiversity of the Urumaco sequence (Socorro, Urumaco and Codore formations) and the habitat of their living representatives								
Taxon			Geologic units			Habitat of living representatives		
			Socorro Formation	Urumaco Formation	Codore Formation	Freshwater	Brackish	Marine
Characiformes	Erythrinidae	? <i>Hoplias</i>		X		X		
	Serrasalminidae	<i>Colossoma</i> sp.		X		X		
		<i>Piaractus</i> sp.		X		X		
Chupeiformes	Engraulidae	<i>Anchoa</i> sp.			X		X	X
		<i>Orthopristis</i> aff. <i>ruber</i>			X		X	X
		<i>Cynoscion</i> aff. <i>jamaicensis</i>		X			X	X
	Sciaenidae	<i>Cynoscion</i> sp.		X	X		X	X
		<i>Equetus</i> sp.		X	X			X
		<i>Larimus</i> sp.		X			X	X
		† <i>Larimus gatumensis</i>		X			X	X
		† <i>Microgogonias coatesi</i>		X			X	X
		<i>Nebris</i> aff. <i>Occidentalis</i>		X			X	X
		† <i>Ophioscion lundbergi</i>		X			X	X
		<i>Paralichthys</i> sp.			X		X	X
		† <i>Plagioscion urumacoensis</i>		X		X	X	X
		<i>Epinephelus itajara</i>		X			X	X
	Serranidae	Indt.		X		?	?	?
		<i>Sphyraena</i> sp.			X		X	X
Pleuronectiformes	Cynoglossidae	<i>Amphiarus rugispinis</i>		X			X	X
Siluriformes	Anidae	† <i>Aspistor verumquadriscutis</i>		X		X	X	X
		<i>Aspistor quadriscutis</i>		X		X	X	X
		<i>Bagre marinus</i>		X			X	X
		<i>Cathorops</i> sp.		X		X	X	X
		<i>Notarius kessleri</i>		X			X	X
		<i>Notarius troscheli</i>		X			X	X
		<i>Sciades couma</i>		X		X	X	X
		<i>Sciades dowii</i>		X		X	X	?
		<i>Sciades herbergii</i>		X		X	X	X
		Indt.	X	X		?	?	?
		<i>Doraops</i> cf. <i>zuloagai</i>		X		X		
	Doradidae	† <i>Doras dionae</i>		X		X		
		<i>Rhinodoras</i> cf. <i>thomersoni</i>		X		X		
		<i>Acanthicus</i> sp.		X		X		
	Loricariidae	† <i>Phractocephalus nassi</i>		X		X		
	Pimelodidae	<i>Platysilurus</i> sp.		X		X		

CHAPTER 5

A new early Miocene (Aquitanian) Elasmobranchii assemblage from the La Guajira Peninsula, Colombia



Kitefin shark *Dalatias* cf. *licha*



A NEW EARLY MIOCENE (AQUITANIAN) ELASMOBRANCHII ASSEMBLAGE FROM THE LA GUAJIRA PENINSULA, COLOMBIA

JORGE D. CARRILLO-BRICEÑO¹, THODORIS ARGYRIOU¹, VLADIMIR ZAPATA², RENÉ KINDLIMANN¹, AND CARLOS JARAMILLO³

¹Paleontological Institute and Museum, University of Zurich, Karl-Schmid-Strasse 4, 8006 Zürich, Switzerland. jorge.carrillo@pim.uzh.ch, thodoris.argyriou@pim.uzh.ch, mad_design@gmx.ch

²Ecopetrol S.A Edificio Colgas, Calle 37 N° 8-43 Piso 8 costado occidental, 11001000 Bogotá, Colombia. vladimir.zapata@ecopetrol.com.co

³Smithsonian Tropical Research Institute, Av. Gorgas, Ed. 235, 0843- 03092 Balboa, Ancón, Panamá. jaramilloc@si.edu

Abstract. Recent field expeditions have led to the discovery of a selachian assemblage from the earliest Miocene (Aquitanian) deposits of the Uitpa Formation in the La Guajira Peninsula, Colombia. This elasmobranch assemblage provides a unique glimpse into the Caribbean biodiversity at the onset of the Neogene. The assemblage consists of 13 taxa, of which some are reported from Miocene deposits for the very first time. There are also new records of taxa in the southern Caribbean region. The taxonomic composition of the selachian assemblage was used to conduct a paleoenvironmental and paleobathymetric analysis of the lower Uitpa Formation. The maximum likelihood estimation of paleobathymetry suggests that the lower part of the Uitpa Formation was probably accumulated at a water depth of 100 to 200 m. This indicates a rapid increase in relative sea level or basin deepening, providing new insights into the possible causes of marine biota changes in the Cocosineta Basin during the Oligocene/Miocene transition.

Key words. Neogene. Tropical America. Caribbean. Sharks. Rays. Paleobathymetry. Paleoenvironments.

Resumen. UN NUEVO ENSAMBLE DE ELASMOBRANCHII DEL MIOCENO TEMPRANO (AQUITANIANO) DE LA PENÍNSULA DE LA GUAJIRA, COLOMBIA. Recientes trabajos de campo han llevado al descubrimiento de un conjunto de seláceos procedente del Mioceno temprano (Aquitano) en los depósitos más antiguos de la Formación Uitpa en la Península de La Guajira, Colombia. Este conjunto de elasmobranchios provee una visión única de la paleobiodiversidad del Caribe a inicios del Período Neógeno. El conjunto aquí descrito se compone de 13 taxones, algunos de los cuales son reportados por vez primera para el Mioceno. Esta fauna también incluye nuevos registros para la región sur del Caribe. La composición taxonómica del nuevo ensamble permitió realizar un análisis paleoambiental y paleobatimétrico en la sección más inferior de la Formación Uitpa. La estimación de paleobatimetría usando *maximum likelihood* sugiere que la parte inferior de la Formación Uitpa fue probablemente depositada a profundidades entre los 100 y 200 m. Esto indica un rápido incremento en el nivel relativo del mar o en la profundización de la cuenca, proveyendo nueva información sobre las posibles causas que dieron origen a los cambios en la biota marina de la Cuenca de Cocinetas durante la transición del Oligoceno/Mioceno.

Palabras clave. Neógeno. América Tropical. Caribe. Tiburones. Rayas. Paleobatimetría. Paleoambientes.


THE beginning of the Neogene was a time when world climate was warmer and sea level was higher than at present (Zachos *et al.*, 2001). Large scale geological processes, including the closure of the Central American Seaway and the rise of the Panamanian Isthmus, had not yet been completed (Woodburne, 2010; Montes *et al.*, 2012a,b; Coates and Stallard, 2013; Montes *et al.*, 2015). Large areas of the northern margin of South America were submerged during the early Miocene (see Iturralde-Vinent and MacPhee, 1999 and references therein). This was also the case in the southern Caribbean Cocosineta Basin, where there are widespread early Miocene marine sedimentary deposits that have

yielded a rich record of invertebrate (Becker and Dusenbury, 1958; Lockwood, 1965; Rollins, 1965; Thomas, 1972; Hendy *et al.*, 2015) and vertebrate fossils of which most are chondrichthyans (Lockwood, 1965; Moreno *et al.*, 2015).


Early Miocene marine chondrichthyan faunas from Tropical America are still poorly known with only a few relevant reports from Barbados (Casier, 1958), Brazil (Santos and Travassos, 1960; Santos and Salgado, 1971; Reis, 2005; Costa *et al.*, 2009), Cuba (Iturralde-Vinent *et al.*, 1996), Mexico (Gonzales-Barba and Thies, 2000), Panama (Pimiento *et al.*, 2013a), Trinidad (Leriche, 1938), the Grenadines (Portell *et al.*, 2008) and Venezuela (Leriche, 1938; Sánchez-Villagra *et*

CHAPTER 6

An Early Neogene Elasmobranch fauna from the southern Caribbean (Western Venezuela)



Devil ray †*Mobula fragilis*



An Early Neogene Elasmobranch fauna from the southern Caribbean (Western Venezuela)

Jorge D. Carrillo-Briceño, Orangel A. Aguilera, Carlos De Gracia, Gabriel Aguirre-Fernández, René Kindlimann, and Marcelo R. Sánchez-Villagra

ABSTRACT

The Cantaure Formation (Burdigalian to ?early Langhian) is located in the Falcón Basin, North Western Venezuela, and includes one of the most diverse Neogene teleostean and benthonic invertebrate faunas in Tropical America. The paleoenvironmental preferences of the members of this fauna, as well as published paleogeographic reconstructions, suggest that the Cantaure Formation was deposited in a highly-productive shallow water environment, associated with coastal upwelling. We documented a paleodiversity of 39 shark and ray species, including 15 previously unreported taxa for Venezuela and six for Tropical America. We performed a bathymetric analysis of the fossil assemblage based on the distribution of closely-related extant chondrichthyan relatives of fossil taxa and discuss the ecological role and stratigraphic significance of the latter. Our results support the hypothesis that the Cantaure Formation was deposited in an insular inner-middle shelf environment. The elasmobranch fauna is characterized by a predominance of benthopelagic sharks with piscivorous feeding preferences (e.g., †*Paratodus*, *Galeorhinus*, *Hemipristis*, *Rhizoprionodon*, *Carcharhinus*, *Isogomphodon*, *Negaprion*, †*Physogaleus* and *Sphyrna*) followed by durophagous/carcitrophic feeders (e.g., *Heterodontus*, *Nebrius*, *Mustelus*, *Rhynchobatus*, *Pristis*, *Dasyatis*, cf. *Pteroplatytrygon*, cf. *Taeniurops*, *Aetobatus*, *Aetomylaeus* and *Rhinoptera*). Filter (e.g., *Mobula* and †*Plinthiscus*), eurytrophic/sarcophagous (e.g., †*Carcharocles* and *Galeocerdo*) and teuthitrophic (e.g., *Alopias*) feeder species were also found. Teeth of *Carcharocles megalodon* found in Burdigalian sediments of the Cantaure Formation support the presence of this species already in the early Miocene. Some taxa (*Nebrius*, *Carcharhinus* cf. *C. macroti* and *Rhynchobatus*) are absent from the extant Caribbean and Western Atlantic fauna, but were present in the region before the closure of the Central American Seaway.

Jorge D. Carrillo-Briceño. Paleontological Institute and Museum, University of Zurich, Karl-Schmid-Strasse 4, Zürich, 8006, Switzerland. jorge.carrillo@pim.uzh.ch

Orangel A. Aguilera. Universidade Federal Fluminense, Instituto de Biologia, Departamento de Biologia Marinha, Programa de Pós-graduação em Biologia Marinha. Campus do Valonguinho, Outeiro São João Batista, s/nº. 24020-141, Niterói, RJ, Brasil. orangel.aguilera@gmail.com

Carlos De Gracia. Institute of Geology and Paleontology, Faculty of Science, Charles University in Prague,

Carrillo-Briceño, Jorge D., Aguilera, Orangel A., De Gracia, Carlos, Aguirre-Fernández, Gabriel, Kindlimann, René, and Sánchez-Villagra, Marcelo R. 2016. An Early Neogene Elasmobranch fauna from the southern Caribbean (Western Venezuela). *Palaeontologia Electronica* 19.2.27A: 1-32

palaeo-electronica.org/content/2016/1541-early-neogene-elasmobranchii

Copyright: August 2016 Palaeontological Association

Albertov 6, 128 43 - Prague 2, Czech Republic. degracia@gmail.com

Gabriel Aguirre-Fernández. Paleontological Institute and Museum, University of Zurich, Karl-Schmid-Strasse 4, Zürich, 8006, Switzerland. gabriel.aguirre@pim.uzh.ch

René Kindlimann. Paleontological Institute and Museum, University of Zurich, Karl-Schmid-Strasse 4, Zürich, 8006, Switzerland. mad_design@gmx.ch

Marcelo R. Sánchez-Villagra. Paleontological Institute and Museum, University of Zurich, Karl-Schmid-Strasse 4, Zürich, 8006, Switzerland. m.sanchez@pim.uzh.ch

Keywords: Early Miocene; Burdigalian; Cantaure Formation; sharks; rays; Megalodon; paleoenvironment

Submission: 22 March 2016 **Acceptance:** 29 June 2016

INTRODUCTION

Sharks and rays are essential and diverse elements of tropical marine ecosystems (Cortés et al., 2008; Klimley, 2013), and their fossils are ubiquitous in many localities (e.g., Cappetta, 2012). In “Tropical America” – the geographic area of the Western Hemisphere located between the Tropic of Cancer and the Tropic of Capricorn, Neogene chondrichthyan assemblages are known from Barbados (Casier, 1958, 1966), Brazil (Santos and Travassos, 1960; Santos and Salgado, 1971; Reis, 2005; Costa et al., 2009), Cuba (Iturralde-Vinent et al., 1996; MacPhee et al., 2003), Southern Mexico (González-Rodríguez et al., 2013), Panama (Pimiento et al., 2013a), Peru (Alván et al., 2006), Trinidad (Leriche, 1938), The Grenadines (Portell et al., 2008) and Venezuela (Leriche, 1938; Sánchez-Villagra et al., 2000; Aguilera, 2010; Aguilera and Rodrigues de Aguilera, 2004; Aguilera and Lundberg, 2010). Despite all previous efforts, the chondrichthyan fossil record from the Caribbean is still fragmentary and poorly known, making it difficult to reconstruct paleofaunas that could help our understanding of the major ecological and biogeographical changes that happened in the area during the Neogene. Among those changes are those associated with the uplift of the Central American land bridge (Coates and Stallard, 2013; Montes et al., 2015), which interrupted the inter-oceanic corridor known as the Central American Seaway (CAS), and influenced the water circulation patterns of Pacific and the Caribbean (O’Dea et al., 2007; Leigh et al., 2014). The geographic position of Venezuelan deposits renders them as relevant for investigating the effects of those major paleogeographic rearrangements.

More than 20 years of paleontological expeditions in the Paraguaná Peninsula, North Western Venezuela, produced a collection of 1836 fossil elasmobranch specimens from the Cantaure Formation. This geological unit is thought to have

been deposited under fully marine conditions (Díaz de Gamero, 1974; Rey, 1996), exhibiting a well-known high diversity of invertebrates (Ingram, 1947; Jung, 1965; Gibson-Smith, 1974; Gibson-Smith and Gibson-Smith, 1974; Aguilera et al., 2010; Landau et al., in press) and teleostean fishes (Nolf and Aguilera, 1998; Aguilera, 2010; Aguilera and Lundberg, 2010). We hereby present a new taxonomic study of the elasmobranch fauna from the Cantaure Formation and a discussion of the paleoenvironmental and feeding preferences of these elasmobranchs, based on comparisons with extant species.

GEOLOGICAL AND STRATIGRAPHIC SETTING

The Cantaure Formation (Jung, 1965) has been dated as early Miocene, based on biozones of planktonic foraminifera (N7-8) and calcareous nannofossils (NN4-5) (Díaz de Gamero, 1974; Rey, 1996). According to Gradstein et al. (2012), biozones N7 and N8 correspond to the Lower to Middle Miocene transition (upper Burdigalian to lower Langhian). Griffiths et al. (2013) assigned a Burdigalian age (16.5 ± 0.4 Ma.) to a horizon of the Cantaure Formation using Sr isotopes from corals (Figure 1.2). Landau et al. (in press) suggested that the gastropod assemblage from the Cantaure Formation is typical for early-middle Miocene deposits. The stratotype of the Cantaure Formation is located approximately 10 km west of Pueblo Nuevo on the Paraguaná Peninsula, Falcón State, Venezuela. Outcrops are found south of Casa Cantaure (Figure 1.1) and are composed of fossiliferous silty shales interbedded with thin algal limestones and shell beds (Hunter and Bartok, 1974). Jung (1965) estimated a thickness of 75 m for the composite section of the Cantaure Formation. Hunter and Bartok (1974) divided the Cantaure Formation into three sections: 1) fossiliferous basal breccia unconformably overlaying the granite basement, 2) lower section comprised mainly of




FIGURE 1. Location and stratigraphy of the Cantaure Formation. 1. Fossiliferous localities. 2. Stratigraphic section of locality 1 (artesian well) and locality 2 (outcrop). Stratigraphic column modified after Aguilera (2010) and Aguilera et al. (2013).

shales and some sandy horizons [including the mollusk rich horizons illustrated by Jung (1965)], and 3) upper section represented by silty shales interbedded with thin algal limestones and shell beds. However, Hunter and Bartok (1974) did not provide a stratigraphic column, while they mentioned that the landscape of the area prevents a detailed measurement of the complete section. In contrast, Rey (1996) constructed a stratigraphic column of the suggested basal and middle sections, and also assumed that the mollusks described by Jung (1965), Gibson-Smith (1974) and Gibson-Smith and Gibson-Smith (1974), were derived from the upper section.

The top of the Cantaure Formation is conformably overlain by limestone layers that have been interpreted by Jung (1965) and Hunter and Bartok (1974) as post-Cantaure and pre-Pliocene sedimentation. Rey (1996) suggested that the upper contact of the Cantaure Formation represents an unconformity with the Amuay Member

(early Pliocene) of the Paraguaná Formation. Our field observations (OAA, JDCB, pers. obs.) indicate possible lateral variations in a short distance, with fossiliferous outcrops (e.g., thick layers of barnacles) that are not exposed in the previously known and described fossiliferous localities (e.g., Jung, 1965; Hunter and Bartok, 1974; Rey, 1996, among others). All the above clearly indicates that the Cantaure Formation is a more complex sequence than previously thought, in need of a new, detailed stratigraphic redescription.

MATERIAL AND METHODS

The fossil elasmobranch fauna described here (Table 1; Appendix 1) consists of 1836 specimens from four localities in the Cantaure Formation (Figure 1.1), which have been collected by the authors and other collaborators during several expeditions since 1992. Locality 1 (11° 56' 20" N, 70° 1' 2" W) corresponds to an artesian well ~44 m deep (Figure 1.2) and locality 2 (a': 11° 56' 23" N,

TABLE 1. Elasmobranchii paleodiversity of the Cantaure Formation.

Superorder	Order	Family	Genus	Taxon
Galeomorphii	Heterodontiformes	Heterodontidae	<i>Heterodontus</i>	<i>Heterodontus</i> sp.
	Orectolobiformes	Ginglymostomatidae	<i>Nebrius</i>	<i>Nebrius</i> sp.
	Lamniformes	†Otodontidae	† <i>Paratodus</i>	† <i>Paratodus benedenii</i> (Le Hon, 1871)
			† <i>Carcharocles</i>	† <i>Carcharocles megalodon</i> (Agassiz, 1843)
	Carcharhiniformes	Alopiidae	<i>Alopias</i>	<i>Alopias</i> cf. <i>A. vulpinus</i> (Bonnaterre, 1788)
			<i>Galeorhinus</i>	<i>Galeorhinus</i> cf. <i>G. galeus</i> (Linnaeus, 1758)
		Hemigaleidae	<i>Mustelus</i>	<i>Mustelus</i> sp.
			<i>Hemipristis</i>	† <i>Hemipristis serra</i> (Agassiz, 1835)
		Carcharhinidae	† <i>Galeocerdo</i>	† <i>Galeocerdo aduncus</i> Agassiz, 1843
				† <i>Galeocerdo mayumbensis</i> Darteville and Casier, 1943
			<i>Rhizoprionodon</i>	<i>Rhizoprionodon</i> sp.
			<i>Carcharhinus</i>	† <i>Carcharhinus ackermannii</i> Santos & Travassos, 1960
				<i>Carcharhinus brachyurus</i> (Günther, 1870)
				<i>Carcharhinus falciformis</i> (Müller and Henle, 1839)
				<i>Carcharhinus</i> cf. <i>C. galapagensis</i> (Snodgrass and Heller, 1905)
				† <i>Carcharhinus gibbesii</i> (Woodward, 1889)
				<i>Carcharhinus</i> cf. <i>C. limbatus</i> (Müller and Henle, 1839)
				<i>Carcharhinus</i> cf. <i>C. macroti</i> (Müller and Henle, 1839)

70° 1' 3" W and b': 11° 56' 2" N, 70° 0' 55" W) is an outcrop (overlying the section of locality 1) with a ~300 m long collecting surface (Figures 1.1-1.2). The corals used by Griffiths et al. (2013) for Sr isotope analysis were collected at locality 2 (Figure 1.1-1.2). This suggests that localities 1 and at least the lower section of locality 2 are of Burdigalian age (Figure 1.2). Localities 3 (11° 55' 59" N, 70° 1' 1" W) and 4 (11° 55' 48" N, 70° 00' 40" W) correspond to surface collecting areas along the south of Barbasco Creek, on the north escarpment of the Cerro San Carlos (Figure 1.1). Without a detailed stratigraphical knowledge of the outcrops south of Barbasco Creek, it is not possible to establish if localities 3 and 4 are different vertical strata (facies) along the Cantaure section or horizontal variations of localities 1 and 2. Field observations (OAA, JDCB and GAF pers. obs.) suggest that localities 3 and 4 are stratigraphically older than

the limestones layers attributed to post-Cantaure sedimentation (Jung, 1965; Hunter and Bartok, 1974).

Large specimens were surface-collected directly from the outcrop, while microsamples were collected after screen-washing approximately 1000 and 380 kg of sediments (mesh sizes: 0.5 and 2 mm) from locality 1 and 2, respectively. Approximately 50 kg of sediment was processed from locality 3, but it did not contain otoliths or elasmobranch micro-teeth. Specimens from locality 4 were only surface collected. The Cantaure specimens (Appendix 2) are housed in the paleontological collections of the Alcaldía Bolivariana de Urumaco (AMU-CURS), Centro de Investigaciones Antropológicas, Arqueológicas y Paleontológicas of the Universidad Experimental Francisco de Miranda (CIAAP, UNEFM-PF), and the Museo de Ciencias de Caracas (MCNC), all in Venezuela.

TABLE 1 (continued).

Superorder	Order	Family	Genus	Taxon
Batomorphii	Rajiformes	Sphyrnidae		<i>Carcharhinus perezii</i> (Poey, 1876)
				<i>Carcharhinus</i> sp. 1
				<i>Carcharhinus</i> spp.
			<i>Isogomphodon</i>	† <i>Isogomphodon acuaris</i> (Probst, 1879)
			<i>Negaprion</i>	† <i>Negaprion eurybathrodon</i> (Blake, 1862)
			† <i>Physogaleus</i>	† <i>Physogaleus contortus</i> (Gibbes, 1849)
			Gen. indet.	sp. indet.
			<i>Sphyrna</i>	† <i>Sphyrna arambourgi</i> Cappelletta, 1970
				<i>Sphyrna</i> cf. † <i>S. laevis</i> (Cope, 1867)
			<i>Rhynchobatus</i>	<i>Rhynchobatus</i> sp.
	Myliobatiformes	Pristidae	<i>Pristis</i>	<i>Pristis</i> sp.
			<i>Dasyatis</i>	<i>Dasyatis</i> sp.
		Dasyatidae	cf. <i>Pteroplatytrygon</i>	cf. <i>Pteroplatytrygon</i> sp.
			cf. <i>Taenirops</i>	cf. <i>Taenirops</i> sp.
				Dasyatidae Indet.
		Myliobatidae	<i>Aetobatus</i>	<i>Aetobatus</i> sp.
			<i>Aetomylaeus</i>	<i>Aetomylaeus</i> sp.
		Rhinopteridae	<i>Rhinoptera</i>	<i>Rhinoptera</i> sp.
		Mobulidae	<i>Mobula</i>	† <i>Mobula fragilis</i> (Cappelletta, 1970)
				<i>Mobula</i> cf. † <i>M. loupianensis</i> (Cappelletta, 1970)
				<i>Mobula</i> sp.
			† <i>Plinthiscus</i>	† <i>Plinthiscus stenodon</i> Cope, 1869
				Myliobatiformes indet.
				Chondrichthyes indet.

The taxonomy follows Cappelletta (2012) and Compagno (2005), with the exception of the extinct genus *Carcharocles* Jordan and Hannibal, 1923, for which we follow the nomenclature of Pimiento et al. (2010). Taxonomic identification was based on an extensive literature review, and also comparative analyses between fossil and recent specimens from the following collections: Mapuka Museum of Universidad del Norte (MUN-STRI), Barranquilla, Colombia; Museu Paraense Emilio Goeldi (MPEG-V), Belem, Brazil; Natural History Museum of Basel (NMB), Switzerland; Paleontological collection of the Institut des Sciences de l'Evolution, University of Montpellier (UM), France; Palaeontological Institute and Museum at the University of Zurich (PIMUZ), Switzerland and René Kindlimann private collection, Uster, Switzerland.

We analyzed the abundance using percentages of specimens by order, families and species

using the 39 securely recognized taxa that are represented by 1711 specimens (out of the 1836 collected in total) (Appendix 1). For conducting the paleoecological interpretation, we compiled information on habitat preference (benthic, benthopelagic, pelagic, neritic and bathyal) and feeding ecology (dietary composition and behavior) of extant taxa recorded in the Cantaure Formation (Appendix 3), following Compagno (1984a, 1984b); Cortés (1999); Compagno et al. (2005); Musick et al. (2004); Kiraly et al. (2003); Voigt and Weber (2011); Cortés et al. (2008); Ebert and Stehmann (2013); and the FishBase website (Froese and Pauly, 2015). The paleobathymetric analysis was performed following the methodology of Nolf and Brzobohatý (1994), adapted to fossil sharks (Carrillo-Briceño et al., 2015a, 2016). For this analysis we included only species/genera with closely related extant taxa. Extinct species and taxa with-

out clear identification to genus were removed from the analysis. A total of 34 (out of 39) taxa, for which ecological information is available, were analyzed.

Taking into consideration the limitations associated with paleoecological interpretations of fossil assemblages, we tried to approach the ecological role that members of the Cantaure elasmobranch fauna played in this ancient marine environment. We use the categorization of elasmobranch feeding preferences suggested by Cortés (1999) and Cortés et al. (2008). Living sharks and rays as a whole exhibit a wide range of diets, but each species has specific preferences. The latter are used to infer the diet of taxa we documented from the Cantaure assemblage (Appendix 3).

RESULTS

Taxonomical Composition

The taxonomical composition of the four study localities includes at least 39 species of galeomorphs and batoids (Table 1; Appendix 1, 4). Galeomorphs are represented by 26 species, 15 genera and eight families of Heterodontiformes, Orectolobiformes, Lamniformes and Carcharhiniformes (Table 1; Figure 2). Batoids include 13 species attributed to 10 genera and six families of Rajiformes and Myliobatiformes (Table 1; Figure 2). Pictures of the chondrichthyan teeth, vertebrae and caudal and rostral denticles in Figures 3-12, were illustrated following the taxonomic classification shown in Table 1.

Heterodontiformes Berg, 1937. The record of this group (Heterodontidae Gray, 1851) consists of a single and undetermined specimen of *Heterodontus* Blainville, 1816. The tooth was collected from locality 2 (Appendix 1, 4), and was previously reported by Aguilera and Lundberg (2010) and Aguilera (2010). The specimen is missing from the UNEFM collection and it is not illustrated here.

Orectolobiformes Applegate, 1972. The orectolobiform specimens are characterized by a long apron and a short main cusp flanked by more than seven pairs of lateral cusplets (Figure 3.1-3.9). This condition is diagnostic for *Nebrius* Rüppell, 1837, a genus recorded from North America going back to the Eocene (e.g., Kent, 1994; Müller, 1999). The specimens from the Cantaure assemblage (Figure 3.1-3.9; Appendix 1, 4) were previously assigned to *Ginglymostoma delfortriei* (Daimeries, 1889) (Aguilera and Rodrigues de Aguilera, 2004; Aguilera, 2010; Aguilera and Lundberg, 2010). Fossils of *Nebrius* in the Americas have not been thoroughly studied, and there is no

clear consensus about their taxonomy at the species level or stratigraphic range (e.g., Kent, 1994; Cicimurri and Knight, 2009). Therefore we refrain from assigning species to our specimens. Cicimurri and Knight (2009) suggested that some fossil teeth of *Nebrius* (from North America) have been misidentified as *Ginglymostoma* Müller and Henle, 1837. The specimens referred to *G. delfortriei* from Panama (Pimiento et al., 2013b) and *Ginglymostoma* sp. from Brazil (Costa et al., 2009) resemble our *Nebrius* teeth from the Cantaure Formation.

Lamniformes Berg, 1937. These sharks are represented by one species of Alopiidae Bonaparte, 1838 and two genera and two species of †Otodontidae Glikman 1964 (Figure 2). *Alopias* cf. *A. vulpinus* (Bonnaterre, 1788) is represented by three specimens (Figure 3.22-3.26; Appendix 1, 3) while the otodontid †*Paratodus benedenii* (Le Hon, 1871) is the least abundant lamniform (Figure 3.10-3.11; Appendix 1, 4). The most abundant lamniform is the otodontid †*Carcharocles megalodon* (Agassiz, 1843), with more than 100 specimens (Figure 3.12-3.21) from all studied localities (Appendix 1, 4). Many *C. megalodon* teeth are broken; however, most of them, especially large specimens, preserve the wide triangular shape (crown without lateral cusplets), a v-shaped 'neck', a fine serrations, and the large size that characterize this species (Purdy et al., 2001; Cappetta, 2012; Pimiento et al., 2010, 2013b). Small teeth having a pair of short lateral cusplets may belong to juvenile and/or subadult individuals of *C. megalodon* (Figure 3.12-3.16), a feature not present in adult teeth (Figure 3.17-3.21). Morphologically, the younger species *Carcharocles chubutensis* (Ameghino, 1901), differs from *C. megalodon* by the presence of lateral cusplets not separated from the crown on the teeth of juveniles, subadults and adults (Cappetta, 2012; Pimiento et al., 2013a). In more derived populations as *C. megalodon*, lateral cusplets disappear, except sometimes on very lateral files or in juvenile individuals (Cappetta, 2012). This absence of lateral cusplets in *C. megalodon* adult teeth has been interpreted as the result of heterochronic processes in the *Carcharocles* clade through geologic time (Pimiento et al., 2010, 2013a, 2013b; Pimiento and Balk, 2015). The absence of lateral cusplets in all well-preserved teeth interpreted as adults (including also some teeth that may belong to juvenile/subadults specimens) allows us to assign our specimens to *C. megalodon*. In addition, adult teeth of *C. megalodon* referred here, are very similar to those of the




FIGURE 2. Elasmobranch paleodiversity of the Cantaure Formation. 1. Orders and 2. Families and genera.

same species, found in the late Burdigalian section (~16.2 Ma.) of the Castilletes Formation at North East Colombia (under study by JDCB).

Carcharhiniformes Compagno, 1973. This order is represented by four families, 10 genera and 21 species, thus being the most abundant and diverse one of the Cantaure assemblage (Figures 2, 4-8; Table 1; Appendix 1, 4). Triakidae Gray, 1851 [two genera and two species: *Galeorhinus* cf. *G. galeus* (Linnaeus, 1758) (Figure 4.1-4.2), and *Mustelus* Linck, 1790 (*Mustelus* sp.) (Figure 4.3-4.7)], Hemigaleidae Hasse, 1879 [*Hemipristis serra* (Agassiz, 1835) (Figure 4.8-4.13)], and Sphyrnidae Gill,

1872 [*Sphyrma arambourgi* Cappetta, 1970 (Figure 7.21-7.25), and *Sphyrma* cf. *S. laevis* (Cope, 1867) (Figure 8.1-8.5)], are the less diverse families of this order in the Cantaure assemblage (Figure 2; Appendix 1, 4). Nevertheless, *Hemipristis serra* with 425 specimens (Appendix 1, 4), is the second most abundant species of the assemblage, representing 23% of the total specimens collected. The specimen referred by Aguilera and Rodrigues de Aguilera (2004) to *Paragaleus* sp. corresponds to a juvenile posterior ?lower tooth of *H. serra* (Figure 4.8).




FIGURE 3. Orectolobiformes and Lamniformes of the Cantaure Formation. 1-9. *Nebrius* sp. (AMU-CURS-986); 10-11. †*Paratodus benedenii* (AMU-CURS-645). 12-21. †*Carcharocles megalodon* (12: AMU-CURS-997; 13: AMU-CURS-512; 14: UNEFM-PF-349; 15: AMU-CURS-996; 16: AMU-CURS-518; 17-18: UNEFM-PF-s/n; 19: AMU-CURS-512 and 20-21: AMU-CURS-515). 22-26. *Alopias* cf. *A. vulpinus* (22-23: AMU-CURS-985; 24-25: AMU-CURS-983 and 26: AMU-CURS-984). Jaw position: upper (10-11? 12, 14? 17, 22-25), lower (13, 15, 18) and indet. (1-9, 16, 19-21, 26). View: labial (1, 6, 10, 13, 20-21, 23, 24), lingual (2, 7, 11-12, 14-19, 22, 25-26), profile (3, 8), occlusal (4) and basal (5, 9).




FIGURE 4. Carcharhiniformes of the Cantaure Formation. 1-2. *Galeorhinus* cf. *G. galeus* (AMU-CURS-974). 3-7. *Mustelus* sp. (AMU-CURS-975). 8-13. †*Hemipristis serra* (8: AMU-CURS-731 and 9-13: AMU-CURS-644). 14-18. †*Galeocерdo aduncus* (14-15: AMU-CURS-730 and 16-18: AMU-CURS-647). 19-23. †*Galeocерdo mayumbensis* (19-20: AMU-CURS-995 and 21-23: AMU-CURS-646). Jaw position: upper (9-12), lower (8? 13) and indet. (1-7, 14-23). View: labial (1, 6, 7, 10, 15, 19, 21-22), lingual (2, 4, 8-9, 11-14, 16-18, 20, 23) and occlusal-lingual (3, 5).

The Carcharhinidae Jordan and Evermann, 1896, with six genera and 16 species, is the most diverse and abundant family of the assemblage (Figure 2; Table 1; Appendix 1, 4). The genera *Rhizoprionodon* Whitley, 1929 [*Rhizoprionodon* sp. (Figure 5.1-5.5)], *Isogomphodon* Gill, 1862 [†*Isogomphodon acuarius* (Probst, 1879) (Figure 7.1-7.5)], *Negaprion* Whitley, 1940 [†*Negaprion eurybathrodon* (Blake, 1862) (Figure 7.6-7.10)], and †*Physoga-*

leus Cappetta, 1980 [†*Physogaleus contortus* (Gibbes, 1849) (Figure 7.11-7.15)], are represented by one species each. *Galeocерdo* Müller and Henle, 1837 is represented by two species [†*Galeocерdo aduncus* Agassiz, 1843 (Figure 4.14-4.18), and †*Galeocерdo mayumbensis* Darteville and Casier, 1943 (Figure 4.19-4.23)]. *Negaprion eurybathrodon* with 586 specimens (Appendix 1, 4) is the most abundant species of the assemblage,




FIGURE 5. Carcharhiniformes of the Cantaure Formation. 1-5. *Rhizoprionodon* sp. (1-3, 5: AMU-CURS-975 and 4: AMU-CURS-960). 6-10. †*Carcharhinus ackermannii* (6-7: AMU-CURS-713 and 8-10: AMU-CURS-714). 11-15. *Carcharhinus brachyurus* (AMU-CURS-990). 16-20. *Carcharhinus falciformis* (AMU-CURS-991). 21-25. *Carcharhinus* cf. *C. galapagensis* (AMU-CURS-973). Jaw position: upper (2, 4-14, 16-25) and lower (1, 3, 15). View: labial (6, 9, 12-13, 17-18, 20-21, 24) and lingual (1-5, 7-8, 10-11, 14-16, 19, 22-23, 25).

representing 32% of the total specimens collected. Specimens referred to *Mustelus* sp. (Triakidae) and *Rhizoprionodon* sp. (Carcharhinidae), are few and fragmentary, lacking diagnostic characters for accurate species assignment.

The genus *Carcharhinus* Blainville, 1816, with nine species, represents the most diverse genus

from the Cantaure assemblage (Figures 2, 5.6-5.25, 6.1-6.24; Table 1; Appendix 1, 4). With the exception of the extinct †*Carcharhinus ackermannii* Santos and Travassos, 1960 (Figure 5.6-5.10) and †*Carcharhinus gibbesii* (Woodward, 1889) (Figure 6.1-6.4), the species *Carcharhinus brachyurus* (Günther, 1870) (Figure 5.11-5.15),




FIGURE 6. Carcharhiniformes of the Cantaure Formation. 1-4. †*Carcharhinus gibbesii* (AMU-CURS-958). 5-8. *Carcharhinus* cf. *C. limbatus* (AMU-CURS-944). 9-12. *Carcharhinus* cf. *C. macroti* (AMU-CURS-957). 13-17. *Carcharhinus perezii* (AMU-CURS-956). 18-22. *Carcharhinus* sp.1 (AMU-CURS-943). 23-24. *Carcharhinus* spp. (23: AMU-CURS-941 and 24: AMU-CURS-940). Jaw position: upper (1-24). View: labial (1, 4, 6-7, 10, 12, 14, 17-18, 20, 22-24) and lingual (2-3, 5, 8-9, 11, 13, 15-16, 19, 21).

Carcharhinus falciformis (Müller and Henle, 1839) (Figure 5.16-5.20), *Carcharhinus* cf. *C. galapagensis* (Snodgrass and Heller, 1905) (Figure 5.21-5.25), *Carcharhinus* cf. *C. limbatus* (Müller and Henle, 1839) (Figure 6.5-6.8) and *Carcharhinus perezii* (Poey, 1876) (Figure 6.13-6.17), have living counterparts and most of them inhabit the Atlantic and Pacific coast of America (Compagno et al., 2005). *Carcharhinus macroti* (Müller and Henle, 1839) (Figure 6.9-6.12) is today an exclusive

inhabitant of the Indo-west Pacific Ocean (Compagno et al., 2005). The specimens referred to *Carcharhinus* sp.1 (Figure 6.18-6.22), have a dental morphology different, especially in the form of the crown, which differentiate these teeth from the rest of the *Carcharhinus* species found in the Cantaure assemblage. Due to the scarcity of specimens (Appendix 1) and the lack of preserved diagnostic characters in *Carcharhinus* teeth we have used for comparison (fossil and recent speci-




FIGURE 7. Carcharhiniformes of the Cantaure Formation. 1-5. †*Isogomphodon acuarius* (1-2: AMU-CURS-950 and 3-5: AMU-CURS-951). 6-10. †*Negaprion eurybathrodon* (6-9: AMU-CURS-989 and 10: AMU-CURS-987). 11-15. †*Physogaleus contortus* (11-12: AMU-CURS-719 and 13-15: AMU-CURS-648). 16-20. Carcharhinidae indet. (AMU-CURS-939). 21-25. †*Sphyrna arambourgi* (21-24: AMU-CURS-961 and 25: AMU-CURS-962). Jaw position: upper (1-3, 6-9, 16-19? 21-22, 25), lower (4-5, 10, 20? 23-24) and indet. (11-15). View: labial (2, 6, 9, 11, 13, 17-18, 22, 24-25) and lingual (1, 3-5, 7-8, 10, 12, 14-16, 19-21, 23).

mens), we prefer to keep these specimens to a generic level. We have referred all specimens that are broken and eroded, without any diagnostic features for specific identification (Figure 6.23-6.24), to *Carcharhinus* spp. Some of the specimens clas-

sified as Carcharhinidae indet. (Figure 7.16-7.20; Table 1; Appendix 1, 4), exhibit a very prominent lingual protuberance with a sigmoid cusp, which could be the basis of a more detailed taxonomical identification in future studies.




FIGURE 8. Carcharhiniformes and Rajiformes of the Cantaure Formation. 1-5. *Sphyrna* cf. †*S. laevisissima* (1-3, 5: AMU-CURS-964 and 4: AMU-CURS-963). 6-11. *Rhynchobatus* sp. (6, 9: AMU-CURS-968; 7-8, 10: AMU-CURS-969 and 11: AMU-CURS-970). 12-20. *Pristis* sp. (rostral denticle 12-14: AMU-CURS-242; buccal teeth 15-20: AMU-CURS-967). Jaw position: upper (1-2) and indet. (3-11, 15-20). View: labial (1, 3), lingual (2, 4-5, 11), occlusal (6-8, 15, 17, 19), profile (18, 20), basal (9-10, 16), posterior (12) and dorsal (13-14).

Rajiformes Berg, 1937. This group of batoids is represented by two families, two genera and two species (Figure 2). Concerning Rhynchobatidae Garman, 1913, we report indeterminate teeth of *Rhynchobatus* Müller and Henle 1837 (Figure 8.6-8.11; Appendix 1, 4), resembling previously figured isolated teeth of *Rhynchobatus* from the late Miocene of Costa Rica, Panama and Venezuela (Laurito, 1999; Pimiento et al., 2013b; Carrillo-Briceño et al., 2015b). We refrain from taxonomic identification at the species level of the Cantaure *Rhynchobatus*, because the range of dental variation in extant species is unknown, and little is known about fossil species from the Americas. The *Pristi-*

dae Bonaparte, 1838 is represented by a few specimens of oral teeth and rostral denticles of *Pristis* Linck, 1790 (Figure 8.12-8.20, Appendix 1, 4). As noted by Carrillo-Briceño et al. (2015b), *Pristis* rostral denticles are not diagnostic at species level, while isolated oral teeth are somewhat difficult to identify, because those of extant species are poorly known and are morphologically diverse.

Myliobatiformes Compagno, 1973. This group is represented by four families, eight genera and 11 species, being thus the second most abundant and diverse order from the Cantaure assemblage (Figure 2; Table 1; Appendix 1, 4). The *Dasyatidae* Jordan, 1888, is represented by three genera and four




FIGURE 9. Myliobatiformes of the Cantaure Formation. 1-10. *Dasyatis* sp. (1-2, 7-8: AMU-CURS-930; 3-4: AMU-CURS-932; 5-6: AMU-CURS-933 and 9-10: AMU-CURS-931). 11-17. cf. *Pteroplatytrigon* sp. (AMU-CURS-926). 18-21. cf. *Taeniurops* sp. (AMU-CURS-925). 22-27. *Dasyatidae* indet. (22-24: AMU-CURS-936 and 25-27: AMU-CURS-937). Jaw position: indet. (1-27). View: labial (4, 9, 11, 15, 21), lingual (2, 16), occlusal (1, 5, 13, 18, 22, 25), profile (3, 6-7, 10, 12, 17, 19, 24, 27) and basal (8, 14, 20, 23, 26).




FIGURE 10. Myliobatiformes of the Cantaure Formation. 1-3. *Aetobatus* sp. (1-3: AMU-CURS-716). 4-14. *Aetomylaeus* sp. (AMU-CURS-750). 15-29. *Rhinoptera* sp. (15-18: AMU-CURS-979; 19-21, 22-23, 24-25, 28-29: AMU-CURS-977 and 26-27: AMU-CURS-981). Jaw position: indet. (1-29). View: labial (11, 18), lingual (5, 12, 17, 21-22, 24, 28), occlusal (1, 4, 8, 10, 15, 19, 23, 26), profile (3, 13-14) and basal (2, 6, 7, 9, 16, 20, 25, 27, 29).

species, and the Myliobatidae Bonaparte, 1838 and Rhinopteridae Jordan and Evermann, 1896 by one genus and one species each (Figure 2; Table 1). Due to the small number of specimens and the lack of diagnostic characters, teeth referred to *Dasyatis* Rafinesque, 1810 (Figure 9.1-9.10), cf. *Pteroplatytrygon* Fowler, 1910 (Figure 9.11-9.17), cf. *Taeniurops* Garman, 1913 (Figure 9.18-9.21), and *Dasyatidae* indet. (Figure 9.22-9.27), cannot be assigned to species. As noted by Carrillo-Briceño et al. (2015b), there is very little known available about the morphological diversity of dental patterns in extant and fossil *Dasyatidae* from the Americas.

Teeth referred to *Aetobatus* Blainville, 1816 (Figure 10.1-10.3), and *Aetomylaeus* Garman, 1913 (Figure 10.4-10.14), are fragmented and incomplete; but preserve diagnostic characters that made their generic identification possible. Teeth of *Rhinoptera* Cuvier, 1829 (Figure 10.15-10.29) are common in all localities (especially at locality 2), being the third most abundant species of the assemblage (Appendix 1, 4). Previous work on the Cantaure Formation has reported the presence of *Myliobatis* Cuvier, 1816 teeth (Aguilera and Rodrigues de Aguilera, 2004; Aguilera and Lundberg, 2010; Aguilera, 2010). However, our revision of fossils led to the identification of isolated medial

and lateral teeth, which exhibit the typical imbrication pattern and morphology of *Rhinoptera*. *Mobulidae* Gill, 1893 is represented by two genera and four species (Figure 2; Table 1; Appendix1, 4). The mobulids †*Mobula fragilis* Cappetta, 1970 (Figure 11.1-11.11), *Mobula* cf. †*M. loupianensis* Cappetta, 1970 (Figure 11.12-11.18) and *Mobula* sp. (Figure 11.19-11.23) are scarce (Appendix1, 4), and their teeth have been recovered from localities 1 and 2 only. One tooth of †*Plinthicus stenodon* Cope, 1869 was found at locality 3.

All eroded, broken and non-diagnostic caudal spines (Figure 12.1-12.6) are referred to Myliobatiformes indet. In addition, specimens referred to Chondrichthyes indet. correspond to fragmented and non-diagnostic vertebrae (Figure 12.7-12.8).

Elasmobranch Paleobathymetric Analysis

The Cantaure elasmobranch fauna is characterized by a predominance of benthopelagic sharks and rays, the extant relatives of which inhabit diverse environments and have diverse bathymetric preferences, especially in neritic/epipelagic environments (Appendix 3). Our paleobathymetric analysis indicates that the most probable deposition depth for the Cantaure assemblage is the 0 to 50 m interval, followed by the 50 to 100 m interval as the second most probable (Figure 13). This sug-




FIGURE 11. Mobulids of the Cantaure Formation. 1-11. †*Mobula fragilis* (1-3, 10-11: AMU-CURS-947; 4-6: AMU-CURS-948 and 7-9: AMU-CURS-949). 12-18. *Mobula* cf. †*M. loupianensis* (AMU-CURS-946). 19-23. *Mobula* sp. (AMU-CURS-927). 24-27. †*Plinthisus stenodon* (AMU-CURS-715). Jaw position: indet. (1-27). View: labial (1, 4, 7, 10, 14-15, 17, 22, 24), lingual (3, 6, 20, 26), occlusal (2, 5, 8, 16, 19, 25), profile (11-12, 18, 23, 27) and basal (9, 13, 21).

gests that the Cantaure Formation was deposited in an inner-middle shelf environment. The presence of *Nebrius* sp., *Carcharhinus* cf. *C. limbatus*, *C. perezi*, *Isogomphodon acuaris*, *Negaprion eurybathrodon*, *Rhynchobatus* sp. and *Rhinoptera* sp. support this hypothesis, due to the fact that their extant representatives inhabit waters shallower than those of middle shelf environments (Appendix 3). Other taxa recognized in the first and second most probable depth intervals (Figure 13) include Alopidae, Carcharhinidae, Sphyrnidae, Rinchobatidae, Pristidae, Dasyatidae, Myliobatidae and Mobulidae, whose extant representatives are associated with coastal environments. However, it is important to keep in mind that some of those species also have a wide vertical distributional range (Appendix 3), or are even able to move

along significant distances over oceanic basins (Compagno, 1984b; Compagno et al., 2005; Voigt and Weber, 2011; Froese and Pauly, 2015).

Dietary Preferences of the Cantaure Elasmobranchs

The dietary composition of the elasmobranch assemblage from the Cantaure Formation can be represented by five food categories, from filter feeders to benthic-pelagic predators (Figure 14; Appendix 3). In this assemblage the most abundant feeder group is that of the piscivorous, representing 49% of the total species count (Figure 14; Appendix 3). We have included in this group the pelagic lamnid *Paratodus benedenii* (Cappetta, 1987, Kent and Powell, 1999) and the carcharhinid *Physogaleus contortus*, extinct species with a cut-




FIGURE 12. Myliobatiformes and other chondrichthyans of the Cantaure Formation. 1-6. Myliobatiformes indet. (AMU-CURS-507). 7-8. vertebra of chondrichthyan Indet. (AMU-CURS-928). 9-10. tooth of †*Galeocерdo aduncus* in association with barnacles of the family Balanidae (AMU-CURS-720). Jaw position: indet. (9). View: lingual (9), dorsal (1, 4, 5-6), ventral (2-3) and indet. (7-8).

ting-clutching dental type that could suggest a fish-based diet. The durophagous/cancritrophic (mollusk, crustacean, coral feeders) group is the second most abundant, corresponding to 30% of the species in the Cantaure assemblage, with a predominance of Myliobatiformes and Rajiformes (Figure 14; Appendix 3). In this group the most abundant taxon is *Rhinoptera* sp., living species of which are opportunistic/generalists consuming mainly mollusks. The filter feeder group (Figure 14) is represented in the Cantaure assemblage only by mobulids (Table 1; Appendix 1, 4), whose living counterparts are characterized by a diet based mainly on planktonic microorganisms (Klimley,

2013). One of its representatives is the extinct mobulid *Plinthiscus stenodon*, whose fragile teeth also suggest a possible filter diet of soft prey (Capetta, 2012). In the eurytrophic/sarcophagous group (diverse food source: fishes, reptiles, birds, mammals, etc.) (Figure 14), we have included large sharks with a broad habitat preference. Those species include *Carcharocles megalodon*, *Galeocерdo aduncus* and *G. mayumbensis*, which could have had the role of apex predators. In reference to the teuthitrophic group, it is represented only by *Alopias* cf. *A. vulpinus*, whose living counterpart has a higher preference for cephalopods (Cortés, 1999; Appendix 3).




FIGURE 13. Bathymetric ranges estimated for Cantaura Formation taxa and their relative abundance. Arrows indicate that the bathymetric range is greater; question mark symbol indicates uncertainty about the extent of the range. A dashed red line indicates that there is no information for the taxa. The gray shadow indicates the more probably depth range for this assemblage. In the relative abundance graphic the y-axis is shortened between 2 and 3 to improve the values visualization; note the greater abundance of shallow water taxa. The extinct species including *P. benedenii*, *C. megalodon*, *P. contortus*, and *P. stenodon*, and taxa without clear identification, including *Carcharhinus* spp., *Carcharhinidae* indet., *Myliobatiformes* indet. and *Chondrichthyes* indet., were removed from the analysis.




FIGURE 14. Dietary preference composition of the elasmobranchs from the Cantaure Formation. Percentages do not include *Carcharhinus* spp., Myliobatiformes indet. and Chondrichthyes indet.

DISCUSSION

Paleodiversity Significance

Forty-four percent of the Cantaure elasmobranch assemblage consists of genera or species with living representatives in Tropical America (Compagno, 1984a, 1984b; Compagno et al., 2005; Voigt and Weber, 2011). Five genera and 15 species present are extinct (Table 1). *Nebrius* sp., *Carcharhinus* cf. *C. macloiti*, and *Rhynchobatus* sp., are present in our fossil sample and are currently extinct in the Caribbean, but have living counterparts in the eastern Atlantic and Indo-West Pacific (Compagno et al., 2005; Froese and Pauly, 2015). Their fossil record confirms that these taxa became extinct in the Caribbean, Western Atlantic and Eastern Pacific, possibly as a consequence of ecological competition with other species or environmental changes during the final stage of the CAS (e.g., Coates and Stallard, 2013).

The galeomorphs *Paratodus benedenii*, *Galeocerdo mayumbensis*, *C. brachyurus*, *Carcharhinus* cf. *C. galapagensis*, *C. gibbesii*, *Carcharhinus* cf. *C. macloiti*, *Isogomphodon acuaris*, *Physogaleus contortus*, *Sphyrna arambourgi*, *Sphyrna* cf. *S. laevissima*, and the batoids cf. *Pteroplatytrygon* sp., cf. *Taeniurops* sp., *Mobula fragilis* and *Mobula* cf. *M. loupianensis* are reported for the first time from Neogene Venezuelan deposits. *Paratodus benedenii*, *Carcharhinus* cf. *C. gala-*

pagensis, *Carcharhinus* cf. *C. macloiti*, cf. *Pteroplatytrygon* sp., *Taeniurops* sp., *Mobula fragilis* and *Mobula* cf. *M. loupianensis* are new records for the Neogene of Tropical America. With the exception of the records mentioned above from Tropical America, the fossil record of most elasmobranch taxa from the Cantaure Formation have been found in other Neogene marine deposits of the Americas and are summarized in Kruckow and Thies (1990), Laurito (1999), Purdy et al. (2001), Aguilera et al. (2011), González-Rodríguez et al. (2013) and Carrillo-Briceño et al. (2014, 2015a, 2016).

Early Neogene marine elasmobranch assemblages known from the Americas are summarized in Figure 15 and include North America (Purdy et al., 2001; Visaggi and Godfrey, 2010), Southern South America (Suarez et al., 2006; Cione et al., 2011) and Tropical America (Leriche, 1938; Casier, 1966; MacPhee et al., 2003; Alván et al., 2006; Portell et al., 2008; Costa et al., 2009; Aguilera and Lundberg, 2010; Pimiento et al., 2013b, Laurito et al., 2014; Carrillo-Briceño et al., 2016). From these assemblages the Uitpa Formation (Colombia) has been referred to the Aquitanian (Carrillo-Briceño et al., 2016), while the Calvert and Pungo River formations in North America (Purdy et al., 2001; Visaggi and Godfrey, 2010) and the Cantaure (in this work) and Castillo (Rincón et al., 2014) formations in Venezuela, have been referred to the Burdigala-




FIGURE 15. Early Neogene chondrichthyan diversity in the Americas. Tropical America: Barbados (Ba), Brazil (Br), Colombia (Co), Costa Rica (Cs), Cuba (Cu), The Grenadines (Gr), Panamá (Pa), Perú (Pu), Trinidad (Tr) and Venezuela (Ve). North America: United States of America (US). Southern South America: Argentina (Ar) and Chile (Ch). Aquitanian (Anq), Burdigalian (Brd), Langhian (Lngh); early Miocene without differentiation (EM), early-Middle Miocene (E-MM). Formation(Fm.), early (E). Note: to references see "Paleodiversity significance" (see Discussion).

lian. The remaining formations (Figure 15) have been assigned to an unresolved early Miocene age. The Pungo River (~38 spp.) and Cantaure (39 spp.) assemblages preserve the most diverse elasmobranch faunas known from early Miocene of the Americas.

The late Burdigalian age (16.5 ± 0.4 Ma.) of the Cantaure Formation based on Sr isotopes from corals from locality 2 of Griffiths et al. (2013) (Figure 1.2), suggests that all specimens found in localities 1 and 2 could be coetaneous or older than 16.5 ± 0.4 Ma. The poorly known stratigraphic position of localities 3 and 4, in reference to localities 1 and 2 (different vertical strata along the Cantaure section or horizontal variation of localities 1 and 2), does not permit an accurate age estimation for them, but their stratigraphic position (underlying the limestones interpreted as post-Cantaure sedimentation; Jung, 1965; Hunter and Bartok, 1974),

suggests that these localities (3 and 4) are close to the top section of the Cantaure Formation. A Burdigalian age (Griffiths et al., 2013), or even that of a Burdigalian-Langhian boundary for the upper section of the Cantaure Formation (assuming that the biozones N7-8 and NN4-5 represent the lower to middle Miocene transition), would extend the stratigraphic range of two galeomorphs and of a batoid found in this assemblage. Fossils of *Carcharhinus* cf. *C. galapagensis*, from the Cantaure Formation, represent the oldest record for this species. It was previously known from the late Miocene of Mexico (González-Rodríguez et al., 2013). The same applies to *Carcharhinus gibbesii*, the youngest record was reported by Carrillo-Briceño et al. (2016) as the Aquitanian of Colombia. In reference to the batoid cf. *Pteroplatytrygon* sp., its record from the Cantaure Formation could be the oldest known to this stingray species.

The presence of *Carcharocles megalodon* in Burdigalian sediments of the Cantaure Formation (in at least localities 1 and 2) confirms the presence of this species during late early Miocene. It has been generally accepted that the record of *Carcharocles chubutensis* spans from the early to the middle Miocene, whereas *Carcharocles megalodon* spans from the middle Miocene to the late Pliocene (Pimiento and Clements, 2014; Pimiento and Balk, 2015). However, other authors have suggested that *C. megalodon* also occurs in the Burdigalian of Europe and North America (Leriche, 1938; Purdy et al., 2001; Visaggi and Godfrey, 2010; Reinecke et al., 2011). According to Leriche (1938), the two *C. megalodon* specimens he collected (Leriche, 1938, Planche III, Figures 5-6) in southern Mexico (Tabasco and Isthmus of Tehuantepec) came from early Miocene strata and are similar to teeth he previously collected in the Burdigalian of Europe. Another early Miocene unpublished occurrence of *C. megalodon* from Tropical America is from the late Burdigalian section (~16.2 Ma.) of the Castilletes Formation at North East Colombia (under study by JDCB). The specimens of *C. megalodon* from the Cantaure sequence have been found in all four localities studied here (Figure 1, Appendix 1), but with a greater abundance in localities 2 and 3 (Appendix 1). The presence of *C. megalodon* tooth (UNEFM-PF-322) at least 6-10 m deep in the artisan well (Figure 1.2) could suggest an older age than the 16.5 ± 0.4 Ma for the species. More detailed information about the American and worldwide Neogene stratigraphic range of the remaining elasmobranch taxa recovered from the Cantaure Formation (Table 1) can be found in Kruckow and Thies (1990), Cicimurri and Knight (2009), Reinecke et al. (2011), Bor et al. (2012), Adnet et al. (2012), Cappetta, (2012), Pimiento et al. (2013b), and Carrillo-Briceño et al. (2014, 2015a, 2016).

Furthermore, it is important to note that the elasmobranch fauna from the Cantaure Formation shows a clear differentiation in paleodiversity amongst the four localities studied here (Appendix 1, 4). As is shown in Appendix 4.2, localities 1 and 2 (Figure 1) have a higher diversity than those of localities 3 and 4, a fact that could be attributable to: 1) less intensive sieving and sampling of the localities 3 and 4 (see Material and Methods section) or 2) different lithologic, taphonomic and preservational conditions. Future, more accurate dating of the deposits could offer new insights into the age of the Cantaure deposits, and especially of the sections located south of Barbasco Creek.

Paleoenvironment and Paleoecology

On the basis of planktonic foraminifera, calcareous nannofossils and mollusks, it has been proposed that the Cantaure Formation was deposited in a shallow, coastal tropical marine environment, with clear waters and normal salinity (Jung, 1965; Díaz de Gamero, 1974; Rey, 1996; Landau et al., in press). A conspicuous absence of typical herbivore gastropods suggests that sea-grass habitats were probably not represented in this environment (Landau et al., in press). The teleostean assemblage reported by Nolf and Aguilera (1998) reflects a shallow environment with a water depth of less than 50 m. The crustacean assemblage (Aguilera et al., 2010) is also suggestive of inner-shelf environments. The barnacle epifauna attached to a shark tooth AMU-CURS-720 (Figure 12.9-12.10) further indicates that the Cantaure Formation was deposited in a shallow marine environment, as barnacles live either in intertidal (26%) or sublittoral areas of ≤ 100 m (73%) (Doyle et al., 1996). Our bathymetric analysis of the Cantaure Formation, suggesting most likely an estimated deep interval depth between 0 to 50 m (Figure 13), supports the assumption of an inner-middle shelf environment (Figure 16). Additional support to this estimation is derived from the presence of the benthopelagic sharks *Nebrius* sp., *Carcharhinus* cf. *C. limbatus*, *C. perezii* and *Isogomphodon acuarius*, whose extant representatives usually prefer waters shallower than 40 m deep (Appendix 3). As shown in Figure 13, many of the benthopelagic and pelagic elasmobranch taxa of the Cantaure fauna have extant representatives associated with coastal environments, but can also occur in adjacent deep waters (Appendix 3), and even move significant distances over oceanic basins, suggesting a wide environmental range (Compagno et al., 2005; Voigt and Weber, 2011).

The Cantaure sequence rests on metamorphic and igneous rocks that were above water level during early Miocene (Gonzalez de Juana et al., 1980; Macellari, 1995). During the Cantaure sedimentation, these positive reliefs played the role of islands or archipelagos, as illustrated in the paleogeographic reconstruction of the Falcon Basin by Gonzalez de Juana et al. (1980) and Macellari (1995). According to Landau et al. (in press), 47.8% of the gastropod species (188 spp.) in the Cantaure Formation are exclusive, representing a very high level of endemism. This species endemism could support the hypothesis of an isolated, insular mollusk-community in the northern part of the Falcon Basin during early Miocene. However,




FIGURE 16. Paleobathymetric interpretation of the Cantaure Formation using several proxies. A, Foraminifera (Díaz de Gamero, 1974), B, corals (Griffiths et al., 2013), C, mollusks (Jung, 1965; Landau et al., in press), D, crustaceans (Aguilera et al., 2010), E, sharks (this study), F, teleosteans (Nolf and Aguilera, 1998, Aguilera and Rodrigues de Aguilera, 2001).

expanded knowledge of their localities will certainly lower that estimate. The Cantaure gastropod assemblage also has a small freshwater component, corresponding to typical species of late Oligocene early Middle Miocene NW South-American inland freshwater faunas (Landau et al. in press). This is likely suggestive of insular areas allowing the existence of terrestrial faunas. In addition, the diverse teleostean fauna from the Cantaure Formation, with more than 65 species (Nolf and Aguilera, 1998; Aguilera, 2010; Aguilera and Lundberg, 2010), suggests a high-productivity environment associated with a coastal upwelling (Aguilera and Rodrigues de Aguilera, 2004).

Although it is difficult to approach the precise ecological role that the Cantaure elasmobranch fauna played in this ancient marine environment, we can speculate about trophic interactions using the dietary (nutrition mode) categorization of taxa in the assemblage (Figure 14). The Cantaure

assemblage is dominated by piscivorous shark species (Figure 14, Appendix, 3). This could be related to the diverse teleostean fauna of the Cantaure Formation (Nolf and Aguilera, 1998; Aguilera, 2010; Aguilera and Lundberg, 2010), whose remains (mainly otoliths) are extremely abundant, with approximately more than 60 specimens per 10 kg of sifted matrix (OAA and JDCB pers. obs.). The durophagous/cancritrophic feeders represent the second most abundant elasmobranch group in the assemblage (Figure 14, Appendix 3). Their potential benthonic prey includes some 445 mollusks species (Jackson et al., 1999; Landau et al., in press) and more than 20 crustacean species (Aguilera et al., 2010) and must have offered wide feeding opportunities. Durophagous rays with capacity to triturate hard shells were abundant in the Cantaure fauna (e.g., *Rhinoptera* sp. being the third most abundant species of the assemblage, Appendix 1, 3). The presence of mobulids (Table 1, Appendix 1, 4), whose living counterparts are characterized by a diet based mainly on planktonic microorganisms (Klimley, 2013), support the assumption of a high-productive environment associated to a coastal upwelling (Aguilera and Rodrigues de Aguilera, 2004).

A few large sharks from the Cantaure assemblage such as *Carcharocles megalodon*, *Galeocerdo aduncus* and *G. mayumbensis*, could have alternatively occupied different environments and fed on various resources, such as teleosteans, chondrichthyans, reptiles, birds and mammals. In extant marine communities many sharks are perceived as apex predators, occupying the same trophic level with marine mammals (Klimley, 2013). In the Cantaure fauna such large sharks (*C. megalodon*, *G. aduncus* and *G. mayumbensis*), together with a medium-sized odontocete (?*Squalodelphinidae*) (Cozzuol and Aguilera, 2008; likely occupied the apex predator role. Moreover, cetacean (squalodontids) and sirenid (dugongids) skeletal remains found in the Cantaure Formation exhibit shark bite marks, suggesting that these mammals also were prey (Cozzuol and Aguilera, 2008). A cetacean lumbar vertebra with an embedded tooth of a *Carcharocles megalodon* was reported from the early Pliocene Paraguaná Formation, in the “El Yacural” locality (Aguilera et al., 2008). Recent field work in the “El Yacural” produced new specimens that include cetacean remains (vertebra and a skull fragment; AMU-CURS-709 and 710, respectively) and a crocodilian vertebra (AMU-CURS-708).

Those specimens lack any evidence for predation. The geographic coordinates of locality 4

(Figure 1.1) are very close to the "El Yacural" locality studied by Aguilera et al. (2008), with a horizontal distance of less than 200 m. Both localities are overlain stratigraphically by limestone layers on the top of the Cerro San Carlos (OAA, JDCB, GAF, pers. obs.), which have been interpreted by Jung (1965) and Hunter and Bartok (1974) as the top of the Cantaure Formation, representing the transition between the latter unit and post-Cantaure and pre-Pliocene sedimentation. We tentatively suggest that the vertebra with an embedded *Carcharocles megalodon* tooth could belong to the Cantaure Formation and not to the early Pliocene Paraguaná Formation (Aguilera et al., 2008). As previously mentioned, the Cantaure Formation is a complex sequence that needs a new detailed stratigraphic redescription, which could offer new geological information. The sections located at the south of the Barbasco Creek are of particular relevance for future research. In addition, the crocodilian vertebra (AMU-CURS-708) and the small eroded turtle-shell fragments, although not offering useful information for taxonomical classification, verify the presence of at least two reptiles in this ancient high-productive marine environment of the Cantaure Formation.

CONCLUSIONS

The elasmobranch fauna from the Cantaure Formation, containing 39 species, is characterized by a predominance of benthopelagic sharks with piscivorous feeding preferences, followed by durophagous/carcitrophic feeders. The elasmobranch habitat preferences and paleobathymetric analyses support the hypothesis that the Cantaure Formation was deposited in a high productivity, inner-middle shelf environment, around positive reliefs (island/archipelago), in the northern part of Falcon Basin during early-middle Miocene. This shark and ray fauna, is one of most diverse assemblages from early Neogene of the Americas, and contains many forms now extinct in the Caribbean, but present in the eastern Atlantic and western Pacific. Crocodile and turtle remains are also preserved in the Cantaure Formation.

ACKNOWLEDGEMENTS

This work was supported by Swiss National Science Foundation SNF 31003A-149605 to MRSV and by the Smithsonian Tropical Research Institute (Panama Paleontological Project) to OAA. The authors wish to especially thank T. Argyriou, B. Landau, J. Reyes, D. Rodrigues de Aguilera, A.

Coates, Dr. H. Cappetta, S. Adnet, R. Sánchez, C. Moron, H. Moreno, I. de Jesus, D. Cruz and A.E. Reyes for their generous and important counseling and collaboration; the Instituto del Patrimonio Cultural de Venezuela, for the authorization and permissions to collect, transport and study fossil material. Special thanks to R. Roth, M. Kirschmann and the Center for Microscopy and Image Analysis of the University of Zurich for their assistance and support performing the scanning electron microscopy analysis. The Alcaldía Bolivariana de Urumaco, the Universidad Experimental Francisco de Miranda, Museo de Ciencias Naturales de Caracas in Venezuela; Mapuka Museum of Universidad del Norte (Barranquilla, Colombia), Universidade Federal Fluminense, Universidade Federal do Pará and Museu Paraense Emílio Goeldi (Brazil), Natural History Museum of Basel (Switzerland), Paleontological collection of the Institut des Sciences de l'Évolution, University of Montpellier (France) and Palaeontological Institute and Museum at the University of Zurich for their valuable assistance and for access to comparative material. We greatly appreciate comments and suggestions from two anonymous reviewers.

REFERENCES

- Adnet, S., Cappetta, H., Guinot, G., and Notarbartolo Di Sciara, G. 2012. Evolutionary history of the devilrays (Chondrichthyes: Myliobatiformes) from fossil and morphological inference. *Zoological Journal of the Linnean Society*, 166(1):132-159.
- Agassiz, L. 1833–1843. *Recherches sur les poisons fossils*, 3, Neuchatel.
- Aguilera, O.A. 2010. *Peces Fósiles del Caribe de Venezuela*. Gorham Printing, Washington.
- Aguilera, O.A. and Rodrigues de Aguilera, D. 2004. Giant-toothed white sharks and wide-toothed mako (Lamnidae) from the Venezuela Neogene: Their role in the Caribbean, shallow-water fish Assemblage. *Caribbean Journal of Science*, 40(3):368-382.
- Aguilera, O.A. and Lundberg, J.G. 2010. Venezuelan Caribbean and Orinocoan Neogene fish, p. 129-152. In Sánchez-Villagra, M.R., Aguilera, O.A., and Carlini, F. (eds.), *Urumaco and Venezuelan Paleontology*. Indiana Press University, Bloomington.
- Aguilera, O.A., García, L., and Cozzuol, M. 2008. Giant-toothed white sharks and cetacean trophic interaction from the Pliocene Caribbean Paraguaná Formation. *Paläontologische Zeitschrift*, 82(2):204-208.
- Aguilera, O.A., Rodrigues de Aguilera, D., Vega, F.J., and Sánchez-Villagra, M.R. 2010. Mesozoic and Cenozoic decapod crustaceans from Venezuela and related trace-fossil assemblages, p. 103-128. In Sánchez-Villagra, M.R., Aguilera, O.A., and Carlini,

- F. (eds.), *Urumaco and Venezuelan Paleontology*. Indiana Press University, Bloomington.
- Aguilera, O.A., Ramos, M.I.F., Paes, E.T., Costa, S.A.R.F., and Sánchez-Villagra, M.R. 2011. The Neogene tropical America fish assemblage and the paleobiogeography of the Caribbean region. *Swiss Journal of Palaeontology*, 130(2):217-240.
- Alván, A., Apolín, J., and Chacaltana, C. 2006. Los dientes de seláceos (Chondrichthyes) y su aplicación estratigráfica en la lomas de Ullujaya (Ica-Perú). In: *XIII Congreso Peruano de Geología, Lima, 17-20 octubre, 2006*:595-598.
- Ameghino, F. 1901. L'âge des formations sédimentaires de Patagonie. *Anales de la Sociedad Científica Argentina*, 51:20-39, 65-91.
- Applegate, S.P. 1972. A revision of the higher taxa of Orectoloboids. *Journal of the Marine Biological Association of India*, 14(2):743-751.
- Berg, L.S. 1937. A classification of fish-like vertebrates. *Bulletin of the Academy of Sciences of the USSR, Division of Chemical Science*, 1937:1277-1280.
- Blainville, H.M.D. 1816. Prodrome d'une nouvelle distribution systematique du regne animal. *Bulletin de la Société Philomathique de Paris*, 8:105-112, 121-124.
- Blake, C.C. 1862. Shark's teeth at Panama. *Geologist*, 5:316.
- Bonaparte, C.L. 1838. Selachorum tabula analytica. *Nuovi Annali della Science Naturali Bologna*, 1:195-214.
- Bonnaterre, J.P. 1788. *Ichthyologie. Tableau encyclopédique et méthodique des trois règnes de la nature*, Paris.
- Bor, T.J., Reinecke, T., and Verschuere, S. 2012. Miocene Chondrichthyes from Winterswijk - Miste, the Netherlands. *Palaeontos*, 21:1-136.
- Cappetta, H. 1970. Les Sélaciens du Miocène de la région de Montpellier. *Palaeovertebrata, Mémoire Extraordinaire*, (1970):1-139.
- Cappetta, H. 1980. Modification du statut générique de quelques espèces de sélaciens crétacés et tertiaires. *Palaeovertebrata*, 10:29-42.
- Cappetta, H. 1987. *Chondrichthyes II: Mesozoic and Cenozoic Elasmobranchii. Handbook of Paleoichthyology*, 3B. Verlag Dr. Friedrich Pfeil, Munich.
- Cappetta, H. 2012. *Chondrichthyes. Mesozoic and Cenozoic Elasmobranchii: Teeth. Handbook of Paleoichthyology*, 3E. Verlag Dr. Friedrich Pfeil, Munich.
- Carrillo-Briceño, J.D., Aguilera, O.A., and Rodríguez, F. 2014. Fossil Chondrichthyes from the central eastern Pacific Ocean and their paleoceanographic significance. *Journal of South American Earth Sciences*, 51:76-90.
- Carrillo-Briceño, J.D., De Gracia, C., Pimiento, C., Aguilera, O.A., Kindlimann, R., Santamarina, P., and Jaramillo, C. 2015a. A new Late Miocene chondrichthyan assemblage from the Chagres Formation, Panama. *Journal of South American Earth Sciences*, 60:56-70.
- Carrillo-Briceño, J.D., Maxwell, E., Aguilera, O.A., Sánchez, R., and Sánchez-Villagra M.R. 2015b. Sawfishes and other elasmobranch assemblages from the Mio-Pliocene of the south Caribbean (Urumaco Sequence, Northwestern Venezuela). *PLoS ONE*, 10(10):e0139230. doi:10.1371/journal.pone.0139230.
- Carrillo-Briceño, J.D., Argyriou, T., Zapata, V., Kindlimann, R., and Jaramillo, C.A. 2016. A new Early Miocene (Aquitania) Elasmobranchii assemblage from the Guajira peninsula, Colombia. *Ameghiniana*, 53:77-99.
- Casier, E. 1958. Contribution à l'étude des poissons fossiles des Antilles. *Mémoire Suisse de Paléontologie*, 74:1-95.
- Casier, E. 1966. Sur la faune ichthyologique de la Formation de Bissex Hill et de la Série océanique, de l'île de la Barbade, et sur l'âge de ces formations. *Eclogae Geologicae Helveticae*, 59:493-516.
- Cicimurri, D.J. and Knight, J.L. 2009. Late Oligocene sharks and rays from the Chandler Bridge Formation, Dorchester County, South Carolina, USA. *Acta Palaeontologica Polonica*, 54(4):627-647.
- Cione, A.L., Cozzuol, M.A., Dozo, M.T., and Acosta Hospitaleche, C. 2011. Marine vertebrate assemblages in the southwest Atlantic during the Miocene. *Biological Journal of the Linnean Society*, 103(2):423-440.
- Coates, A.G. and Stallard, R.F. 2013. How old is the Isthmus of Panama? *Bulletin of Marine Sciences*, 89(4):801-813.
- Compagno, L.J.V. 1973. Interrelationships of living elasmobranchs, p. 15-61. In Greenwood, P.H., Miles, R.S., and Patterson, C. (eds.), *Interrelationships of Fishes*. Academic Press, London.
- Compagno, L.J.V. 1984a. *Sharks of the World: An Annotated and Illustrated Catalogue of Shark Species Known to Date. Part 1: Hexanchiformes to Lamniformes*. FAO Species Catalogue, 4. Food and Agriculture Organization of the United Nations, Rome.
- Compagno, L.J.V. 1984b. *Sharks of the World. An Annotated and Illustrated Catalogue of Shark Species Known to Date. Part 2: Carcharhiniformes*. FAO Species Catalogue, 4. Food and Agriculture Organization of the United Nations, Rome.
- Compagno, L.J.V. 2005. *Appendix 1: Global Checklist of Living Fishes. Sharks, Rays and Chimaeras: The Status of the Chondrichthyan Fishes*. IUCN-The World Conservation Union, Oxford, UK.
- Compagno, L.J.V., Dando, M., and Fowler, S. 2005. *Sharks of the World*. Princeton University Press, Princeton.
- Cope, E.D. 1867. An addition to the vertebrate fauna of the Miocene period, with a synopsis of the extinct Cetacea of the United States. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 19:138-156.
- Cope, E.D. 1869. Descriptions of some extinct fishes previously unknown. *Proceedings of the Boston Society of Natural History*, 12:310-317.

- Cortés, E. 1999. Standardized diet compositions and trophic levels of sharks. *ICES Journal of Marine Science*, 56:707-717.
- Cortés, E., Papastamatiou, Y.P., Carlson, J.K., Ferry-Graham, L., and Wetherbee, B.M. 2008. An overview of the feeding ecology and physiology of elasmobranch fishes, p. 393-443. In Cyrino, J.E.P., Bureau, D.P., and Kapoor, B.G. (eds.), *Feeding and Digestive Functions of Fishes*. Taylor and Francis Group, Florida.
- Costa, S.A.F., Richter, M., Toledo, P.M., and Moraes-Santos, H.M. 2009. Shark teeth from Pirabas Formation (Lower Miocene), northeastern Amazonia, Brazil. *Boletín do Museo Paraense Emílio Goeldi, Ciências Naturais*, 4:221-230.
- Cozzuol, M.A. and Aguilera, O.A. 2008. Cetacean remains from the Neogene of northwestern Venezuela. *Paläontologische Zeitschrift*, 82(2):196-203.
- Cuvier, G.L.C.F.D. 1816. *Le Règne Animal distribué d'après son organisation pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée. Les reptiles, les poissons, les mollusques et les annélides*. Deterville, Paris.
- Cuvier, G.L.C.F.D. 1829. *Le règne animal, distribué d'après son organisation, pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée*. Chez Deeterville, Paris.
- Daimeries, A. 1889. Notes ichthyologiques. V. *Annales de la Société royale malacologique de Belgique. Bulletin des Séances*, 24:39-44.
- Dartevelle, E. and Casier, E. 1943. Les poissons fossiles du Bas-Congo et des régions voisines. *Annales du Musée du Congo Belge, Série. A (Minéralogie Géologie, Paléontologie)*, 2(1):1-200.
- Díaz de Gamero, M.L. 1974. Microfauna y edad de la Formación Cantaure, Península de Paraguaná, Venezuela. *Boletín Informativo, Asociación Venezolana de Geología*, 17:41-47.
- Doyle, P., Mather, A.E., Bennett, M.R., and Bussell, M.A. 1996. Miocene barnacle assemblages from southern Spain and their palaeoenvironmental significance. *Lethaia*, 29(3):267-274.
- Ebert, D.A. and Stehmann, M.F.W. 2013. *Sharks, Batoids, and Chimaeras of the North Atlantic. FAO Species Catalogue for Fishery Purposes, 7. Food and Agriculture Organization of the United Nations, Rome*.
- Fowler, H.W. 1910. Notes on batoid fishes. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 62:468-475.
- Froese, R. and Pauly, D. 2015. FishBase. World Wide Web electronic publication. www.fishbase.org, version (2015).
- Garman, S. 1913. The Plagiostomia (Sharks, Skates and Rays). *Memoirs of the Museum of Comparative Zoology at Harvard College*, 36:1-528.
- Gibbes, R.W. 1849. Monograph of the fossil Squalidae of the United States. *Journal of the Academy of Natural Sciences of Philadelphia*, 1(2):191-206.
- Gibson-Smith, J. 1974. On two new members of the family Oculidae (Mollusca: Gastropoda) from the Cantaura Formation, Venezuela. *Boletín Informativo, Asociación Venezolana de Geología, Minería y Petróleo*, 17(4-6):87-89.
- Gibson-Smith, J. and Gibson-Smith, W. 1974. The Genus *Strombina* in Venezuela, with description of a new Recent and some fossil species. *Boletín Informativo, Asociación Venezolana de Geología, Minería y Petróleo*, 17(4-6):49-85.
- Gill, T. 1862. Note on some genera of fishes of western North America. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 14:329-332.
- Gill, T. 1872. *Arrangement of the Families of Fishes or Classes Pisces, Marsipobranchii, and Leptocardi* Prepared for the Smithsonian Institution by Theodore Gill, 247. Smithsonian Institution Miscellaneous Collection.
- Gill, T. 1893. Families and subfamilies of fishes. *Memoirs of the National Academy of Sciences*, 6:125-138.
- Glikman, L.S. 1964. *Sharks of the Paleogene and Their Stratigraphic Significance*. Nauka Press, Moscow. (In Russian).
- González de Juana, C., Iturralde de Arozena, J.M., and Picard, C.X. 1980. *Geología de Venezuela y de sus Cuencas Petrolíferas*. Ediciones Foninves, Caracas.
- González-Rodríguez, K.A., Espinosa-Arrubarrena, L., and González-Barba, G. 2013. An overview of the Mexican fossil fish record, p. 9-34. In Arratia, G., Schultze, H.P., and Wilson, M.V.H. (eds.), *Mesozoic Fishes 5 – Global Diversity and Evolution*. Verlag Dr. Friedrich Pfeil, München, Germany.
- Gradstein, F.M., Ogg, J.G., Schmitz, M.D., and Ogg, G. 2012. *The Geologic Time Scale 2012*. Elsevier New York, pp. 437-1144.
- Gray, J. 1851. *List of the specimens of fish in the collection of the British Museum. Part 1. Chondropterygii*. British Museum (Natural History).
- Griffiths, N., Müller, W., Johnson, K.G., and Aguilera, O.A. 2013. Evaluation of the effect of diagenetic cements on element/Ca ratios in aragonitic Early Miocene (~ 16 Ma) Caribbean corals: Implications for 'deep-time' palaeo-environmental reconstructions. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 369:185-200.
- Günther, A. 1870. *Catalogue of the Fishes in the British Museum*, 8. British Museum (Natural History), London.
- Hasse, K.E. 1879. *Das natürliche System der Elasmobranchier auf Grundlage des Baues und der Entwicklung ihrer Wirbelsäule: eine morphologische und paläontologische Studie*, 1. Gustav Fischer Verlag, Jena.
- Hunter, V.F. and Bartok, P. 1974. The age and correlation of the Tertiary sediments of the Paraguaná Peninsula, Venezuela. *Boletín Informativo, Asociación Venezolana de Geología, Minería y Petróleo*, 17:143-154.

- Ingram, W.M. 1947. New fossil Cypræidae from Venezuela and Colombia. *Bulletins of American Paleontology*, 31(121):1-12.
- Iturralde-Vinent, M.A., Hubbell, G., and Rojas, R. 1996. Catalogue of Cuban fossil Elasmobranchii (Paleocene to Pliocene) and paleogeographic implications of their lower to middle Miocene occurrence. *Boletín de la Sociedad Jamaicana de Geología*, 31:7-21.
- Jackson, J.B.C., Todd, J.A., Fortunato, H., and Jung, P. 1999. Diversity and assemblages of Neogene Caribbean Mollusca of lower Central America, p. 193-230. In Collins, L.S. and Coates, A.G. (eds.), *A Paleobiotic Survey of the Caribbean Faunas from the Neogene of the Isthmus of Panama*. *Bulletins of American Paleontology*, Ithaca, NY.
- Jordan, D.S. 1888. Description of two new species of fishes from South America. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 39:387-388.
- Jordan, D.S. and Evermann, B.W. 1896. The fishes of North and Middle America: a descriptive catalogue of the species of fish-like vertebrates found in the waters of North America, north of the Isthmus of Panama. Part I. *Bulletin of the United States National Museum*, 47:1-1240.
- Jordan, D.S. and Hannibal, H. 1923. Fossil sharks and rays of the Pacific Slope of North America. *Bulletin of the Southern California Academy of Sciences*, 22:27-63.
- Jung, P. 1965. Miocene Mollusca from the Paraguaná Peninsula, Venezuela. *Bulletins of American Paleontology*, 49(223):387-644.
- Kent, B.W. 1994. *Fossil Sharks of the Chesapeake Bay Region*. Egan Ress & Boyer, Inc., Columbia, Maryland.
- Kent, B.W. and Powell, J.G.W. 1999. Reconstructed dentition of the rare lamnoid shark *Parotodus benedicti* (Le Hon) from the Yorktown Formation (Early Pliocene) at Lee Creek Mine, North Carolina. *Mosa-saur*, 6:1-10.
- Kiraly, S.J., Moore, J.A., and Jasinski, P.H. 2003. Deep-water and other sharks of the US Atlantic ocean exclusive economic zone. *Marine Fisheries Review*, 65(4):1-20.
- Klimley, P.A. 2013. *The Biology of Sharks and Rays*. The University of Chicago Press, Chicago.
- Kruckow, T. and Thies, D. 1990. Die Neoselachier der Paleokaribik (Pisces: Elasmobranchii). *Courier Forschungsinstitut Senckenberg*, 119:1-102.
- Landau, B.M., Silva, C.A. da., and Heitz, A. in press (Accepted in January 2016). Systematics of the Gastropods of the Lower-Middle Miocene Cantaure Formation, Paraguaná Peninsula, Venezuela. *Bulletins of American Paleontology*.
- Laurito, C.A. 1999. Los seláceos fósiles de la localidad de Alto Guayacán (y otros ictiolitos asociados), Mioceno superior-Plioceno inferior de la Formación Uscari, provincia de Limón, Costa Rica. Laurito, C. (ed.), San José.
- Laurito, M.C.A., Calvo, C., Valerio, A.L., Calvo, A., and Chacón, R. 2014. Ictiofauna del Mioceno inferior de la localidad de Pacuare de Tres Equis, formación río Banano, provincia de Cartago, Costa Rica, y descripción de un nuevo género y una nueva especie de scaridae. *Revista Geológica de América Central*, 50:153-192.
- Le Hon, H. 1871. *Préliminaires d'un mémoire sur les poissons tertiaires de Belgique*, Brussels.
- Leigh, E.G., O'Dea, A., and Vermeij, G.J. 2014. Historical biogeography of the Isthmus of Panama. *Biological Reviews*, 89(1):148-172.
- Lerliche, M. 1938. Contribution à l'étude des Poissons fossils des pays riverains de la Méditerranée américaine, Venezuela, Trinité, Antilles, Mexique. *Mémoires de la Société paléontologique du Suisse*, 61:1-52.
- Linck, H.F. 1790. Versuch einer Eintheilung der Fische nach den Zähnen. *Magazin für das Neueste aus der Physik und Naturgeschichte*, 6(3):28-38.
- Linnaeus, C. 1758. *Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species; cum characteribus, differentiis, synonymis, locis*. 10th Ed. Larentii Salvii, Stockholm.
- Macellari, C.E. 1995. Cenozoic sedimentation and tectonics of the southwestern Caribbean Pull-Apart Basin, Venezuela and Colombia, p. 757-780. In Tankard, A.J., Suárez, R., and Welsink, H.J. (eds.), *Petroleum basins of South America*. The American Association of Petroleum Geologists, Memoir.
- MacPhee, R.D.E., Iturralde-Vinent, M.A., and Gaffney, E.S. 2003. Domo de Zaza, an Early Miocene vertebrate locality in South-Central Cuba, with notes on the tectonic evolution of Puerto Rico and the Mona Passage. *American Museum Novitates*, 3394:1-42.
- Montes, C., Cardona, A., Jaramillo, C., Pardo, A., Silva, J.C., Valencia, V., Ayala, C., Pérez-Angel, L.C., Rodríguez-Parra, L.A., Ramirez, V., and Niño, H. 2015. Middle Miocene closure of the Central American Seaway. *Science*, 348(6231):226-229.
- Müller, A. 1999. Ichthyofaunen aus dem atlantischen Tertiär der USA. *Leipziger Geowissenschaften*, 9-10:1-360.
- Müller, J. and Henle, J. 1837. Gattungen der Haifische und Rochen nach einer von ihm mit Hr. Henle unternommenen gemeinschaftlichen Arbeit über die Naturgeschichte der Knorpelfische. *Akademie der Wissenschaften zu Berlin*, 1837(2):111-118.
- Müller, J. and Henle, J. 1839-1841. *Systematische Beschreibung der Plagiostomen*, Berlin.
- Musick, J.A., Harbin, M.M., and Compagno, L.J.V. 2004. Historical zoogeography of the Selachii, p. 33-78. In Carrier, J.C., Musick, J.A., and Heithaus, M.R. (eds.), *Biology of Sharks and Their Relatives*. CRC Press, Washington, D.C.
- Nolf, D. and Brzobohatý, R. 1994. Fish otoliths as paleobathymetric indicators. *Paleontologia y Evolució*, 24:255-264.
- Nolf, D. and Aguilera, O.A. 1998. Fish otoliths from the Cantaure Formation (Early Miocene of Venezuela).

- Bulletin de l'Institut Royal des Science Naturelles de Belgique, Sciences de la Terre*, 68:237-262.
- O'Dea, A., Jackson, J.B.C., Fortunato, H., Smith, J.T., D'Croz, L., Johnson, K.G., and Todd, J.A. 2007. Environmental change preceded Caribbean extinction by 2 million years. *Proceedings of the National Academy of Sciences*, 104(13):5501-5506.
- Pimiento, C. and Balk, M.A. 2015. Body-size trends of the extinct giant shark *Carcharocles megalodon*: a deep-time perspective on marine apex predators. *Paleobiology*, 41(03):479-490.
- Pimiento, C. and Clements, C.F. 2014. When did *Carcharocles megalodon* become extinct? A new analysis of the fossil record. *PLoS ONE*, 9(10):e111086.
- Pimiento, C., Ehret, D.J., MacFadden, B.J., and Hubbell, G. 2010. Ancient nursery area for the extinct giant shark *Megalodon* from the Miocene of Panama. *PLoS ONE*, 5(5):e10552.
- Pimiento, C., Gonzalez-Barba, G., Hendy, A.J.W., Jaramillo, C., MacFadden, B.J., Montes, C., Suarez, S.C., and Shippritt, M. 2013a. Early Miocene chondrichthyans from the Culebra Formation, Panama: A window into marine vertebrate faunas before closure the Central American Seaway. *Journal of South American Earth Sciences*, 42:159-170.
- Pimiento, C., González-Barba, G., Ehret, D.J., Hendy, A.J., MacFadden, B.J., and Jaramillo, C. 2013b. Sharks and rays (Chondrichthyes, Elasmobranchii) from the late Miocene Gatun Formation of Panama. *Journal of Paleontology*, 87(5):755-774.
- Poe, F. 1876. Enumeratio piscium cubensium (Parte III). *Anales de la Sociedad Española de Historia Natural*, 5:373-404.
- Portell, R.W., Hubbell, G., Donovan, S.K., Green, J.L., Harper, D.A., and Pickerill, R. 2008. Miocene sharks in the Kendeace and Grand Bay formations of Carriacou, The Grenadines, Lesser Antilles. *Caribbean Journal of Science*, 44(3):279-286.
- Probst, J. 1879. Beiträge zur Kenntniss der fossilen Fische aus der Molasse von Baltringen. Hayfische. *Jahreshefte des Vereins für vaterländische Naturkunde in Württemberg*, 35:127-191.
- Purdy, R., Clellan, J.H.M., Schneider, V.P., Applegate, S.P., Meyer, R., and Slaughter, R. 2001. The Neogene sharks, rays and bony fishes from Lee Creek Mine, Aurora, North Carolina. *Smithsonian Contributions to Paleobiology*, 90:71-202.
- Rafinesque, C.S. 1809-1810. *Caratteri di alcuni nuovi generi e nuove specie di animali e piante della Sicilia con varie osservazioni sopra i medesimi*. Per le stampe di Sanfilippo, Palermo.
- Reinecke, T., Louwye, S., Havekost, U., and Moths, H. 2011. The elasmobranch fauna of the late Burdigalian, Miocene, at Werder-Uesen, Lower Saxony, Germany, and its relationships with Early Miocene faunas in the North Atlantic, Central Paratethys and Mediterranean. *Palaeontos*, 20:1-170.
- Reis, M.A.F. 2005. Chondrichthyan fauna from the Pirabas Formation, Miocene of northern Brazil, with comments on paleobiogeography. *Anuário do Instituto de geociências*, 28(2):31-58.
- Rey, O.T. 1996. Estratigrafía de la Península de Paraguaná, Venezuela. *Revista de la Facultad de Ingeniería, Universidad Central de Venezuela*, 11(1):35-45.
- Rincón, A.D., Solórzano, A., Benammi, M., Vignaud, P., and McDonald, H.G. 2014. Chronology and geology of an Early Miocene mammalian assemblage in North of South America, from Cerro La Cruz (Castillo Formation), Lara State, Venezuela: implications in the 'changing course of Orinoco River' hypothesis. *Andean Geology*, 41(3):507-528.
- Rüppell, W.P.E.S.E. 1835-1838. *Fische des Rothen Meeres*, Frankfurt am Main.
- Sánchez-Villagra, M.R., Burnham, R.J., Campbell, D.C., Feldmann, R.M., Gaffney, E.S., Kay, R.F., Lozán, R., Purdy, R., and Thewissen, J.G.M. 2000. A new near-shore marine fauna and flora from the early Neogene of Northwestern Venezuela. *Journal of Paleontology*, 74(5):957-968.
- Santos, R.S. and Travassos, H. 1960. *Contribuição à Paleontologia do estado do Pará. Peixes fósseis da Formação Pirabas. Monografia*, 16. Ministerio da Agricultura, Departamento Nacional da Produção Mineral, Divisão de Geologia e Mineralogia Rio de Janeiro.
- Santos, R.S. and Salgado, M.S. 1971. Contribuição à paleontologia do estado do Pará. Novos restos de peixes da Formação Pirabas. *Boletim do Museu Paraense Emílio Goeldi*, 16:1-13.
- Snodgrass, R.E. and Heller, E. 1905. Papers from the Hopkins-Stanford Galapagos Expedition, 1898-1899. XVII. Shore Fishes of the Revillagigedo, Clipperton, Cocos and Galapagos Islands. *Proceedings of the Washington Academy of Sciences*, 6:333-427.
- Suárez, M.E., Encinas, A., and Ward, D. 2006. An Early Miocene elasmobranch fauna from the Navidad Formation, Central Chile, South America. *Cainozoic Research*, 4(1-2):3-18.
- Visaggi, C.C. and Godfrey, S.J. 2010. Variation in composition and abundance of Miocene shark teeth from Calvert Cliffs, Maryland. *Journal of Vertebrate Paleontology*, 30(1):26-35.
- Voigt, M. and Weber, D. 2011. *Field Guide for Sharks of the Genus Carcharhinus*. Verlag Dr. Friedrich Pfeil, München.
- Whitley, G.P. 1929. Additions to the check-list of the fishes of New South Wales. No. 2. *Australian Zoologist*, 5(4):353-357.
- Whitley, G.P. 1940. *The Fishes of Australia, Pt. I: The Sharks, Rays, Devilfish, and Other Primitive Fishes of Australia and New Zealand*. Royal Zoological Society of New South Wales, Sydney.
- Woodward, A.S. 1889. *Catalogue of the Fossil Fishes in the British Museum. Part. I. British Museum (Natural History)*, London.

APPENDIX 1.

Elasmobranchii of the Cantaure Formation and their record per localities, jaw position and tooth measurements. Abbreviations: number of specimens (N°), total number of specimens (N° T) and indeterminate (indet.).

Taxon	Localities									Jaw position			Teeth measurements (mm)		
	1	N°	2	N°	3	N°	4	N°	N° T	Upper	Lower	indet.	Height	Width	Length
<i>Heterodontus</i> sp.	—	—	X	1	—	—	—	—	1	—	—	1	?	?	?
<i>Nebris</i> sp.	X	3	X	1	—	—	—	—	4	—	—	4	2 to 11	3 to 7	—
† <i>Paratodus benedenii</i>	—	—	X	1	—	—	—	—	1	1?	—	—	32	?	—
† <i>Carcharocles megalodon</i>	X	1	X	18	X	80	X	7	106	50	31	25	23 to 127	23 to ~ 66	—
<i>Alopias</i> cf. <i>Alopias vulpinus</i>	X	1	X	1	X	1	—	—	3	2	—	1	9 to 11	10 to 11	—
<i>Galeorhinus</i> cf. <i>G. galeus</i>	X	1	—	—	—	—	—	—	1	—	—	1	2.5	5	—
<i>Mustelus</i> sp.	—	—	X	3	—	—	—	—	3	—	—	3	<3	<3	<3
† <i>Hemipristis serra</i>	X	2	X	182	X	195	X	46	425	230	70	125	5 to 32	4.5 to 35	—
† <i>Galeocerdo aduncus</i>	X	1	X	13	—	—	—	—	14	—	—	14	11 to 18	16 to 24	—
† <i>Galeocerdo mayumbensis</i>	—	—	X	8	X	1	—	—	9	—	—	9	12 to 22	20 to 25.5	—
<i>Rhizoprionodon</i> sp.	X	3	X	14	—	—	—	—	17	4	6	7	2 to 3.8	2.2 to 4	—
† <i>Carcharhinus ackermanni</i>	X	5	—	—	—	—	—	—	5	3	2	—	11 to 18.2	12.7 to 20	—
<i>Carcharhinus brachyurus</i>	—	—	X	32	—	—	—	—	32	29	3	—	8 to 14	8 to 16.1	—
<i>Carcharhinus falciformis</i>	—	—	X	85	—	—	X	17	102	96	6	—	3 to 17	3 to 15.2	—
<i>Carcharhinus</i> cf. <i>C. galapagensis</i>	—	—	—	—	X	17	—	—	17	17	—	—	12 to 17.5	13 to 16	—
† <i>Carcharhinus gibbesii</i>	—	—	X	2	—	—	—	—	2	2	—	—	9 to 9.9	7.8 to 8	—
<i>Carcharhinus</i> cf. <i>C. limbatus</i>	X	2	—	—	—	—	—	—	2	2	—	—	5 to 9	6.5 to 11	—
<i>Carcharhinus</i> cf. <i>C. macrotis</i>	—	—	X	2	—	—	—	—	2	2	—	—	3.4 to 6.3	4 to 6.7	—
<i>Carcharhinus perezii</i>	X	11	X	3	X	1	—	—	15	15	—	—	4 to 8	5 to 13	—
<i>Carcharhinus</i> sp. 1	X	8	—	—	—	—	—	—	8	8	—	—	2 to 4.5	3.2 to 6	—
<i>Carcharhinus</i> spp.	X	21	X	50	X	17	X	10	98	40	52	6	5 to 15	4 to 14	—
† <i>Isogomphodon acurarius</i>	X	8	X	26	—	—	—	—	34	18	16	—	3 to 10.1	3.1 to 9.5	—
† <i>Negaprion eurybathron</i>	X	13	X	101	X	437	X	35	586	280	306	—	6 to 19	6 to 22	—
† <i>Physogaleus contortus</i>	—	—	X	30	X	3	X	1	34	—	—	34	10 to 19	19 to 19	—
<i>Carcharhinidae</i> indet.	X	2	X	8	—	—	—	—	10	2	8	—	3 to 6.2	4.5 to 8	—
† <i>Sphyrna arambourgi</i>	X	14	X	13	—	—	—	—	27	14	12	1	3 to 7	4 to 7.2	—
<i>Sphyrna</i> cf. † <i>S. laevisima</i>	X	2	X	1	X	5	—	—	8	—	3	5	3.5 to 11.9	5.5 to 14.9	—
<i>Rhynchobatus</i> sp.	X	8	X	7	—	—	—	—	15	—	—	15	<4	<4	<4
<i>Pristis</i> sp.	X	5	X	13	X	1	—	—	19	—	—	19	<3	<3	<3
<i>Dasyatis</i> sp.	X	21	X	12	—	—	—	—	33	—	—	33	<3	<3	<3
cf. <i>Pteroplatytrygon</i> sp.	X	5	—	—	—	—	—	—	5	—	—	5	<3	<3	<3
cf. <i>Taenirops</i> sp.	X	1	—	—	—	—	—	—	1	—	—	1	<3	<3	<3
<i>Dasyatidae</i> indet.	X	2	—	—	—	—	—	—	2	—	—	2	<3	<3	<3
<i>Aetobatus</i> sp.	X	6	X	9	X	4	—	—	19	—	—	19	4.1 to 6	?	~ 4.5 to 10
<i>Aetomylaeus</i> sp.	X	2	X	1	—	—	—	—	3	—	—	3	?	?	?
<i>Rhinoptera</i> sp.	X	15	X	129	X	1	—	—	145	—	—	145	2 to 6	5 to 29	3 to 8
† <i>Mobula fragilis</i>	X	4	—	—	—	—	—	—	4	—	—	4	2.5 to 3	3.5 to >6	1 to 1.6
<i>Mobula</i> cf. † <i>M. loupianensis</i>	X	3	—	—	—	—	—	—	3	—	—	3	<2	<2	<2
<i>Mobula</i> sp.	—	—	X	3	—	—	—	—	3	—	—	3	<2	<2	<2
† <i>Plinthiscus stenodon</i>	—	—	—	—	X	1	—	—	1	—	—	1	8	15.5	4
<i>Myliobatiformes</i> indet.	—	—	X	11	—	—	—	—	11	—	—	—	?	?	?
<i>Chondrichthyes</i> indet.	—	—	X	6	—	—	—	—	6	—	—	—	?	?	?
Total:		170		786		764		116	1836						

APPENDIX 2.

Referred fossil specimens and collection numbers.

Referred fossil specimens

- *Nebrius* sp. [1 tooth (AMU-CURS-982), 1 tooth (AMU-CURS-986)].
- †*Paratodus benedenii* [1 tooth (AMU-CURS-645)].
- †*Carcharocles megalodon* [4 teeth (AMU-CURS-512), 3 teeth (AMU-CURS-513), 2 teeth (AMU-CURS-514), 2 teeth (AMU-CURS-515), 3 teeth (AMU-CURS-516), 4 teeth (AMU-CURS-517), 3 teeth (AMU-CURS-518), 1 tooth (AMU-CURS-519), 4 teeth (AMU-CURS-650), 1 tooth (AMU-CURS-996), 1 tooth (AMU-CURS-997), 1 tooth (AMU-CURS-998), 7 teeth (AMU-CURS-1034), 1 tooth (UNEFM-PF-321), 1 tooth (UNEFM-PF-322), 13 teeth (UNEFM-PF-s/n), 17 teeth (UNEFM-PF-s/n), 21 teeth (UNEFM-PF-s/n), 17 teeth (MCNC-s/n)].
- *Alopias* cf. *A. vulpinus* [1 tooth (AMU-CURS-983), 1 tooth (AMU-CURS-984), 1 tooth (AMU-CURS-985)].
- *Galeorhinus* cf. *G. galeus* [1 tooth (AMU-CURS-974)].
- *Mustelus* sp. [3 teeth (AMU-CURS-975)].
- †*Hemipristis serra* [66 teeth (AMU-CURS-502), 1 tooth (AMU-CURS-506), 6 teeth (AMU-CURS-520), 24 teeth (AMU-CURS-644), 3 teeth (AMU-CURS-712), 1 tooth (AMU-CURS-731), 1 tooth (AMU-CURS-973), 1 tooth (AMU-CURS-993), 1 tooth (AMU-CURS-994), 1 tooth (AMU-CURS-1014), 46 1 tooth (AMU-CURS-1031), 156 (UNEFM-PF-s/n), 118 teeth (UNEFM-PF-s/n)].
- †*Galeocерdo aduncus* [13 teeth (AMU-CURS-647), 1 tooth (AMU-CURS-730)].
- †*Galeocерdo mayumbensis* [8 teeth (AMU-CURS-646), 1 tooth (AMU-CURS-995)].
- *Rhizoprionodon* sp. [14 teeth (AMU-CURS-959), 3 teeth (AMU-CURS-960)].
- †*Carcharhinus ackermanni* [3 teeth (AMU-CURS-713), 2 teeth (AMU-CURS-714)].
- *Carcharhinus brachyurus* [28 teeth (AMU-CURS-668), 4 teeth (AMU-CURS-990)].
- *Carcharhinus falciformis* [72 teeth (AMU-CURS-669), 6 teeth (AMU-CURS-991), 11 teeth (AMU-CURS-953), 17 teeth (AMU-CURS-1035)].
- *Carcharhinus* cf. *C. galapagensis* [17 teeth (AMU-CURS-973)].
- †*Carcharhinus gibbesii* [2 teeth (AMU-CURS-958)].
- *Carcharhinus* cf. *C. limbatus* [2 teeth (AMU-CURS-944)].
- *Carcharhinus* cf. *C. macroti* [2 teeth (AMU-CURS-957)].
- *Carcharhinus perezii* [1 tooth (AMU-CURS-954), 3 teeth (AMU-CURS-955), 11 teeth (AMU-CURS-956)].
- *Carcharhinus* sp. 1 [8 teeth (AMU-CURS-943)].
- *Carcharhinus* spp. [22 teeth (AMU-CURS-651), 26 teeth (AMU-CURS-652), 14 teeth (AMU-CURS-940), 7 teeth (AMU-CURS-941), 10 teeth (AMU-CURS-942), 7 teeth (AMU-CURS-972), 1 tooth (AMU-CURS-1015), 1 tooth (AMU-CURS-1017), 10 teeth (AMU-CURS-1085)].
- †*Isogomphodon acuarius* [8 teeth (AMU-CURS-950), 25 teeth (AMU-CURS-951), 1 tooth (AMU-CURS-952)].
- †*Negaprion eurybathrodon* [165 teeth (AMU-CURS-503), 65 teeth (AMU-CURS-504), 56 teeth (AMU-CURS-505), 1 tooth (AMU-CURS-509), 47 teeth (AMU-CURS-649), 13 teeth (AMU-CURS-987), 1 tooth (AMU-CURS-998), 52 teeth (AMU-CURS-989), 1 tooth (AMU-CURS-1013), 35 (AMU-CURS-1032), 150 teeth (UNEFM-PF-s/n)].
- †*Physogaleus contortus* [29 teeth (AMU-CURS-648), 2 teeth (AMU-CURS-992), 1 tooth (AMU-CURS-1033), 1 tooth (AMU-CURS-719), 1 tooth (AMU-CURS-720)].

APPENDIX 2 (CONTINUED).

- Carcharhinidae indet. [2 teeth (AMU-CURS-938), 8 teeth (AMU-CURS-939)].
- †*Sphyrna arambourgi* [14 teeth (AMU-CURS-961), 13 teeth (AMU-CURS-962)].
- Sphyrna* cf. †*S. laevissima* [1 tooth (AMU-CURS-963), 5 teeth (AMU-CURS-964), 1 tooth (AMU-CURS-965)].
- Rhynchobatus* sp. [4 teeth (AMU-CURS-653), 2 teeth (AMU-CURS-654), 3 teeth (AMU-CURS-968), 4 teeth (AMU-CURS-969), 1 tooth (AMU-CURS-970), 1 tooth (AMU-CURS-971)].
- Pristis* sp. [7 rostral denticles and 1 rostrum fragment (AMU-CURS-242), 1 rostral denticle (AMU-CURS-717), 3 teeth (AMU-CURS-655), 1 tooth (AMU-CURS-1019), 1 tooth (AMU-CURS-966), 5 teeth (AMU-CURS-717)].
- Dasyatis* sp. [1 tooth (AMU-CURS-657), 1 tooth (AMU-CURS-658), 12 teeth (AMU-CURS-929), 5 teeth (AMU-CURS-930), 10 teeth (AMU-CURS-931), 1 tooth (AMU-CURS-932), 1 tooth (AMU-CURS-933), 1 tooth (AMU-CURS-934), 1 tooth (AMU-CURS-935)].
- cf. *Pteroplatytrygon* sp. [5 teeth (AMU-CURS-926)].
- cf. *Taeniurops* sp. [1 tooth (AMU-CURS-925)].
- Dasyatidae indet. [1 tooth (AMU-CURS-936), 1 tooth (AMU-CURS-937)].
- Aetobatus* sp. [1 tooth (AMU-CURS-508), 1 tooth (AMU-CURS-659), 4 teeth (AMU-CURS-663), 4 teeth (AMU-CURS-665), 6 teeth (AMU-CURS-716), 2 teeth (AMU-CURS-718), 1 tooth (AMU-CURS-729)].
- Aetomylaeus* sp. [1 tooth (AMU-CURS-750), 2 teeth (AMU-CURS-766)].
- Rhinoptera* sp. [80 teeth (AMU-CURS-660), 1 tooth (AMU-CURS-664), 4 teeth (AMU-CURS-977), 6 teeth (AMU-CURS-978), 28 teeth (AMU-CURS-979), 11 teeth (AMU-CURS-980), 15 teeth (AMU-CURS-981)].
- †*Mobula fragilis* [2 teeth (AMU-CURS-947), 1 tooth (AMU-CURS-948), 1 tooth (AMU-CURS-949)].
- Mobula* cf. †*M. loupianensis* [3 teeth (AMU-CURS-946)].
- Mobula* sp. [3 teeth (AMU-CURS-945)].
- †*Plinthiscus stenodon* [1 tooth (AMU-CURS-715)].
- Myliobatiformes indet. [10 caudal spines (AMU-CURS-507), 1 tooth (AMU-CURS-510), 1 tooth (AMU-CURS-1018)].
- Chondrichthyes indet. [4 vertebrae (AMU-CURS-511), 2 denticles (AMU-CURS-656)].


APPENDIX 3.

Bathymetric and feeding preferences of the Cantaure Formation elasmobranch taxa, based on the biology of their extant relatives. The gray shadow indicates the feeding preferences. Abbreviations: minimum (Mn), maximum (Mx) and meters (m).

Cantaure Formation taxa	Extant species	Habitat of living representatives						Feeding ecology								
								Behavior		Diet						
		Benthic	Benthopelagic	Pelagic	Neritic / Epipelagic	Bathyal / Mesobathypelagic	Bathymetry depth (m)		Comments	Filter feeders	Benthic / pelagic predators	Piscivorous	Durophagous / Cancritophic	Teuthitrophic	Eurytrophic / Sarcophagous	Mammalian
Mn	Mx															
<i>Heterodontus</i> sp.	<i>Heterodontus</i> spp.	X			X	X	0 280			X	X	X				
<i>Nebrius</i> sp.	<i>Nebrius ferrugineus</i>	X			X		1 70	usually less than 30 m		X	X	X	X			
† <i>Paratodus benedenii</i>	—			X	X	?	?	?		X	X				?	
† <i>Carcharocles megalodon</i>	—		?	X	X	?	0 ?				X				X	
<i>Alopias</i> cf. <i>A. vulpinus</i>	<i>Alopias vulpinus</i>			X	X	X	0 550	usually less than 200 m			X		X			
<i>Galeorhinus</i> cf. <i>G. galeus</i>	<i>Galeorhinus galeus</i>		X		X	X	0 1100	usually 2-400 m		X	X	X	X	X		
<i>Mustelus</i> sp.	<i>Mustelus</i> spp.		X		X	X	0 900				X	X	X			
† <i>Hemipristis serra</i>	<i>Hemipristis elongata</i>		X		X		0 ~130			X	X			?	?	
† <i>Galeocerdo aduncus</i>	<i>Galeocerdo cuvier</i>		X		X	?	0 ~370	usually less than 150 m		X	X	X	X	X	X	
† <i>Galeocerdo mayumbensis</i>	<i>Galeocerdo cuvier</i>		X		X	?	~370	usually less than 150 m		X	X	X	X	X	X	
<i>Rhizoprionodon</i> sp.	<i>Rhizoprionodon</i> spp.		X		X		0 500	usually less than 100 m		X	X	X	X	X		
† <i>Carcharhinus ackermanni</i>	<i>Carcharhinus</i> spp.		?	X	X	?	0 >100?			X	X					
<i>Carcharhinus brachyurus</i>	<i>Carcharhinus brachyurus</i>		X		X	X	0 ~360	usually less than 100 m		X	X			X		
<i>Carcharhinus falciformis</i>	<i>Carcharhinus falciformis</i>			X	X	X	0 4000	most in waters > 200 m		X	X			X		
<i>Carcharhinus</i> cf. <i>C. galapagensis</i>	<i>Carcharhinus galapagensis</i>		X		X		0 180			X	X					
† <i>Carcharhinus gibbesii</i>	<i>Carcharhinus</i> spp.		?	X	X	?	0 >100?			X	X			X		
<i>Carcharhinus</i> cf. <i>C. limbatus</i>	<i>Carcharhinus limbatus</i>		X		X		0 100	usually less than 30 m		X	X			X		
<i>Carcharhinus</i> cf. <i>C. macloiti</i>	<i>Carcharhinus macloiti</i>			X	X		0 170			X	X			X		
<i>Carcharhinus perezii</i>	<i>Carcharhinus perezii</i>		X		X		0 65	usually less than 30 m		X	X					
<i>Carcharhinus</i> sp. 1	<i>Carcharhinus</i> spp.		?	?	X	?	0 >100?				X					
<i>Carcharhinus</i> spp.	<i>Carcharhinus</i> spp.		X	X	X	?	0 >100?			X	?	?	?	?		
† <i>Isogomphodon acurarius</i>	<i>Isogomphodon oxyrhynchus</i>		X		X		0 40			X	X					
† <i>Negaprion eurybathron</i>	<i>Negaprion brevirostris</i>		X		X		0 92			X	X					
† <i>Physogaleus contortus</i>	—		?	X	X	?	?	?		X	X					
<i>Carcharhinidae</i> indet.	<i>Carcharhinidae</i> spp.		?	?	X	?	0 >100?			X	X					
† <i>Sphyrna arambourgi</i>	<i>Sphyrna</i> spp.		X		X	?	0 >100?			X	X					
<i>Sphyrna</i> cf. † <i>S. laevisima</i>	<i>Sphyrna zygaena</i>		X		X		0 >100?	usually less than 20 m		X	X					
<i>Rhynchobatus</i> sp.	<i>Rhynchobatus</i> spp.		X		X		0 ~60			X	X	X				
<i>Pristis</i> sp.	<i>Pristis</i> spp.		X		X		0 ~120	most in waters < 20 m		X	X	X				
<i>Dasyatis</i> sp.	<i>Dasyatis</i> spp.	X			X	X	0 ~470	most in waters < 50 m		X	X	X				
cf. <i>Pteroplatytrygon</i> sp.	<i>Pteroplatytrygon violacea</i>	X			X	X	0 381	usually 1-100 m		X	X	X		X		
cf. <i>Taeniurops</i>	<i>Taeniurops</i> spp.	X			X	X	10 500	usually 20-60 m		X	X	X				
<i>Dasyatidae</i> indet.	<i>Dasyatidae</i> spp.	X			X	X	0 ~470			X	X	X				
<i>Aetobatus</i> sp.	<i>Aetobatus</i> spp.		X		X		0 ~100	usually less than 40 m		X	X	X	X	X		
<i>Aetomylaeus</i> sp.	<i>Aetomylaeus</i> spp.		X				0 110			X	X	X	X	X		
<i>Rhinoptera</i> sp.	<i>Rhinoptera</i> spp.		X		X		0 <60	usually less than 30 m		X	X	X				
† <i>Mobula fragilis</i>	<i>Mobula</i> spp.			X	X	?	0 >200	usually less than 100 m	X							X
<i>Mobula</i> cf. † <i>M. loupianensis</i>	<i>Mobula</i> spp.			X	X	?	0 >200	usually less than 100 m	X							X
<i>Mobula</i> sp.	<i>Mobula</i> spp.			X	X	?	0 >200	usually less than 100 m	X							X
† <i>Plinthiscus stenodon</i>	—		X		X	?	?	?	X							X
<i>Myliobatiformes</i> indet.	—	?	?	?	?	?	?	?	?	?						
<i>Chondrichthyes</i> indet.	—	?	?	?	?	?	?	?								

APPENDIX 4.

Relative abundance of elasmobranchs in the Cantaure Formation. 1. Abundance percentage per locality per taxon. 2. Total abundance per locality. Ht: Heterodontiformes; Or: Orectolobiformes; La: Lamniformes; Ca: Carcharhiniformes; Ra: Rajiformes; My: Myliobatiformes; In: Indeterminate; Loc.: locality; †: extinct taxon. *: Percentage does not include *Carcharhinus* spp., Myliobatiformes indet. and Chondrichthyes indet.



CHAPTER 7

Conclusions and future perspectives

Conclusions and future perspectives

This dissertation, as a whole, with the description and publication of eleven elasmobranch assemblages from the Caribbean region (Colombia, Panama and Venezuela) and Eastern Pacific (Ecuador), plus the ongoing description of four more assemblages from the early Miocene of Colombia and Brazil (e.g. Carrillo-Briceño et al., and Aguilera et al., both in prep.), capitalizes on long-term, previous fieldwork efforts, providing new insights of the chondrichthyan paleodiversity inhabiting Tropical America during the Neogene. The new assemblages document shallow and deep-water habitats. Among the findings are 17 taxa reported for the first time for Tropical America. Fossil remains of chimaeras (Holocephali) were not found in our assemblages, being until now the record of *Chimaera* sp. from the late Miocene-Pliocene of Costa Rica (Laurito, 2008) and *Callorhinchus* cf. *C. callorhinchus* from the Pliocene of Peru (De Muizon and Devries, 1985), the only representatives of this group known from the Neogene of Tropical America. The comprehensive interpretation about habitat and biology of the living counterpart taxa of the fossil ones presented in the studied assemblages was the tool used for the understanding of the paleoecological and the paleoenvironmental conditions of these geologic units.

Previous to this contribution, knowledge about chondrichthyan assemblages of deep-water affinities from Tropical America was very limited, with only general descriptions from the late Miocene-Pliocene of Costa Rica and Venezuela (Laurito, 1999; Aguilera and Rodríguez de Aguilera, 2001). Thus the novelty in this work represented by the description of new faunas associated with deep-water affinities from Colombia (Uitpa Formation), Ecuador (Angostura Formation) and Panama (Chagres Sandstone Member of the Chagres Formation) (chapters 2-3 and 5), including the description of the new species of the frilled shark †*Chlamydoselachus landinii* sp. nov. (Figure 1A; Chapter 2).

Fossil assemblages characterized by taxa with shallow-water affinities from the Neogene of Tropical America are the most common ones referred in previous contributions (e.g. Leriche, 1938; Santos and Travassos, 1960; Santos and Salgado, 1971; Kindlimann, 1990; Kruckow and Thies, 1990; Iturralde-Vinent et al., 1996; Reis, 2005; Alván et al., 2007; Costa et al., 2009; Aguilera, 2010; Pimiento et al., 2013a, 2013b). The assemblages described in chapters 2-4 and 6, from mainly Miocene and Pliocene units of Ecuador (Onzole, Canoa and Jama formations), Panama (Rio Indio Member, Chagres Formation), and Venezuela (Cantaure, Socorro, Urumaco and Codore formations), provide new insights about neritic faunas and their associated paleoenvironments. An example constitute the assemblages from Ecuador (Onzole, Canoa and Jama formations), which represent the most diverse Neogene paleodiversity known from the Eastern Pacific coast of Tropical America. In the case of the assemblages from Venezuela, which include the description of the new carcharhinid †*Carcharhinus caquetius* sp. nov. (Figure 1B), the habitat preference of the living counterparts of the fauna recovered from the Urumaco sequence (Socorro, Urumaco and Codore formations), suggest marine shallow waters and estuarine habitats associated with a geographic area connected with a hydrographic system that flowed from western Amazonia up to the Proto-Caribbean Sea during the Miocene (e.g. Sánchez-Villagra et al., 2010; Aguilera et al., 2013). These assemblages from the Urumaco sequence could be useful to understand the ecological role that elasmobranchs played in these ancient transitional environments (Aguilera and Rodrigues de Aguilera, 2004; Aguilera 2010). In addition, the assemblage from the Cantaure Formation, which is hypothetically associated to a shallow-insular high productive marine environment, represents the most diverse elasmobranch fauna known to the early Miocene of the America continent. Its paleodiversity characterized by 39 shark and ray species, with a predominance of benthopelagic sharks with piscivorous

durophagous/cancritrophic feeding preferences, represent a unique glimpse into elasmobranch dietary composition and food chain in the ancient proto-Caribbean Sea.




Figure 1. Specimens of new species of deep (A) and shallow (B) waters described in this work. A. *†Chlamydoselachus landinii* sp. nov. (PPP-3455-T-1), from the late Miocene of Ecuador, and B. *†Carcharhinus caquetius* sp. nov. (AMU-CURS-499), from the late Miocene of Venezuela.

The known Tropical America extant chondrichthyan diversity is characterized by a total of 276 species of chimaeras, sharks and rays (Eastern Pacific: 145 spp.; Western Atlantic 181 spp.; see Chapter 1). Notwithstanding the fragmentary nature of the fossil record, it can offer new insights to understand the origin and distribution pattern of the living species from this marine tropical area. One approximation to the chondrichthyan palaeodiversity from Tropical America is presented in the Figure 2, where a comprehensive Miocene-Pleistocene total faunal compilation based on new and published data is presented. The data were standardized at the generic level; in order to remove biases caused by taxonomic and nomenclatural uncertainties at the species level.

Throughout the Neogene the paleodiversity of Tropical America has not been the same, especially shark genera, in both Eastern Pacific and Western Atlantic (Figure 2). The differences could be the result to a significant extent of biases resulting from the scarce chondrichthyan fossil record known of the region, especially in the Eastern Pacific. The clear

difference between sharks and rays paleodiversity could be affected by biases introduced by sampling of dental elements, especially those associated with size. Usually, shark and ray (e.g. Myliobatidae and Rhinopteridae) teeth, due their bigger size, have more probabilities to be surface-collected directly from the outcrop. In contrast, dental elements of many ray and shark species [e.g. many species of Squaliformes and Carcharhiniformes (Scyliorhinidae)], are characterized by micro-teeth, requiring special collecting technics such as screen-washing and microscopically picking. With the exceptions of the assemblages described herein (chapters 2-6), and those previously referred by Laurito (1999), Aguilera and Rodrigues de Aguilera (2001) and Pimiento et al. (2013a, b), little was known about the Tropical America shark/ray paleodiversity based on micro-dental elements. In addition, an important contrast in the number of genera affecting both shark and ray paleodiversity, can also be observed in the Pleistocene fossil record (Figure 2). At first glance it appears that this low Pleistocene diversity could be related to an extinction process. However, the answer for this low paleodiversity is sampling bias. Pleistocene units from Tropical America with chondrichthyan assemblages are extremely rare, including only few taxa from Eastern Pacific (Ecuador and Panama) and Western Atlantic (Venezuela) (Aguilera, 2010; Chapter 1; Carrillo-Briceño and Sánchez-Villagra in prep.).




Figure 2. Neogene chondrichthyan paleodiversity of Tropical America in both Eastern Pacific and Western Atlantic. From Carrillo-Briceño and Sánchez-Villagra (in prep).

Tropical marine areas in both Eastern Pacific and Western Atlantic are environmentally heterogeneous (Spalding et al., 2007), with a marked differentiation referable to abiotic variables such as temperature, salinity, bathymetry, coastal length, currents, continental platform, rivers discharge to the oceans, among others (e.g. Miloslavich et al., 2011). These variables changed especially after the rise of the Central American land bridge (e.g. O'Dea et al., 2007a; Coates, and Stallard, 2013), and affected the marine biota (e.g. O'Dea et al., 2007a). The differential pattern in extant chondrichthyan diversity from Tropical America in both Eastern Pacific and Western Atlantic shown in Figure 4 (Chapter 1) could be a direct consequence of changes in the above mentioned variables, as well as other biotic/ecological variables (e.g. food resources, physiological conditions). Assuming the above, the question arises: Did the biotic/abiotic variables in both Eastern Pacific and Western Atlantic explain the chondrichthyan paleodiversity patterns shown in Figures 2-3, or are these mainly the result of sampling bias? The pattern surely reflects real phenomena. However, with the currently paleodiversity known (Figure 2), accurate interpretations are still preliminary. These strategies would be followed to improve this situation. 1) Increase the number of studied assemblages with accurate environmental (e.g. paleobathymetric analysis) and paleoecological inferences (e.g., dietary composition and feeding and habitat preferences). 2) Use of geochemical analysis (e.g. stable isotope on dental elements) of the known chondrichthyan assemblages, used to infer paleotemperatures and paleosalinity levels. Our ongoing work of the assemblages from the early Miocene of Brazil (Pirabas Formation) and Colombia (Jimol and Castilletes formation) (Carrillo-Briceño et al., and Aguilera et al., both in prep.), includes the use of stable isotope in sharks and rays teeth.




Figure 3. Shared/Unshared fossil genera in both the Eastern Pacific and Western Atlantic. From Carrillo-Briceño and Sánchez-Villagra (in prep).

Taking under consideration a plausible chondrichthyan faunal interchange through marine corridors in the Panamanian area before the definite closure of the isthmus (e.g. Montes et al., 2012, 2015; Coates and Stallard, 2013) we should expect a high taxonomic commonality between Eastern Pacific and Western Atlantic chondrichthyans, with a faunal turnover suggesting environmental changes towards the end of the Neogene. In contrast, Figure 3 shows a pattern where unshared genera (especially sharks) are dominant in all Miocene-Pliocene assemblages. The strong difference observed in early Miocene assemblages, could be assumed due the scarce Eastern Pacific assemblages known from this time (see chapter 6). The numbers of shared/unshared genera are more similar in middle Miocene-Pliocene assemblages; besides the fossil biases, the high proportion of unshared shark genera could be a response to differences in the proportion of the studied paleoenvironments. Most of the fossil assemblages known from Tropical America are

associated to shallow waters, while those associated to deep environments have been scarcely studied (chapters 2-3, 5). One example are Squalomorphii sharks (especially Hexanchiformes and Squaliformes taxa), associated to deep water environments, better known in the fossil record from the Western Atlantic assemblages (chapters 3, 5; Carrillo-Briceño and Sánchez-Villagra, in prep).

Our survey (Figures 2-3), still does not offer a direct biostratigraphic inference to support any of the two postulated timings proposed for the closure of the Panama Isthmus (e.g. Montes et al., 2012, 2015; Coates and Stallard, 2013). However, clues related with possible consequence of “post-Isthmus” environmental changes can be observed. Our analysis (Carrillo-Briceño and Sánchez-Villagra, in prep) reveal regional or complete extirpation of many Eastern Pacific and Western Atlantic shark/ray taxa (mainly neritic/epipelagic species) from Tropical America (Table 1), especially during late Miocene-Pliocene times. At least three genera became regionally extirpated in the Eastern Pacific (*Centrophorus*, *Pristiophorus* and *Carcharhias*) and two in the western Atlantic (*Heterodontus* and *Aetomylaeus*). Moreover, another 13 genera became extinct in both Eastern Pacific and Western Atlantic oceans (*Trigonognathus*, *Nebrius*, *Chiloscyllium*, †*Cosmopolitodus*, †*Carcharocles*, †*Anotodus*, *Hemipristis*, *Paragaleus*, †*Paratodus*, †*Pachyscyllium*, †*Kruckowlamna*, *Rhynchobatus*, and †*Plinthicus*). This late Neogene chondrichthyan extirpation/extinction in both Eastern Pacific and Western Atlantic relatively coincides with the extinction rate of other marine groups (corals and mollusks) in the Caribbean area (O'Dea et al., 2007a, fig. 3).

Taxon				Status		Habitat preferences	
				Eastern Pacific	Western Atlantic	Neritic/Epipelagic	Bathyal/Meso-Bathypelagic
Squalomorphii	Squaliformes	Squalidae	<i>Centrophorus</i>	RE	P		X
		Etmopteridae	<i>Trigonognathus</i>	RE	RE		X
	Pristiophoriformes	Pristiophoridae	<i>Pristiophorus</i>	RE	P	X	X
Galeomorphii	Heterodontiformes	Heterodontidae	<i>Heterodontus</i>	P	RE	X	
	Orectolobiformes	Hemiscyllidae	<i>Chiloscyllium</i>	RE	RE	X	
		Ginglymostomatidae	<i>Nebrius</i>	RE	RE	X	
	Lamniformes	Odontaspidae	<i>Carcharias</i>	RE	P	X	
		Lamnidae	† <i>Cosmopolitodus</i>	W-Ext		X	?
		†Otodontidae	† <i>Carcharocles</i>	W-Ext		X	?
			† <i>Paratodus</i>	W-Ext		X	?
			† <i>Anotodus</i>	W-Ext		X	?
	Carcharhiniformes	Scyliorhinidae	† <i>Pachyscyllium</i>	W-Ext		X	?
		Hemigaleidae	<i>Hemipristis</i>	RE	RE	X	
			<i>Paragaleus</i>	RE	RE	X	
		Carcharhinidae	† <i>Kruckowlamna</i>	W-Ext		X	?
Batiomorphii	Rajiformes	Rhynchobatidae	<i>Rhynchobatus</i>	RE	RE	X	
		Myliobatidae	<i>Aetomylaeus</i>	P	RE	X	
		Mobulidae	† <i>Plinthiscus</i>	W-Ext		X	?

Table 1. Extirpated/extinct chondrichthyan taxa from Tropical America. Status: Regional extirpated (RE), extant species presented on the region (P), species worldwide extinct (W-Ext). †: extinct taxon.

The future study of the chondrichthyan paleodiversity of Tropical America could be focused on these objectives:

- Continuation of the studies of fossil assemblages in both Eastern Pacific and Western Atlantic, especially in search of micro-dental elements, which could offer new information about poorly studied fossil groups, such as Rajiformes and Torpediniformes, among others, especially those with dental remains below a size of 2 mm (not sampled in the current work).

- Increased use of stable isotope analysis in fossil assemblages [e.g. the ongoing oxygen isotope composition analysis for assemblages from the early Miocene of Colombia and Brazil (Carrillo-Briceño et al., and Aguilera et al., both in prep.)], which are used as proxies for describing environmental and ecological conditions.

- Increased efforts in Pleistocene geological units, due the scarce diversity known from that period. It was in that time, after the definitive closure of the Isthmus of Panama, when different oceanographic events occurred in the region, triggering the beginning of the currently chondrichthyan diversity in both Eastern Pacific and Western Atlantic oceans. Added to this, the incursion of new methods, such as molecular analysis of chondrichthyan genera present in both side of the Isthmus of Panama, using time-calibrated hypothesis, could be useful to infer divergences and speciation patterns.

References

- Aguilera, O.A. 2010. Peces Fósiles del Caribe de Venezuela. Gorham Printing, Washington.
- Aguilera, O., and Rodrigues De Aguilera, D. 2001. An exceptional coastal upwelling fish assemblage in the Caribbean Neogene. *Journal of Paleontology*, 75(3):732-742.
- Aguilera, O.A., and Rodrigues de Aguilera, D. 2004. Giant-toothed White Sharks and Wide-toothed Mako (Lamnidae) from the Venezuela Neogene: Their Role in the Caribbean, Shallow-water Fish Assemblage. *Caribbean Journal of Science*, 40(3):368-382.
- Aguilera, O.A., Moraes-Santos, H., Costa, S., Ohe, F., Jaramillo, C., and Nogueira, A. 2013. Ariid sea catfishes from the coeval Pirabas (Northeastern Brazil), Cantaura, Castillo (Northwestern Venezuela), and Castilletes (North Colombia) formations (Early Miocene), with description of three new species. *Swiss Journal of Palaeontology*, 132(1):45-68.

- Alván, A., Apolín, J., and Chacaltana, C. 2006. Los dientes de seláceos (Chondrichthyes) y su aplicación estratigráfica en la lomas de Ullujaya (Ica-Perú). In Perú, S.G.P. (ed.), XIII Congreso Peruano de Geología, Lima, 17-20 octubre, 2006, pp. 595-598.
- Coates, A.G., and Stallard, R.F. 2013. How old is the Isthmus of Panama? . *Bulletin of Marine Sciences* 89(4):801–813.
- Costa, S.A.F., Richter, M., Toledo, P.M., and Moraes-Santos, H.M. 2009. Shark teeth from Pirabas formation (Lower Miocene), northeastern Amazonia, Brazil. *Boletín do Museu Paraense Emílio Goeldi, Ciencias Naturais* 4:221-230.
- De Muizon, C., and Devries, T.J. 1985. Geology and paleontology of late Cenozoic marine deposits in the Sacaco area (Peru). *Geologische Rundschau*, 74(3):547-563.
- Iturralde-Vinent, M.A., Hubbell, G., and Rojas, R. 1996. Catalogue of Cuban fossil Elasmobranchii (Paleocene to Pliocene) and paleogeographic implications of their lower to middle Miocene occurrence. *Boletín de la Sociedad Jamaicana de Geología* 31:7-21.
- Kindlimann, R. 1990. Selacios del Terciario Tardío de Sacaco, Departamento de Arequipa. *Boletín de Lima* 69:91-95.
- Kruckow, T., and Thies, D. 1990. Die Neoselachier der Paleokaribik (Pisces: Elasmobranchii). *Courier Forschungsinstitut Senckenberg* 119:1-102.
- Laurito, C.A. 1999. Los seláceos fósiles de la localidad de Alto Guayacán (y otros ictiolitos asociados), Mioceno superior-Plioceno inferior de la Formación Uscari, provincia de Limón, Costa Rica. Laurito, C. (Ed.), San José.

- Laurito, M.C.A. 2008. Additional fish record from the Uscari Formation, Upper Miocene-Lower Pliocene of Costa Rica: a Chimaeroidei tooth plate (Pisces, Chondrichthyes, Holocephali). *Revista Geológica de América Central*, 38:135-137.
- Leriche, M. 1938. Contribution à L'étude des Poissons fossils des pays riverains de la Méditerranée américaine, Venezuela, Trinité, Antiles, Mexique. *Mémoires de la Société Paléontologique du Suisse* 61:1-52.
- Miloslavich, P., Klein, E., Díaz, J.M., Hernández, C.E., Bigatti, G., Campos, L., Artigas, F., Castillo, J., Penchaszadeh, P.E., Neill, P.E., Carranza, A., Retana, M.V., Díaz de Astarloa, J.M., Lewis, M., Yorio, P., Piriz, M.L., Rodríguez, D., Yoneshigue-Valentin, Y., Gamboa, L., and Martín, A. 2011. Marine Biodiversity in the Atlantic and Pacific Coasts of South America: Knowledge and Gaps. *PLoS ONE*, 6(1):e14631.
- Montes, C., Cardona, A., McFadden, R., Morón, S.E., Silva, C.A., Restrepo-Moreno, S., Ramírez, D.A., Hoyos, N., Wilson, J., and Farris, D. 2012. Evidence for middle Eocene and younger land emergence in central Panama: implications for Isthmus closure. *Geological Society of America Bulletin*, 124(5-6):780-799.
- Montes, C., Cardona, A., Jaramillo, C., Pardo, A., Silva, J.C., Valencia, V., Ayala, C., Pérez-Angel, L.C., Rodríguez-Parra, L.A., Ramirez, V., and Niño, H. 2015. Middle Miocene closure of the Central American Seaway. *Science*, 348(6231):226-229.
- O'Dea, A., Jackson, J.B.C., Fortunato, H., Smith, J.T., D'Croz, L., Johnson, K.G., and Todd, J.A. 2007a. Environmental change preceded Caribbean extinction by 2 million years. *Proceedings of the National Academy of Sciences*, 104(13):5501-5506.
- Pimiento, C., González-Barba, G., Hendy, A.J.W., Jaramillo, C., MacFadden, B.J., Montes, C., Suarez, S.C., and Shippritt, M. 2013a. Early Miocene chondrichthyans from the

- Culebra Formation, Panama: A window into marine vertebrate faunas before closure the Central American Seaway. *Journal of South American Earth Sciences*, 42:159-170.
- Pimiento, C., González-Barba, G., Ehret, D.J., Hendy, A.J., MacFadden, B.J., and Jaramillo, C. 2013. Sharks and rays (Chondrichthyes, Elasmobranchii) from the late Miocene Gatun Formation of Panama. *Journal of Paleontology*, 87(5):755-774.
- Reis, M.A.F.d. 2005. Chondrichthyan fauna from the Pirabas Formation, Miocene of northern Brazil, with comments on paleobiogeography. *Anuário do Instituto de Geociências*, 28(2):31-58.
- Sánchez-Villagra, M.R., Burnham, R.J., Campbell, D.C., Feldmann, R.M., Gaffney, E.S., Kay, R.F., Lozsán, R., Purdy, R., and Thewissen, J.G.M. 2000. A New Near-Shore Marine Fauna and Flora from the Early Neogene of Northwestern Venezuela. *Journal of Paleontology*, 74(5):957-968.
- Sánchez-Villagra, M.R., Aguilera, O.A., and Carlini, F. (eds.). 2010. *Uruguay and Venezuelan Paleontology*. Indiana Press University, Bloomington.
- Santos, R.S., and Travassos, H. 1960. Contribuição à Paleontologia do estado do Pará. Peixes fósseis da Formação Pirabas. Monografia, 16. Ministério da Agricultura, Departamento Nacional da Produção Mineral, Divisão de Geologia e Mineralogia Rio de Janeiro.
- Santos, R.S., and Salgado, M.S. 1971. Contribuição à paleontologia do estado do Pará. Novos restos de peixes da Formação Pirabas. *Boletim do Museu Paraense Emílio Goeldi*, 16:1-13.

APPENDIX I

Late Neogene elasmobranch fauna from the Coquimbo Formation, Chile

LATE NEOGENE ELASMOBRANCH FAUNA FROM THE COQUIMBO FORMATION, CHILE

FELIPE STAIG

Departamento de Biología Marina, Facultad de Ciencias del Mar, Universidad Católica del Norte, Coquimbo, Chile.
fstaig@gmail.com

SEBASTIÁN HERNÁNDEZ

Sala de Colecciones Biológicas, Facultad de Ciencias del Mar, Universidad Católica del Norte, Larrondo 1281, Coquimbo, Chile. Veritas University, Center for International Programs, Costa Rica. *shernandez@veritas.cr*

PATRICIO LÓPEZ

Departamento de Antropología, Facultad de Ciencias Sociales, Universidad de Chile. Centre for Maritime Archaeology Research of the Southeastern Pacific, Santiago, Chile. *patriciolopezmendoza@gmail.com*

JAIME A. VILLAFANA

Programa de Magister en Ciencias del Mar and Laboratorio de Paleobiología, Centro de Estudios Avanzados en Zonas Áridas, Coquimbo, Chile. *jaime.villafana@ceaza.cl*

CRISTIAN VARAS

Subsecretaría de Pesca, Avenida Ossandón, 890, Coquimbo, Chile.
cvaras@subpesca.cl

LUIS PATRICIO SOTO

Florida Museum of Natural History, University of Florida, Gainesville, Florida, USA.
luispatricio.soto@gmail.com

JORGE D. CARRILLO-BRICEÑO

Paleontological Institute and Museum, University of Zürich, Karl-Schmid-Strasse 4, Zürich, 8006, Switzerland.
jorgedcb100@gmail.com

ABSTRACT – Neogene marine sediments from Chilean geological formations contain a diverse marine fossil fauna. In Chile, the “Norte Chico” (27°S to 32°S) is composed of two important sedimentary marine deposits, the Bahía Inglesa and Coquimbo formations. Diverse vertebrate taxa including fish, birds, mammals and abundant chondrichthyans have been described from Bahía Inglesa Formation. However, the vertebrate fauna from Coquimbo Formation has been poorly documented. Based upon field trips and the analysis of collections from the Coquimbo Formation, the elasmobranch fossil fauna is composed of at least nine taxa, two of which are extinct (*Carcharocles megalodon* and *Carcharodon plicatilis*). The rest of the taxa are related with living elasmobranch species that are inhabitants of the Eastern Pacific Ocean and Tropical America coast.

Key words: Neogene, shark, elasmobranch, chondrichthyans, Coquimbo Formation, Pacific.

RESUMO – Sedimentos marinhos de formações geológicas chilenas do Neógeno contêm uma diversa fauna fóssil marinha. No Chile, o “Norte Chico” (27°S to 32°S) é composto de dois importantes depósitos sedimentológicos marinhos, as formações Bahía Inglesa e Coquimbo. Uma diversidade de táxons de vertebrados, que inclui peixes, aves, mamíferos além de abundantes chondrichthies, têm sido descritos para a Formação Bahía Inglesa. Entretanto, a fauna de vertebrados da Formação Coquimbo está pobremente documentada. Com base em trabalhos de campo, bem como na análise de coleções da Formação Coquimbo, observa-se uma fauna de elasmobrânquios composta de pelo menos nove táxons, dois dos quais estão extintos (*Carcharocles megalodon* e *Carcharodon plicatilis*). Os demais táxons estão relacionados com as espécies de elasmobrânquios recentes que habitam as costas leste do Oceano Pacífico Leste e da América Tropical.

Key words: Neógeno, tubarão, elasmobrânquio, condricthios, Formação Coquimbo, Pacífico.

INTRODUCTION

Evidence of fossil elasmobranchs has been collected from Neogene marine sediments distributed all over the world (Cappetta, 2012). As an example, fossil elasmobranch teeth have been collected and studied from four main Chilean geological formations: Bahía Inglesa (27°S), Coquimbo (between 29°S to 30°S), Horcón (32°S) and Navidad (33°S) (Philippi, 1887; Gigoux, 1944; Long, 1993; Arratia & Cione, 1996; Suárez & Encinas, 2002; Suárez & Marquardt, 2003; Suárez *et al.*, 2004; Carrillo-Briceño *et al.*, 2013). These marine fossil deposits have been commonly found in the northern coast of Chile, and characterized by a high proportion of fossil elasmobranch teeth (*e.g.* Suárez & Marquardt, 2003). This particular geographic area is called the “Norte Chico”, and is composed of two geological formations: Bahía Inglesa and Coquimbo. The Bahía Inglesa Formation was deposited under marine conditions, and has been well studied geologically (Rojo, 1985; Marquardt, 1999). Different taxa of fish, birds and mammals have been referred from Bahía Inglesa (Long, 1993; Arratia & Cione, 1996; Walsh & Hume, 2001; Walsh & Naish, 2002; Suárez *et al.*, 2004; Acosta *et al.*, 2006; Chávez, 2008), and at least 23 taxa of chondrichthyan species have been described as well (Long, 1993; Arratia & Cione, 1996; Suárez & Marquardt, 2003; Suárez *et al.*, 2004). Unfortunately, craftsmen and private collectors recurrently visit the marine sediments in Bahía Inglesa to remove and sometimes sell the fossils.

In comparison with the knowledge of the marine vertebrate fauna from Bahía Inglesa, little is known about the vertebrate marine fossil fauna from the Coquimbo Formation (Philippi, 1887; Long, 1993). The area of Coquimbo has experienced urban and industrial expansion during the last two decades. Consequently, most of the marine fossil deposits have been disturbed, and many of them are now destroyed or buried. In this context, the aim of this study was to characterize the elasmobranch fossil marine deposits from the Coquimbo Formation by describing fossil teeth found under the surface ground layer before the complete destruction of the locality.

MATERIAL AND METHODS

The material examined consists of isolated elasmobranch fossil teeth that have been collected in different locations from the Coquimbo Formation: Caleta Chañaral, Punta de Choros, Caleta Hornos, La Cantera Baja, La Herradura, Quebrada Las Rosas and El Rincón (Figure 1). The localities La Cantera Baja, La Herradura and Quebrada Las Rosas were already impacted by urbanization. The sampling design consisted of alternating between surface surveys and bulk samples with excavations of up to 1 m depth. Fossil shark teeth are commonly found in the surface layer in the Coquimbo Formation.

This material is deposited at the Museo Arqueológico de La Serena (MALS), La Serena, Chile. The classifications follow Compagno (1973, 1977) and the terminology is based on Cappetta (2012). Measurements taken include height and width, and these refer to the entire tooth including



Figure 1. A, map showing the regional representation of Chile. The Coquimbo region is shown in gray color. B, zoom in of the Coquimbo region; the gray square represents the study area. The light gray lines are indicating the fourteen areas distributed in this region. C, map showing the locations where the elasmobranch fossils were found. The black dot represents the locations: 1, Caleta Chañaral; 2, Punta de Choros; 3, Caleta Hornos; 4, La Cantera Baja; 5, Quebrada El Culebrón; 6, Quebrada Las Rosas; 7, La Herradura; 8, El Rincón.

the root. Taxonomic identification included an extensive bibliographical review and comparative studies of fossil and extant specimens from: Museo Nacional de Historia Natural de Santiago (SGO-PV) in Chile, Museo Paleontológico de Caldera (MPC), Atacama, both in Chile and Natural History Museum of Basel (NMB), Switzerland; Paleontological collections of the Alcaldía Bolivariana de Urumaco (AMUCURS), Venezuela; Palaeontological Institute and Museum at the University of Zurich (PIMUZ), Switzerland.

GEOLOGICAL SETTINGS

Flat terraces of sedimentation characterize the Coquimbo Formation. These are composed of neritic or sublittoral marine sediments that are fine to medium-grained (Le Roux *et al.*, 2004). Small-cemented yellowish sand is also present, along

with blocks displaced from the bedrock. Shell middens are commonly found in banks between 0.40 and 2 m thickness, on the surface or in sectors with a whitish calcareous crust on the ground (Paskoff, 1970; Le Roux *et al.*, 2004; Acosta *et al.*, 2006). Although the shell middens are mostly from the Plio-Pleistocene, the Miocene transgression seems to be mostly responsible for most of the marine sediment (Le Roux *et al.*, 2006). In this study, shark fossil teeth were obtained from areas near beaches such as marine terraces close to the sea or streams. The terraces have a relative distance of up to 2 km from the coastline and a peak height of 81 meters (Le Roux *et al.*, 2006). Paskoff (1970) suggests a height of 80 m for the outcrop of the Miocene transgression, but some shark fossil teeth can be found in secondary deposits due to relief changes associated with neotectonics. These changes were very important in structuring the relief of the coast during the Plio-Quaternary (Radtke, 1989).

The age of the Coquimbo Formation could be extended backward in time to about 15 Ma (middle Miocene). Fossil invertebrates suggest that all locations sampled in this study originated from Pliocene sedimentary depositions of marine fauna (Herm, 1969). A study based on the potassium-argon radioisotope suggests that the Coquimbo Formation is upper Miocene in age, of around 6 million years ago (Chávez *et al.*, 2007). In addition, thirteen stratigraphic units were identified in the Coquimbo Formation using faunal remains and lithological features (Le Roux *et al.*, 2004). A maximum age of 2 Ma was estimated using strontium from microfossil remains found in the top of the stratigraphic unit (middle Miocene to late Pliocene; Le Roux *et al.*, 2004). The second stratigraphic unit is from Langhian sediment and it has associated with the base of the Coquimbo Formation (Le Roux *et al.*, 2004; Acosta *et al.*, 2006). As a result of these estimations, we propose an age close to the middle Miocene-late Pliocene for the Coquimbo Formation.

The stratigraphic sequence from Quebrada Las Rosas is composed of marine invertebrates and vertebrates (see Figure 2). The basal stratum comprised lime and clay with fossil mollusks and shark teeth. The median stratum is sandy and made of conglomerates of mollusks. The upper stratum is formed of fine sand with a layer of invertebrates. However, this is a general description and we are not able to provide further details because of the lack of stratigraphic studies. A similar formation is observed in Bahía Inglesa Formation (BIF) with common records of shark teeth composed of a large spectrum of taxa (Suárez & Marquardt, 2003; Rivadeneira & Varas, 2012).

SYSTEMATIC PALEONTOLOGY

Order LAMNIFORMES Berg 1937

Family †OTODONTIDAE Glikman, 1964

†*Carcharocles* Jordan & Hannibal, 1923

†*Carcharocles megalodon* (Agassiz, 1843)
(Figures 3A-C)

Material. Four lower lateral teeth (MNHN SGO. 58, Coquimbo; MALS P-228, Punta de Choros; MALS P-6564, Quebrada Chañaral de Aceitunas; MALS P-227, Quebrada El Culebrón).

Description. Teeth have a triangular, broad and slightly symmetrical crown. The lingual face of the crown is flat and the labial is convex with a typical large neck, clearly observed in the specimen MNHN SGO. 58. The crown in the rest of the specimens is poorly preserved. Both cutting edges have a fine serration. The root has well developed lobes and a weak lingual protuberance. The teeth range in height between 32 and 98 mm and width between 35 to 93 mm. The largest tooth is the specimen MALS P-228, with 98 mm in height and 93 mm in width.

Remarks. *Carcharocles* was a cosmopolitan species and its fossil record has been reported from almost all Neogene sedimentary deposits that contain shark teeth (García *et al.*, 2009). In the Neogene deposits, *Carcharocles* is mainly represented by *C. chubutensis* Ameghino, 1901 and *C. megalodon* Agassiz, 1843 (e.g. Pimiento *et al.*, 2010, 2013a,b; Cappetta, 2012; Pimiento & Clements, 2014). *C. megalodon* has a stratigraphic range from the middle Miocene to the late Pliocene (Pimiento & Clements, 2014), whereas *C. chubutensis* has been restricted to the early/middle Miocene (Pimiento & Clements, 2014). However, taxonomical distinctions and generic identification between the taxa have been debated over the years. The classification of *Carcharocles* lineage is still discussed (Pimiento *et al.*, 2010, 2013b; Reinecke *et al.*, 2011; Cappetta, 2012; Bor *et al.*, 2012). From the Neogene of Chile, *C. megalodon* has been reported from the Bahía Inglesa and Lo Abarca formations (Long, 1993; Walsh, 2001; Encinas, 2002; Suárez & Encinas, 2002; Suárez *et al.*, 2004). Philippi (1887) reported an isolated tooth of *C. megalodon* with unclear origin from Coquimbo region. The specimens of *C. megalodon* from the Coquimbo Formation extend the southern distribution on the Eastern Pacific during the late Neogene. An overview of the *C. megalodon* is shown by Carrillo-Briceño *et al.* (2013) from America fossil records.

Family LAMNIDAE Müller & Henle, 1838

Carcharodon Smith in Müller & Henle, 1838

Carcharodon carcharias Linnaeus, 1758
(Figure 4A)

Material. Fifty upper lateral teeth (MALS P-59, MALS P-60, MALS P-64, Quebrada El Culebrón; MALS P-79, MALS P-80, MALS P-83, MALS P-84, MALS P-86, MALS P-87, MALS P-88, MALS P-90, MALS P-92, MALS P-105, La Cantera Baja; MALS P-119, MALS P-126, MALS P-127, Quebrada Las Rosas; MALS P-131, MALS P-133, Caleta Chañaral; MALS P-142, MALS P-143, MALS P-144, MALS P-150, MALS P-151, MALS P-152, MALS P-159, MALS P-160, MALS P-161, MALS P-163, MALS P-164, MALS P-165, MALS P-166, MALS P-169, MALS P-170,




Figure 2. A, geological stratigraphic units showing the association of elasmobranchs found in the Coquimbo Formation. B, *Carcharodon carcharias* fossil tooth found on silt-clay mixed with stones and sedimentary soil. C, landscape of sedimentary terraces of the Coquimbo Formation located in Quebrada Las Rosas. D, general view of stratigraphic units from Quebrada Las Rosas.




Figure 3. *Carcharocles megalodon* fossil teeth. A, MNHN SGO.58, lower lateral tooth in labial view; B, MALS P-6564, lower lateral tooth in labial view; C, MALS P-228, lower lateral in labial view. Scale bars = 10 mm.




Figure 4. Elasmobranch fossil teeth. **A**, *Carcharodon carcharias*, MALS P-87, upper lateral tooth in labial view. **B**, *Carcharodon plicatilis*, MALS P-63, upper lateral tooth in labial view. **C**, *Isurus cf. I. oxyrinchus*, MALS P-199, upper anterior tooth in labial view. **D**, *Odontaspis ferox*, MALS P-104, lateral tooth in labial view. **E**, *Carcharhinus* sp. MALS P-111, lower lateral tooth in labial view. **F**, *Pristiophorus* sp., MALS P-75, rostral tooth. **G**, *Hexanchus* cf. *H. griseus*, MALS P-108, upper anterior tooth in lingual view. **H**, *Hexanchus* cf. *H. griseus*, MALS P-225, lower lateral tooth in lingual view. **I**, *Myliobatis* sp., MALS P-231, symphyseal tooth of in lingual view. Scale bars = 10 mm.

MALS P-176, MALS P-177, MALS P-178, MALS P-179, MALS P-184, MALS P-189, MALS P-190, MALS P-193, MALS P-197, MALS P-198, La Herradura; MALS P-203, MALS P-206, MALS P-208; MALS P-212, MALS P-216, MALS P-217, MALS P-219, Quebrada El Culebrón), one upper posterior tooth (MALS P-114, La Cantera Baja), three upper teeth (MALS P-147, MALS P-181, MALS P-185, La Herradura), ten upper anterior teeth (MALS P-85, MALS P-93, La Cantera Baja; MALS P-116, Quebrada Las Rosas; MALS P-146, MALS P-155, MALS P-186, MALS P-191,

MALS P-194, La Herradura; MALS P-210, MALS P-220, Quebrada El Culebrón), three lower posterior teeth (MALS P-78, MALS P-113, La Cantera Baja; MALS P-148, La Herradura), fourteen lower lateral teeth (MALS P-81, MALS P-82, MALS P-91, La Cantera Baja; MALS P-129, MALS P-130, Quebrada Las Rosas; MALS P-145, MALS P-149, MALS P-171, MALS P-196, La Herradura; MALS P-207, MALS P-209, MALS P-211, MALS P-2014, MALS P-215, Quebrada El Culebrón), six lateral teeth (MALS P-117, MALS P-128, Quebrada Las Rosas; MALS P-153, MALS

P-162, MALS P-173, La Herradura; MALS P-223, Quebrada El Culebrón), twenty lower anterior teeth (MALS P-89, La Cantera Baja; MALS P-141, MALS P-154, MALS P-156, MALS P-168, MALS P-172, MALS P-174, MALS P-180, MALS P-182, MALS P-183, MALS P-192, MALS P-195, La Herradura; MALS P-204, MALS P-205, MALS P-213, MALS P-218, MALS P-221, MALS P-222, MALS P-224, Quebrada El Culebrón) and one unknown position tooth (MALS P-157, La Herradura).

Description. Upper and lower teeth have a broad, triangular and slightly elongated labio-lingually compressed crown. The labial face is flattened whereas the lingual is convex; both cutting edges are coarse serrated. The basal portion of the crown is in contact with roots and lacks enamel, forming a thin dental band, observable on the labial and lingual surface. Root with two well-defined lobes that are straight or it may present a U-shaped arch in some specimens. The teeth range in height between 10 and 73 mm and width between 10 to 47 mm.

Remarks. The fossil record of *Carcharodon carcharias* is known from the lower Pliocene to recent, and it has been referred to Africa, Asia, Australia, Europa and North and South America (Kemp, 1991; Cappetta, 2012; Ehret *et al.*, 2012; Cione *et al.*, 2012; Carrillo-Briceño *et al.*, 2013). The fossil record of *C. carcharias* from the late Neogene of Chile is abundant, mainly from the Pliocene strata of the Bahía Inglesa Formation (Long, 1993; Walsh, 2001; Quilodran & Marquardt, 2001; Walsh & Naish, 2002; Suárez & Marquardt, 2003). However, some specimens have been dubiously referred to the late Miocene strata of this unit (*e.g.* Bianucci *et al.*, 2006). Other fossil record of *C. carcharias* from Chile include the following localities: Bahía Salado, Bahía Tongoy, Cuenca del Tiburon, Horcón, La Cueva, and La Portada (Suárez & Brito, 2000; Emslie & Correa, 2003; Suárez & Marquardt, 2003; Le Roux *et al.*, 2006; Carrillo-Briceño *et al.*, 2013). Cione *et al.* (2012) and Carrillo-Briceño *et al.* (2013) provided a summary of the fossil record of *C. carcharias* in South America.

†*Carcharodon plicatilis* Agassiz, 1843
(Figure 4B)

Material. Upper lateral tooth (MALS P-63, El Rincón).

Description. The tooth has a wide, triangular and labio-lingually compressed crown. The cutting edges are sharp without any serration. The root is compressed with two lobes, and the base shows a concavity forming an inverted V. The tooth measures 50 mm in height and 36 mm in width.

Remarks. Several morphological aspects of the taxonomic classification of this species have been debated over the years, without consensus (*e.g.* Purdy *et al.*, 2001; Cappetta, 2012; Cione *et al.*, 2012; Ehret *et al.*, 2012). Previous taxonomic assignments have included some species of “wide-toothed shape” such as *Isurus planus* Agassiz, 1856, *Isurus hastalis* Agassiz, 1838 and *Isurus xiphodon* Agassiz, 1838 to *Cosmopolitodus* (Cappetta, 2006; Cione *et al.*, 2012). Ehret *et al.* (2012) included *Cosmopolitodus* to

Carcharodon, proposing a transition between *Carcharodon hastalis* and *Carcharodon carcharias*. Ward & Bonavia (2001) suggested the recognition of only one species, and proposed that *C. xiphodon* is a *nomen dubium* because of the uncertainty implied in differentiating *Cosmopolitodus hastalis* from *Cosmopolitodus xiphodon*. According to Cione (1988), Purdy *et al.* (2001) and Cione *et al.* (2012), there is a species that appears to be the putative sister group of *Carcharodon carcharias* and different from *Cosmopolitodus hastalis*. Cione *et al.* (2012) proposed this species as “*Isurus*” *plicatilis* Agassiz, 1843, which is recognized as an available and valid taxon. According to Cione *et al.* (2012), only *plicatilis* or *xiphodon* [adapting latter from Purdy *et al.* (2001)] should be referred to the genus *Carcharodon*. Hereby, we adopt the taxonomic proposal suggested by Cione (1988) and Cione *et al.* (2012). In the Neogene of Chile, *Carcharodon plicatilis* has been reported from the Bahía Inglesa, Navidad and Coquimbo formations (Long, 1993; Quilodran & Marquardt, 2001; Walsh, 2001; Suárez & Encinas, 2002; Walsh & Naish, 2002; Suárez & Marquardt, 2003; Le Roux *et al.*, 2006).

Isurus Rafinesque, 1810

Isurus cf. *oxyrinchus* Rafinesque, 1810
(Figure 4C)

Material. Upper lateral tooth (MALS P-58, Caleta Chañaral), upper anterior tooth (MALS P-199, Caleta de Hornos), lower anterior tooth (MALS P-134, Caleta Chañaral) and anterior tooth, unknown jaw position (MALS P-188, La Herradura).

Description. Three teeth are incomplete with damaged roots (MALS P-134, MALS P-188, MALS P-199). In anterior teeth the crown is elongated, asymmetrical, distally inclined and with smooth sharp cutting edges; its apex is sharp-pointed. The lateral has a triangular crown slightly inclined distally. There is an evident lingual protuberance with a transverse groove. The only complete tooth (MALS P-58) measures 25 mm in height and 12 mm in width.

Remarks. *Isurus oxyrinchus* has a stratigraphic record from the late Oligocene to present with a cosmopolitan paleodistribution (Reinecke, 2011). Despite the fact that the specimens from the Coquimbo Formation are mostly incomplete, they exhibit a close resemblance to teeth of the extant *I. oxyrinchus*, allowing the association of the fossils with this taxon. In Chile, the fossil record of *I. oxyrinchus* appears regularly along the central coast of the Bahía Inglesa Formation (Suárez *et al.*, 2002; Walsh, 2001; Long, 1993; Suárez & Marquardt, 2003; Suárez *et al.*, 2004). In contrast to Bahía Inglesa, *I. oxyrinchus* fossil teeth are less frequent southward. However, this species has been found in marine sediments from Lo Abarca, Navidad and Horcón formations (Encinas, 2002; Suárez & Encinas, 2002; Suárez & Marquardt, 2003; Suárez *et al.*, 2006; Carrillo-Briceño *et al.*, 2013).

Family ODONTASPIDIDAE Müller & Henle, 1839

Odontaspis Agassiz, 1838

Odontaspis ferox Risso, 1810
(Figure 4D)

Material. Lateral tooth (MALS P-104, La Cantera Baja).

Description. The crown is elongated and slender with sharp and smooth cutting edges. Fossil and extant specimens of *Odontaspis ferox* show between three and one lateral cusplets (e.g. Purdy *et al.*, 2001). However, the specimen MALS P-104 preserves only one cusplet as a consequence of the damaged root. The root shows an evident lingual protuberance. The incomplete tooth is 22 mm in height.

Remarks. *Odontaspis ferox* has a fossil record from the Miocene to recent (Purdy *et al.*, 2001; Aguilera & Rodríguez de Aguilera, 2001; Marsili, 2007). During the Paleocene-Eocene, this species is considered as the dominant group of Patagonia of Chile and Argentina, as well as of other ancient temperate seas around the world (e.g. Eocene of Seymour Island, Antarctica) (Arratia & Cione, 1996). This species has been reported to the Bahía Inglesa (Suárez *et al.*, 2002) and Navidad formations (Suárez & Encinas, 2002; Suárez & Marquardt, 2003).

Order CARCHARHINIFORMES Compagno, 1973

Family CARCHARHINIDAE Jordan & Evermann, 1896

Carcharhinus Blainville, 1816

Carcharhinus sp.
(Figure 4E)

Material. Three lower lateral teeth (MALS P-110, MALS P-112, La Cantera Baja; MALS P-111, La Cantera).

Description. The fossil teeth are fragmented, preserving only the crown or the root. The crown is triangular and decreases in size toward the apex of the tooth. The crown is distally inclined towards the right with fine serration along the cutting edges (MALS P-110). The root has broad lobes with slightly basal concavity and a shallow nutritive groove (MALS P-111). Diagnostic characters of the species are difficult to identify because of the fragmented condition and poor preservation of the specimens.

Remarks. Morphological features of the tooth within the genus *Carcharhinus* closely resembles between species. This confusion makes difficult the taxonomic classification of the carcharhinids fossil teeth, especially when they are incomplete and eroded. At least two species have been identified from Chilean marine sediments i.e. *Carcharhinus albimarginatus* (Rüppell, 1837) (Long, 1993; Suárez & Marquardt, 2003) and *Carcharhinus brachyurus* (Günther, 1870) (Suárez *et al.*, 2002; Suárez *et al.*, 2004; Carrillo-Briceño *et al.*, 2013). However, previous studies in modern species have confirmed the presence of other carcharhinids species in Chilean coast. Pequeño & Saez (2003) have reported the presence of *Carcharhinus galapagensis* (Snodgrass & Heller, 1905) from Salas y Gomez

Island and Hernández *et al.* (2008) identified the species *Carcharhinus obscurus* (Lesueur, 1818) from shark fins traded in Chilean market. According to these antecedents, the presence of extinct and modern carcharhinids should be better studied in Chile because there are uncertainly in some of the species used to occur and/or still found in Chilean coast. In the Neogene and Quaternary of Chile, species of genus *Carcharhinus* has been reported from the Miocene and Pliocene in the Bahía Inglesa Formation (Long, 1993; Walsh, 2001; Suárez & Marquardt, 2003; Suárez *et al.*, 2002), Bahía Salado (Suárez & Brito, 2000), Cuenca del tiburón (Emslie & Correa, 2003; Suárez & Marquardt, 2003), La Portada (Suárez & Marquardt, 2003) and Horcón (Carrillo-Briceño *et al.*, 2013).

Order PRISTIOPHORIFORMES Berg, 1958

Family PRISTIOPHORIDAE Bleeker, 1859

Pristiophorus Müller & Henle, 1837

Pristiophorus sp.
(Figure 4F)

Material. Thirteen rostral teeth (MALS P-69, MALS P-70, MALS P-71, MALS P-72, MALS P-73, MALS P-74, MALS P-75, MALS P-76, MALS P-77, La Cantera Baja; MALS P-118, Quebrada Las Rosas; MALS P-138, MALS P-139, MALS P-140, La Herradura).

Description. The rostral teeth are elongated and slender, with a dorso-ventrally compressed crown. The cutting edges are smooth and without serration. The root is robust with a flat base and is wider than the crown. The root also has a conical shape with the widest part being the base. Most of the specimens are complete and measure between 31 and 32 mm in length.

Remarks. *Pristiophorus* is known from the upper Cretaceous to present (Cappetta, 2012). The Neogene fossil record of *Pristiophorus* showed a wide paleobiogeographic distribution in America (Carrillo-Briceño *et al.*, 2015). Nowadays, there is only one species in America *Pristiophorus schroederi* Müller and Henle, 1837, which is distributed in the Bahamas region, Western Central Atlantic (Compagno *et al.*, 2005). The taxonomy of the fossil species of *Pristiophorus* found in the Americas, and especially the rostral teeth have been poorly studied. For this reason we prefer to keep the *Pristiophorus* specimen from the Coquimbo Formation in open nomenclature. The fossil record of *Pristiophorus* in Chile includes Bahía Inglesa (Walsh, 2001; Suárez *et al.*, 2002), Navidad (Suárez & Encinas, 2002; Suárez *et al.*, 2006), Horcón (Carrillo-Briceño *et al.*, 2013) and La Cueva formations (Suárez & Marquardt, 2003).

Order HEXANCHIFORMES Buen, 1926

Family HEXANCHIDAE Gray, 1851

Hexanchus Rafinesque, 1810

Hexanchus cf. *griseus*
(Figures 4G-H)

Material. Five upper teeth (MALS P-107, MALS P-108, La Cantera Baja; MALS P-125, Quebrada Las Rosas; MALS P-61, MALS P-226, Quebrada el Culebrón) and a lower lateral tooth (MALS P-109, La Cantera Baja; MALS P-225, Quebrada el Culebrón).

Description. Upper teeth are elongated acrocones that are distally inclined, with two or five distally directed cusplets. The root is flattened and subquadrangular. The lower tooth is incomplete with a missing acrocone (MALS P-225). However, five small and distally directed cusplets are present. The complete teeth range between 16 and 21 mm in height and between 8 to 18 mm in width.

Remarks. *Hexanchus griseus* is known from the Miocene to the recent (Cione & Reguero, 1994). In Chile, *Hexanchus griseus* and *Hexanchus* sp. have been referred to the Bahía Inglesa and Horcón formations (Long, 1993; Walsh, 2001; Suárez & Marquardt, 2003; Carrillo-Briceño *et al.*, 2013). Suárez & Marquardt (2003) referred isolated teeth of *Hexanchus* associated with cetacean remains from phosphatic layer in Caleta Inglesa and Bahía Chañaral. Arratia & Cione (1996) and Carrillo-Briceño *et al.* (2013) presented a brief summary of the fossil record of *Hexanchus* in South America.

Order MYLIOBATIFORMES Compagno 1973
Family MYLIOBATIDAE Bonaparte 1838

Myliobatis Cuvier, 1816

Myliobatis sp.
(Figure 4I)

Material. Six isolated teeth (MALS P-66, MALS P-230, MALS P-231, MALS P-232, MALS P-233, MALS P-234, La Cantera Baja).

Description. The specimens consist of complete isolated teeth of the median files. These teeth are broader than long, with a slightly curved hexagonal contour. In all teeth the crown is practically flat with a smooth occlusal surface. Labial and lingual faces are slightly ornamented. The root displays vascularization type polyaulacorhize. The teeth are between 30 and 50 mm wide.

Remarks. The fossil record of *Myliobatis* extends worldwide from the Paleocene to present (Cappetta, 2012). The taxonomic identification of the isolated teeth of *Myliobatis* is extremely difficult because of the high dental variation within this group (Welton & Zinsmeister, 1980; Nishida, 1990). For this reason, a specific taxonomic assignation was not possible and we assigned all specimens from the Coquimbo Formation to *Myliobatis* sp. In the Neogene of Chile, the genus *Myliobatis* has been found in the Bahía Inglesa, La Cueva and Navidad formations (Walsh, 2001; Suárez & Marquardt, 2003; Suárez *et al.*, 2004; Suárez *et al.*, 2002). Summary of the fossil record of *Myliobatis* is discussed in Carrillo-Briceño *et al.* (2013).

DISCUSSION

Our results revealed at least nine elasmobranch taxa present during the Neogene at the Coquimbo Formation. These include *Carcharocles megalodon*, *Carcharodon carcharias*, *C. plicatilis*, *Isurus* cf. *I. oxyrinchus*, *Pristiophorus* sp., *Odontaspis ferox*, *Carcharhinus* sp., *Hexanchus* cf. *H. griseus* and *Myliobatis* sp. Of these species, *C. megalodon*, *C. plicatilis*, *C. carcharias* and *I. oxyrinchus* have been previously reported from the Coquimbo Formation (*e.g.* Philippi, 1887; Long, 1993; Suárez & Marquardt, 2003; Le Roux *et al.*, 2006). According to our results, *Odontaspis ferox* and *Pristiophorus* sp. represent the first fossil records for the Coquimbo Formation, although Long (1993) included two further species, *Cetorhinus maximus* and *Aetobatus* sp. for this locality. Our assemblage includes at least two extinct species: *C. megalodon* and *C. plicatilis*. With the exception of *Pristiophorus* sp., the rest of the elasmobranch taxa found in this study are related to the living species distributed along the Pacific coast of South America (*e.g.* Egaña & McCosker, 1984; Acuña *et al.*, 2002; Hernández & Lamilla, 2004; Hernández *et al.*, 2008).

The Neogene is characterized as a period of intense climatic, tectonic and biotic change (Zachos, 2001; Dekens *et al.*, 2007; Dowsett & Robinson, 2009; Garreaud *et al.*, 2010). For instance, events such as the hyper-aridity of the Atacama Desert and the reactivation of the modern flow of the Humboldt Current produced important changes in the Neogene climate conditions (Garreaud *et al.*, 2010; Tsuchi, 2002). Studies of elasmobranch fossils suggest vast variations in species richness in the South Pacific (Villafaña & Rivadeneira, 2014). The elasmobranch fossil record of the Coquimbo Formation would suggest differences in the dynamics of elasmobranch biodiversity. Species, such as *Carcharocles megalodon*, *Carcharodon plicatilis* and *Pristiophorus* sp. probably became extinct due to climatic and biotic changes that occurred during the late Neogene. Rivadeneira & Marquet (2007) suggested that the diversity in the composition of mollusk species decreased during the Miocene and Pliocene along the South American Pacific coast. A similar tendency in decreasing diversity was shown with the extinction of marine mammals (Valenzuela-Toro *et al.*, 2013), and sea birds (Chávez *et al.*, 2007). The extinction of several macro-vertebrates may reflect the macro-evolutionary dynamics of those taxa in the temperate Pacific coast of South America.

In addition, we found differences in the distribution of extinct elasmobranch species in comparison to modern species. For example, the fossil distribution of *Carcharodon carcharias* and *Hexanchus* increased southward (Villafaña & Rivadeneira 2013). In contrast, the presence of shark species such as *Isurus oxyrinchus*, *Odontaspis ferox*, *Carcharhinus* sp. and *Myliobatis* sp. decreased southward (Villafaña & Rivadeneira 2013). Analyzing the distribution of genus *Carcharias* in the South Pacific Ocean, it might be attributed to the abrupt temperature decrease during the middle Pliocene and Pleistocene (Cione *et al.*, 2007). This change in the temperature eventually also affected the

distribution other species. However, Villafañá & Rivadeneira (2013) suggested that the fluctuation in elasmobranch biodiversity might also be explained by ecological and life history aspects such as body size, modes of reproduction, and migration patterns.

In order to explain the variability in richness and composition of marine fossil records in Chile, it would be valuable to increase the sampling effort of fossil records from the Coquimbo Formation and other Neogene units. Furthermore, more sampling sites should be determined and reviewed. The well-preserved sedimentary sequence of the Coquimbo Formation is an unexplored field. A better description of the elasmobranch fossil records will set the baseline to gain a better understanding of the ecology, distribution and evolution of taxa that ranged the oceans millions of years ago.

ACKNOWLEDGMENTS

We would like to thank several people from the MALS for their friendly help and collaboration in this research. The manuscript was greatly improved by the very constructive comments and suggestions made by A.L. Cione and G. Cuny. Thank to V. Castelletto for taking photographs for this study, and also thank to M. Heidemeyer and J. Neenam for the proof editing of the manuscript.

REFERENCES

- Acosta, C.H.; Canto, J. & Tambussi, C. 2006. Pingüinos (Aves, Spheniscidae) en Coquimbo (Mioceno medio-Plioceno tardío), Chile y su vinculación con las corrientes oceánicas. *Revista Española de Paleontología*, **21**:115-121.
- Acuña, E.; Villaroel, J.C. & Grau, R. 2002. Fauna íctica asociada a la pesquería de pez espada (*Xiphias gladius* Linnaeus). *Gayana*, **66**:263-267. doi: 10.4067/S0717-65382002000200025
- Agassiz, L. 1833-1843. *Recherches sur les poissons fossiles*. Volume 3, Neuchâtel, p. 390.
- Agassiz, L. 1856. Notice on the fossil fishes found in California by Blake, W.P. *American Journal of Science and Arts*, **21**:272-275.
- Aguilera, O. & Rodríguez de Aguilera, D. 2001. An exceptional coastal upwelling fish assemblage in the Caribbean Neogene. *Journal of Paleontology*, **75**:732-742. doi: 10.1666/0022-3360(2001)075<0732:AECUEA>2.0.CO;2
- Alván, A.; Apolín, J. & Chacaltana, C. 2004. Los dientes de Seláceos (Chondrichthyes) y su aplicación estratigráfica en Las Lomas de Ullujaya (Ica, Perú). In: CONGRESO PERUANO DE GEOLOGÍA, 13, 2004. *Resúmenes Extendidos*, Lima, p. 595-598.
- Ameghino, F. 1901. L'âge des formations sédimentaires de Patagonie. *Anales de la Sociedad Científica Argentina*, **51**:20-39, 65-91.
- Arratia, G. & Cione, A.L. 1996. The fish fossil record of southern South America. In: G. Arratia (ed.) *The vertebrate fossil record of southern South America*, Münchener Geowissenschaft Abhandlungen, p. 565-593.
- Balbotín, F. & Reyes, E. 1981. Ataque de tiburón registrado en la costa de Chile Central. *Noticiero Mensual del Museo Historia Natural*, **298**:3-7.
- Beal, C.H. 1948. Reconnaissance of the geology and oil possibilities of Baja California, Mexico. *Geological Society of America Memoir*, **31**:1-150. doi: 10.1130/mem31
- Berg, L.S. 1937. A classification of fish-like vertebrates. *Bulletin of the Academy of Sciences of the USSR*, **1937**:1277-1280.
- Berg, L.S. 1958. *System der rezenten und fossilen Fischartigen und Fische*. Berlin, Verlag Wissenschaften, 310 p.
- Bianucci, G.; Sorbi, S.; Suárez, M. & Landini, W. 2006. The southernmost sirenian record in the Eastern Pacific Ocean, from the late Miocene of Chile. *Comptes Rendus Palevol*, **5**:945-952. doi: 10.1016/j.crpv.2006.06.001
- Blainville, H.M.D. 1816. Prodrome d'une nouvelle distribution systématique du règne animal. *Bulletin de la Société Philomathique de Paris*, **8**:105-112, 121-124.
- Bleeker, P.R. 1859. Enumeratio specierum piscium hucusque in Archipelago indico observatarum. *Acta Societatis Scientiarum Indo-Neerlandae*, **6**:1-276.
- Bonaparte, C.L. 1838. Selachorum tabula analytica. *Nuovi Annali della Scienze Naturali Bologna*, **1**:195-214.
- Bor, T.; Reinecke, T. & Verschuere, S. 2012. Miocene Chondrichthyes from Winterswijk-Miste, the Netherlands. *Palaeontos*, **21**:1-136.
- Boreske, J.R.; Goldberg, L. & Cameron, B. 1972. A reworked cetacean with clam borings: Miocene of North Carolina. *Journal of Paleontology*, **46**:130-139.
- Campbell, H.J.; Andrews, P.B.; Beu, A.G.; Maxwell, P.A.; Edwards, A.R.; Laird, M.G.; Hornibrook, N.B.; Mildenhall, D.C.; Watters, W.A.; Buckeridge, J.S.; Lee, D.E.; Strong, C.P.; Wilson, G.J. & Hayward, B.W. 1993. Cretaceous-Cenozoic geology and biostratigraphy of the Chatham Islands, New Zealand. *Institute of Geological and Nuclear Sciences Monograph*, **2**:1-269.
- Cappetta, H. 1970. *Les Sélaciens du Miocène de la région de Montpellier*. Montpellier, Laboratoire de Paléontologie, 139 p.
- Cappetta, H. 1980. Les selaciens du Cretacé supérieur du Liban. II: Batoides. *Palaeontographica A*, **168**:149-229.
- Cappetta, H. 1987. *Chondrichthyes II-Mesozoic and Cenozoic Elasmobranchii. Handbook of Paleichthyology*. New York, Verlag, 193 p.
- Cappetta, H. 2006. Elasmobranchii post-Triadici (index generum et specierum). In: W. Riegraf (ed.) *Fossilium Catalogus I: Animalia*, Backhuys Publish, p. 472.
- Cappetta, H. 2012. Chondrichthyes II: Mesozoic and Cenozoic Elasmobranchii: Teeth. In: H.P. Schultze (ed.) *Handbook of Paleichthyology*, Verlag, p. 512.
- Carrillo-Briceño, J.D.; De Gracia, C.; Pimiento, C.; Aguilera, O.A.; Kindlimann, R.; Santamarina, P. & Jaramillo, C. 2015. A New Late Miocene Chondrichthyan assemblage from the Chagres Formation, Panama. *Journal of South America Earth Science*, **60**:56-70. doi: 10.1016/j.jsames.2015.02.001
- Carrillo-Briceño, J.D.; González-Barba, G.; Landaeta, M.F. & Sven, N. 2013. Condriactios del Plioceno superior de la Formación Horcón, Región de Valparaíso, Chile central. *Revista Chilena de Historia Natural*, **86**:191-206. doi: 10.4067/s0716-078x2013000200008
- Chávez, M.F. 2008. *La ornitofauna de la Formación Bahía Inglesa, Caldera, Chile*. Escuela de Ciencias, Universidad Austral de Chile, Tesis de grado, 162 p.
- Chávez, M.; Stucchi, M. & Urbina, M. 2007. El registro de Pelagornithidae (Aves: Pelecaniformes) y la avifauna Neógena del Pacífico sudeste. *Boletín del Instituto Francés de Estudios Andinos*, **36**:175-197. doi: 10.4000/bifea.3780
- Cione, A.L. 1988. *Los peces de las formaciones marinas del Cenozoico de Patagonia*. Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Thesis Doctoral, 536 p.

- Cione, A.L.; Cabrera, D. & Barla, M.J. 2012. Oldest record of the great white shark (Lamnidae, *Carcharodon*, Miocene) in the Southern Atlantic. *Geobios*, **45**:167-172. doi:10.1016/j.geobios.2011.06.002
- Cione, A.L.; Mennucci, J.A.; Santalucita, F. & Acosta-Hospitaleche, C. 2007. Local extinction of sharks of genus *Carcharias* Rafinesque, 1810 (Elasmobranchii, Odontaspidae) in the Eastern Pacific Ocean. *Revista Geológica de Chile*, **34**:139-145. doi:10.4067/s0716-02082007000100007
- Cione, A.L. & Reguero, M.A. 1994. New records of the sharks *Isurus* and *Hexanchus* from the Eocene of Seymour Island, Antarctica. *Proceedings of the Geologists' Association*, **105**:1-14. doi:10.1016/s0016-7878(08)80134-4
- Compagno, L.J.V. 1973. Interrelationships of living elasmobranchs. In: P.H. Greenwood; C. Patterson & H. Miles (eds.) *Interrelationships of fishes*, Zoological Journal of the Linnean Society, p. 15-61.
- Compagno, L.J.V. 1977. Phyletic relationships of living sharks and rays. *American Zoologist*, **17**:303-322. doi:10.1093/icb/17.2.303
- Compagno, L.J.V. 1984. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. *FAO Fisheries Synopsis*, **125**:1-249.
- Compagno, L.J.V.; Dando, V.M. & Flower, S. 2005. *Sharks of the world*. Princeton, Princeton University Press, 368 p.
- Cope, E.D. 1868. An addition to the vertebrate fauna of the Miocene period, with a synopsis of the extinct Cetacea of the United States. *Proceedings of the Academy of Natural Sciences of Philadelphia*, **19**:138-157.
- Cuvier, G. 1816. *Le Règne Animal Distribué D'après son Organisation Pour Servir de Base à L'histoire Naturelle des Animaux et D'introduction à L'anatomie Comparée. Les Reptiles, les Poissons, les Mollusques et les Annélides*. Deterville. Paris, p. 532.
- De Buen, F. 1926. Catálogo ictológico del Mediterráneo español y de Marruecos. *Resultados Campanas Internacionales Instituto Español Oceanografía*, **2**: 153-161.
- Dekens, P.S.; Ravelo, A.C. & McCarthy, M.D. 2007. Warm upwelling regions in the Pliocene warm period. *Paleoceanography*, **22**:1-12. doi:10.1029/2006PA001394
- Dowsett, H.J. & Robinson, M.M. 2009. Mid-Pliocene equatorial Pacific sea surface temperature reconstruction: a multi-proxy perspective. *Philosophical Transactions of the Royal Society A*, **367**:109-125. doi:10.1098/rsta.2008.0206
- Egaña, A.C. & McCosker, J.E. 1984. Attacks on divers by white sharks in Chile. *California fish and game*, **70**:173-179.
- Ehret, D.J.; Hubbell, H. & MacFadden, B.J. 2009. Exceptional preservation of the white shark *Carcharodon* (Lamniformes, Lamnidae) from the early Pliocene of Peru. *Journal of Vertebrate Paleontology*, **29**:1-13. doi:10.1671/039.029.0113
- Ehret, D.J.; MacFadden, B.J.; Jones, D.S.; Devries, T.J.; Foster, D.A. & Salas-Gismondi, R. 2012. Origin of the white shark *Carcharodon* (Lamniformes: Lamnidae) based on recalibration of the upper Neogene Pisco Formation of Peru. *Paleontology*, **55**:1139-1153. doi:10.1111/j.1475-4983.2012.01201.x
- Emslie, S.D. & Correa, C.G. 2003. A new species of penguin (Spheniscidae: Spheniscus) and other birds from the late Pliocene of Chile. *Proceedings of the Biological Society of Washington*, **116**:308-316.
- Encinas, A. 2002. Presencia de vertebrados marinos neógenos en la Formación Lo Abarca, Chile Central. In: CONGRESO LATINOAMERICANO DE PALEONTOLOGÍA DE VERTEBRADOS, 1, 2002. *Actas*, Santiago, p. 32.
- Fitzgerald, E.M.G. 2004. The fossil record of cetaceans (Mammalia) on the Australian continent. *The Fossil Collector*, **72**:5-32.
- García, E.X.M.; Antunes, M.T.; Balbino, A.C.; Ruiz-Muñoz, F. & Cívís-Llovera, J. 2009. Los tiburones Lamniformes (Chondrichthyes, Galeomorphii) del Plioceno inferior de la Formación Arenas de Huelva, suroeste de la cuenca del Guadalquivir, España. *Revista Mexicana de Ciencias Geológicas*, **26**:674-686.
- Garreaud, R.D.; Molina, A. & Farias, M. 2010. Andean uplift, ocean cooling and Atacama hyperaridity: a climate modeling perspective. *Earth and Planetary Science Letters*, **292**:39-50. doi:10.1016/j.epsl.2010.01.017
- Gigoux, E. 1944. Fósiles de Atacama. *Boletín del Museo Nacional de Historia Natural*, **22**:45-48.
- Glikman, L.S. 1964. *Sharks of the Paleogene and their stratigraphic significance*. Moscow, Nauka Press, 229 p.
- Gray, J. 1851. *List of the specimens of fish in the collection of the British Museum. Part 1. Chondropterygii*. London, British Museum (Natural History), 160 p.
- Günther, A.C.L.G. 1870. *Catalogue of the Fishes in the British Museum*. The Trustees, 369 p.
- Herm, D. 1969. Marines Pliozän und Pleistozän in Nord-und Mittel-Chile unterbesonderer Berücksichtigung der Entwicklung der Mollusken-Faunen. *Zitteliana*, **2**:1-159.
- Hernández, S.; González, M.T.; Villaroel, J.C. & Acuña, E. 2010. Seasonal variation in fish bycatch associated with an artisanal flounder fishery on Coquimbo Bay, Chile. *Revista de Biología Marina y Oceanografía*, **45**:695-703. doi:10.4067/s0718-19572010000400013
- Hernández, S.; Haye, P. & Shivji, M. 2008. Characterization of the pelagic shark fin trade in north central Chile by genetic identification and trader surveys. *Journal of Fish Biology*, **73**:2293-2304. doi:10.1111/j.1095-8649.2008.02016.x
- Hernández, S. & Lamilla, J. 2004. Presencia del gran Tiburón blanco *Carcharodon carcharias* en las costas de Chile. In: CONGRESO DE CIENCIAS DEL MAR, 24, 2004. *Resúmenes*, La Serena.
- Jordan, D.S. & Evermann, B.W. 1896. The fishes of North and middle America: a descriptive catalogue of the species of fish-like vertebrates found in the waters of America, North of the Isthmus of Panama. Part I. *Bulletin of the United States National Museum*, **47**:1-1240.
- Jordan, D.S. & Hannibal, H. 1923. Fossil sharks and rays of the Pacific Slope of North America. *Bulletin of the Southern California Academy of Sciences*, **22**:27-63.
- Kazar, E. 2002. Revised phylogeny of the Physeteridae (Mammalia: Cetacea) in light of *Placoziphius* Van Beneden, 1869 and *Aulophyseter* Kellogg, 1927. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Sciences de la Terre*, **72**:151-170.
- Kemp, N.R. 1991. Chondrichthyan in the Cretaceous and Tertiary of Australia. In: P. Vickers-Rich; J.M. Monaghan; R.E. Baird & T.H. Rich (eds.) *Vertebrate Palaeontology of Australasia*, Pioneer Design Studio/Monash University Publications Committee, p. 498-531.
- Kent, B.W. 1994. *Fossil Shark of the Chesapeake Bay Region*. Maryland, Egan Rees & Boyer Inc., 146 p.
- Keyes, I.W. 1982. The Cenozoic sawshark *Pristiophorus lanceolatus* (Davis) (Order Selachii) of New Zealand and Australia, with a review of the phylogeny and distribution of world fossil and extant Pristiophoridae. *New Zealand Journal of Geology and Geophysics*, **25**:459-474. doi:10.1080/00288306.1982.10421510

- Le Roux, J.P.; Gómez, C.; Fennerand, J.H. & Middleton, H. 2004. Sedimentological processes in a scarp-controlled rocky shoreline to upper continental slope environment, as revealed by unusual sedimentary features in the Neogene Coquimbo Formation, North-Central Chile. *Sedimentary Geology*, **165**:67-92. doi:10.1016/j.sedgeo.2003.11.006
- Le Roux, J.P.; Olivares, D.M.; Nielsen, S.N.; Smith, N.D.; Middleton, H.; Fenner, J. & Ishman, S.E. 2006. Bay sedimentation as controlled by regional crustal behaviour, local tectonics and eustatic sea-level changes: Coquimbo formation (Miocene-Pliocene), Bay of Tongoy, central Chile. *Sedimentary Geology*, **184**:133-153. doi:10.1016/j.sedgeo.2005.09.023
- Lesueur, C.A. 1818. Description of several new species of North American fishes (part 1). *Journal of the Academy of Natural Sciences of Philadelphia*, **1**:222-235.
- Linnaeus, C. 1758. *Systema naturae*. 10th ed. Stockholm, Larentii Salvii, 824 p.
- Long, D.J. 1993. Late Miocene and early Pliocene fish assemblages from the North Central coast of Chile. *Tertiary Research*, **14**:117-126.
- Marquardt, C. 1999. *Neotectónica de la franja costera y aportes a la geología regional entre Caldera y Caleta Pajonal (27°-27°45' S), III Región de Atacama, Chile*. Departamento de Geología, Universidad de Chile, Tesis de grado, 297 p.
- Marsili, S. 2007. *Analisi sistematica, paleoecologica e paleobiogeografica della Selaciofauna plio-pleistocenica del Mediterraneo*. Dipartimento di Scienze della Terra, Università di Pisa, Tesi di dottorato, 245 p.
- Martínez-Pardo, R. 1979. Hallazgo de foraminíferos miocénicos cerca de Puerto Aldea, bahía de Tongoy, Provincia de Coquimbo, Chile. *Revista Geológica de Chile*, **8**:65-78.
- Morocco, R. & Muizon, C. 1988. Los vertebrados del Neogeno de La Costa Sur del Perú: Ambiente sedimentario y condiciones de fosilización. *Bulletin de l'Institut Frances d'Etudes Andines*, **17**:105-117.
- Muizon, C. 1978. *Arctocephalus (Hydrarctos) lomasiensis*, subgen. nov. et nov. sp., un nouvel Otariidae du Mio-Pliocene de Sacaco. *Bulletin de l'Institut Francais d'Etudes Andines*, **7**:169-189.
- Muizon, C. 1988. Les vertebres fossiles de la formation Pisco (Peru). Troisieme partie: les Odontocetes (Cetacea, Mammalia) du Miocene. *Editions Recherche sur les Civilisations*, **78**:1-244.
- Muizon, C. & DeVries, T.J. 1985. Geology and paleontology of late Cenozoic marine deposits in the Sacaco area (Peru). *Geologische Rundschau*, **74**:547-563. doi:10.1007/bf01821211
- Müller, J. & Henle, F.G.J. 1837. Gattungen der Haifische und Rochen nach einer von ihm mit Hrn. Henle unternommenen gemeinschaftlichen Arbeit über die Naturgeschichte der Knorpelfische. *Akademie der Wissenschaften zu Berlin*, **2**:111-118.
- Müller, J. & Henle, F.G.J. 1838. Ueber die Gattungen der Plagiostomen. *Archiv für Naturgeschichte*, **4**:83-85.
- Müller, J. & Henle, F.G.J. 1838-1841. *Systematische Beschreibung der Plagiostomen*. Berlin, Veit, p. 1-28, 1838d; p. 27-28 (reset), p. 29-102, 1839, p. 103-200.
- Nel, A. et al. 1999. Un gisement sparnacien exceptionnel à plantes, arthropodes et vertébrés (Éocène basal, MP7): le Quesnoy (Oise, France). *Comptes Rendus de l'Académie des Sciences - Series IIA - Earth and Planetary Science*, **329**:65-72. doi:10.1016/S1251-8505(99)80229-8
- Nishida, K. 1990. *Phylogeny of the Suborder Myliobatoidei*. Sapporo, Hokkaido University, p. 1-108 (Memoirs 37).
- Ohmi, K. 1986. Three fossil marine mammals from the Neogene of Sakiyama Peninsula, Ishikawa Prefecture, Central Japan. *Chikyu kagaku*, **40**:449-452.
- Paskoff, R. 1970. *Recherches géomorphologiques dans le Chili semi-aride*. Bordeaux, Biscaye, 420 p.
- Pequeño, G. 1989. Peces de Chile. Lista sistemática revisada y comentada. *Revista de Biología Marina*, **24**:1-132.
- Pequeño, G. & Sáez, S. 2003. The shark *Carcharhinus galapagensis* from Salas y Gómez Island, Chile (Chondrichthyes: Carcharhinidae). *Revista de Biología Tropical*, **51**:795-796.
- Phillipi, R.A. 1887. *Los fósiles Terciarios i Cuaternarios de Chile*. Leipzig, Brockhaus, 256 p.
- Pimiento, C. & Clements, C.F. 2014. When did *Carcharocles megalodon* become extinct? A new analysis of the fossil record. *PLOS ONE*, **9**: e111086. doi:10.1371/journal.pone.0117877
- Pimiento, C.; Ehret, D.J.; MacFadden, B.J. & Hubbell, G. 2010. Ancient nursery area for the extinct giant shark *Megalodon* from the Miocene Panama. *PLOS ONE*, **5**: e10552. doi:10.1371/journal.pone.0010552
- Pimiento, C.; González-Barba, G.; Ehret, D.J.; Hendy, A.J.W.; MacFadden, B.J. & Jaramillo, C. 2013b. Sharks and rays (Chondrichthyes, Elasmobranchii) from the late Miocene Gatun Formation of Panama. *Journal of Paleontology*, **87**:755-774. doi:10.1666/12-117
- Pimiento, C.; González-Barba, G.; Hendy, A.; Jaramillo, C.; MacFadden, B.; Montes, C.; Suárez, S., & Shippritt, M. 2013a. Early Miocene Chondrichthyes from the Culebra Formation, Panama: a window into marine vertebrate faunas before closure the Central American Seaway. *Journal of South American Earth Sciences*, **42**:159-170. doi:10.1016/j.jsames.2012.11.005
- Portell, R.W.; Hubbel, G.; Donovan, S.K.; Green, J.L.; Harper, D.A.T. & Pickerill, R. 2008. Miocene sharks in the Kendeace and Grand Bay formations of Carriacou, The Grenadines, Lesser Antilles. *Caribbean Journal of Science*, **44**:279-286.
- Purdy, R.W.; Schneider, V.P.; Applegate, S.P.; McLellan, J.H.; Meyer, R.L. & Slaughter, B.H. 2001. The Neogene sharks, rays, and bony fishes from Lee Creek Mine, Aurora, North Carolina. *Smithsonian Contributions to Paleobiology*, **90**:71-202.
- Quilodrán, P. & Marquardt, C. 2001. Catastro y condiciones de fosilización de cetáceos neógenos de la Formación Bahía Inglesa, 27-28S Tercera Región, norte de Chile. *Ameghiniana*, **38**:15-16.
- Radtke, U. 1989. Marine Terrassen und Korallenriffe. Das Problem der quartären Meeresspiegelschwankungen erläutert an Fallstudien aus Chile, Argentinien und Barbados. *Düsseldorfer Geographische Schriften*, **27**:1-246.
- Rafinesque, C.S. 1810. *Caratteri di alcuni nuovi generi e nuove specie di Animali e Pianta della Sicilia con varie osservazioni sopra I medesimi*. Palermo, Del Regno, 105 p.
- Reinecke, T.; Louwye, S.; Havekost, U. & Moths, H. 2011. The elasmobranch fauna of the late Burdigalian, Miocene, at Werder-Uesen, lower Saxony, Germany, and its relationships with early Miocene faunas in the North Atlantic, Central Paratethys and Mediterranean: *Palaeontos*, **20**:1-170.
- Risso, A. 1810. *Ichthyologie de Nice, ou histoire naturelle des poissons du département des Alpes Maritimes*. Paris, F. Schoell, 388 p.
- Rivadeneira, M.M. & Marquet, P.A. 2007. Selective extinction of late Neogene bivalves on the temperate Pacific coast of South America. *Paleobiology*, **33**:455-468. doi:10.1666/06042.1
- Rivadeneira, M.M. & Varas, C. 2012. *Línea base sector paleontológico del área marina y costera protegida de múltiples usos Isla Grande de Atacama*. Proyecto "Centro de Investigación y Conservación del Patrimonio Paleontológico del AMCP-MU-Isla Grande de Atacama", 33 p.
- Rojo, M. 1985. Un aporte al conocimiento del Terciario marino: Formación Bahía Inglesa. In: CONGRESO GEOLÓGICO CHILENO, 4, 1985. *Resúmenes*, p. 514-533.

- Rüppell, E. 1837. *Neue Wirbelthiere zu der Fauna von Abyssinien gehörig. Fische des Rothen Meeres*. Frankfurt am Main, 1-148, p. 1-33. 1835:1-28, p. 1-7; 1836:29-52, p. 8-14; 1837:53-80, p. 15-21; 1838:81-148, p. 22-33.
- Schindler, T.; Nungesser, K.; Mueller, A. & Grimm, K.I. 2009. Die Alzey-Formation der klassischen Lokalität Walschberg bei Waldböckelheim (Rupelium, Oligozän, Mainzer Becken)-Ergebnisse neuer Grabungen. *Jahresberichte und Mitteilungen des Oberrheinischen Geologischen Vereines*, **91**:37-87. doi: 10.1127/jmogn/91/2009/37
- Snodgrass, R.E. & Heller, E. 1905. Papers from the Hopkins-Stanford Galapagos Expedition, 1898-1899. XVII. Shore fishes of the Revillagigedo, Clipperton, Cocos and Galapagos Islands. *Proceedings of the Washington Academy of Sciences*, **6**:333-427.
- Suárez, M.E. & Brito, J.L. 2000. Una nueva localidad con vertebrados fósiles del Terciario de Bahía Salado, Chile. In: JORNADAS ARGENTINAS DE PALEONTOLOGÍA DE VERTEBRADOS, 15, 2000. *Resúmenes*, San Luis, p. 57.
- Suárez, M.E. & Encinas, A. 2002. Vertebrados marinos del miembro inferior de la Formación Navidad (Mioceno temprano), Chile Central. In: CONGRESO LATINOAMERICANO DE PALEONTOLOGIA DE VERTEBRADOS, 5, 2002. *Resúmenes*, Santa Cruz de la Sierra, p. 51-52.
- Suárez, M.E.; Encinas, A. & Ward, D. 2006. An early Miocene elasmobranch fauna from the Navidad Formation, Central Chile, South America. *Cainozoic Research*, **4**:3-18.
- Suárez, M.E.; Lamilla, J. & Marquardt, C. 2004. Peces Chimaeriformes (Chondrichthyes, Holocephali) del Neógeno de la Formación Bahía Inglesa, Región de Atacama, Chile. *Revista Geológica de Chile*, **31**:105-117. doi: 10.4067/s0716-02082004000100006
- Suárez, M.E. & Marquardt, C. 2003. Revisión preliminar de las faunas de peces elasmobranchios del Mesozoico y Cenozoico de Chile: su valor como indicadores cronoestratigráficos. In: CONGRESO GEOLÓGICO CHILENO, 10, 2003. *Actas*, Concepción, p. 9.
- Taniuchi, T. 1990. The role of elasmobranchs in Japanese fisheries. In: H.L. Pratt; S. Gruber & T. Taniuchi (eds.) *Elasmobranchs as living resources: advances in the biology, ecology, systematics, and the status of the fisheries*, NOAA, p. 415-426 (Technical Report 90).
- Tsuchi, R. 2002. Neogene evolution of surface marine climate in Pacific and notes on related events. *Revista Mexicana de Ciencias Geológicas*, **19**:260-270.
- Underwood, C.J.; Michell, S.F. & Veltkamp, K.J. 1999. Shark and Ray teeth from the Hauterivian (lower Cretaceous) of North-East England. *Palaeontology*, **42**:287-302. doi: 10.1111/1475-4983.00074
- Valenzuela-Toro, A.M.; Gutstein, C.S.; Varas-Malca, R.M.; Suárez, M.E. & Pyenson, N.D. 2013. Pinniped turnover in the South Pacific Ocean: new evidence from the Plio-Pleistocene of the Atacama Desert, Chile. *Journal of Vertebrate Paleontology*, **33**:216-223. doi: 10.1080/02724634.2012.710282
- Villafañá, J.A. & Rivadeneira, M.M. 2013. Cambios paleobiogeográficos de elasmobranchios del Neógeno en el Pacífico de Sudamérica. In: CONGRESO DE CIENCIAS DEL MAR, 33, 2013. *Resúmenes*, Antofagasta, p. 60.
- Villafañá, J.A. & Rivadeneira, M.M. 2014. The rise and fall of the diversity of marine vertebrates during the Neogene on the temperate Pacific coast of South America. *Paleobiology*, **40**:659-674. doi: 10.1666/13069
- Walsh, S.A. 2001. *The Bahía Inglesa formation bonebed: genesis and palaeontology of a Neogene konzentrat lagerstätte from North-Central Chile*. University of Portsmouth, Ph.D. thesis, 443 p.
- Walsh, S. & Hume, J. 2001. A new Neogene marine avian assemblage from North-Central Chile. *Journal of Vertebrate Paleontology*, **21**:484-491. doi: 10.1671/0272-4634(2001)021[0484:annmaaj2.0.co;2
- Walsh, S. & Naish, D. 2002. Fossil seals from late Neogene deposits in South America: a new pinniped (Carnivora, Mammalia) assemblage from Chile. *Paleontology*, **45**:821-842. doi: 10.1111/1475-4983.00262
- Ward, D.J. & Bonavia, C.G. 2001. Additions to, and a review of, the Miocene shark and ray fauna of Malta. *The Central Mediterranean Naturalist*, **3**:131-146.
- Welton, B.J. & Zinsmeister, W.J. 1980. Eocene Neoselachians from the La Meseta Formation, Seymour Island, Antarctic Peninsula. *Contributions in Science, Los Angeles County Museum*, **329**:1-10.
- Whitmore, F. 1987. Cetacea from the Sahabi formation. In: N.T. Boaz; A. El-Arnauti; A.W. Gaziry; J. Heinzelin & D.D. Boaz (eds.) *Neogene Paleontology and Geology of Sahabi*, Liss, p. 145-151.
- Zachos, J.; Pagani, M.; Sloan, L.; Thomas, E. & Billups, K. 2001. Trends, rhythms, and aberrations in global climate 65 Ma to Present. *Science*, **292**:686-693. doi: 10.1126/science.1059412

Received in September, 2014; accepted in June, 2015.

APPENDIX II

The Path towards Endangered Species: Prehistoric Fisheries in Southeastern Brazil

RESEARCH ARTICLE

The Path towards Endangered Species: Prehistoric Fisheries in Southeastern Brazil

Mariana Samôr Lopes¹, Thayse Cristina Pereira Bertucci^{1*}, Luciano Rapagnã¹, Rafael de Almeida Tubino¹, Cassiano Monteiro-Neto¹, Acácio Ribeiro Gomes Tomas², Maria Cristina Tenório³, Tânia Lima³, Rosa Souza¹, Jorge Domingo Carrillo-Briceño⁴, Manuel Haimovici⁵, Kita Macario⁶, Carla Carvalho⁶, Orangel Aguilera Socorro¹

1 Universidade Federal Fluminense, Instituto de Biologia, Campus do Valonguinho, Outeiro São João Batista, s/n°. CEP: 24020–141, Niterói, Rio de Janeiro, Brasil, **2** Instituto de Pesca, Centro APTA Pescado Marinho, Av. Bartolomeu de Gusmão 192, Santos, São Paulo, CEP: 11030–906, Brasil, **3** Universidade Federal do Rio de Janeiro, Museu Nacional, Departamento de Antropologia, Quinta da Boa Vista, Rio de Janeiro, CEP: 20940–040, Brasil, **4** Palaeontological Institute and Museum, University of Zürich, Karl-Schmid-Strasse 4, Zürich 8006, Switzerland, **5** Universidade Federal do Rio Grande, Instituto de Oceanografia, Campus Carreiros, Av. Itália, Rio Grande, Rio Grande do Sul, CEP: 96201–900, Brasil, **6** Universidade Federal Fluminense, Instituto de Física, Campus da Praia Vermelha, Boa Viagem, CEP: 24210–310, Niterói, Rio de Janeiro, Brasil

* thathab@gmail.com



OPEN ACCESS

Citation: Lopes MS, Bertucci TCP, Rapagnã L, Tubino RdA, Monteiro-Neto C, Tomas ARG, et al. (2016) The Path towards Endangered Species: Prehistoric Fisheries in Southeastern Brazil. PLoS ONE 11(6): e0154476. doi:10.1371/journal.pone.0154476

Editor: John P. Hart, New York State Museum, UNITED STATES

Received: December 18, 2015

Accepted: April 14, 2016

Published: June 29, 2016

Copyright: © 2016 Lopes et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files. All specimens studied here were deposited in the zooarchaeological collection of Museu Nacional, Universidade Federal do Rio de Janeiro (MNUFRJ-ZA), Parque da Boa Vista, São Cristóvão, Rio de Janeiro, Brazil. <http://www.museunacional.ufrj.br/> - refers to [S1 Appendix](#). All relevant species specific characters were identified based on comparative anatomy, using otoliths, dry finfish skeletons and shark tooth collections housed at the Universidade Estadual do Rio de Janeiro (UERJ) and Universidade

Abstract

Brazilian shellmounds are archaeological sites with a high concentration of marine faunal remains. There are more than 2000 sites along the coast of Brazil that range in age from 8,720 to 985 cal BP. Here, we studied the ichthyoarchaeological remains (i.e., cranial/post-cranial bones, otoliths, and teeth, among others) at 13 shellmounds on the southern coast of the state of Rio de Janeiro, which are located in coastal landscapes, including a sandy plain with coastal lagoons, rocky islands, islets and rocky bays. We identified patterns of similarity between shellmounds based on fish diversity, the ages of the assemblages, littoral geomorphology and prehistoric fisheries. Our new radiocarbon dating, based on otolith samples, was used for fishery characterization over time. A taxonomical study of the ichthyoarchaeological remains includes a diversity of 97 marine species, representing 37% of all modern species (i.e., 265 spp.) that have been documented along the coast of Rio de Janeiro state. This high fish diversity recovered from the shellmounds is clear evidence of well-developed prehistoric fishery activity that targeted sharks, rays and finfishes in a productive area influenced by coastal marine upwelling. The presence of adult and neonate shark, especially oceanic species, is here interpreted as evidence of prehistoric fisheries capacity for exploitation and possibly overexploitation in nursery areas. Various tools and strategies were used to capture finfish in seasonal fisheries, over rocky reef bottoms and in sandy littoral environments. Massive catches of whitemouth croaker, main target demersal species of South Atlantic coast, show evidence of a reduction in body size of approximately 28% compared with modern fisheries. Fishery activity involving vulnerable species, especially in nursery areas, could mark the beginning of fish depletion along the southeastern Brazilian coast and the collapse of natural fish populations.

Federal Fluminense (UFF), see more details in the [S2 Appendix](#). Relevant species specific osteological characters were also identified based on extensive bibliographic review.

Funding: This work was supported by Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro, APQ1, Process E-26/111.278/2014 to OA, <http://www.faperj.br/>; Pró-Reitoria de Pesquisa, Pós-Graduação e Inovação da Universidade Federal Fluminense, Process 46094044157201298 to OA, <http://www.propqi.uff.br/posgraduacao/>; and Conselho Nacional de Desenvolvimento Científico e Tecnológico, Fellowship, Process 306053/2013-6 to CMN; Process 305037/2012-9 to MCT, <http://lattes.cnpq.br/>. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Introduction

When investigating early archaeological settlements in Brazil, South America [1], the existence of an undisturbed marine fauna predating European colonization is expected. The archaeological evidence of prehistoric fisheries shows high abundance and diversity of marine faunal remains recovered from Brazilian shellmounds [2,3,4,5,6,7,8,9,10,11,12]. However, the main goal of these studies is to elucidate the archaeological context of fisher-gatherer settlements. The ichthyoarchaeological remains testify to well-developed fisheries for sharks, rays and fin-fishes, as well as to shellfish gathering and hunting of marine reptiles and mammals. Prehistoric subsistence fisheries could have been the cause of early differential disturbances in local fish fauna resources due to the assumed use of beach seines, gillnets, hook and line, traps and spearfishing.

A key study on prehistoric fisheries in the Caribbean [13] strongly supports the claim that overexploitation did not occur. However, the debate about fish depletion, overexploitation, extinction and environmental degradation in prehistoric and colonial times continues [14,15]. Prehistoric overfishing could be associated with local environmental degradation as a consequence of human uses of the landscape affecting the marine environment, community dynamics and spatially subsidized food webs [16,17,18,19], and overexploitation based exclusively on prehistoric fisheries might not have been the exclusive cause [15,20,21]. This hypothesis was widely discussed using evidence from comparative analyses of Pacific islands and channel ecosystems under archaeological investigation [22,23]. However, tropical Western Atlantic prehistoric fisheries differ in terms of Holocene paleoenvironments, fish assemblages and large-scale vulnerability at the time of prehistoric fisheries along the South American coast ([1]; [S1 Appendix](#)).

Prehistoric fishery exploitation patterns along the Atlantic coast of South America, from Brazil to Argentina are not well described. Brazilian shellmounds ranges from 8,720 to 985 years calibrated before present (cal BP) [1,24,25,26,27,28,29,30,31]. Previous ichthyological records from these archaeological sites were documented only by faunal lists without illustrative diagnostic species characters or museum catalog numbers to corroborate the species identifications. The present contribution is an attempt to improve our understanding of Rio de Janeiro's shellmounds by providing new, detailed and accurate taxonomic lists and analyses of the relevant fish assemblages.

Contiguous with south Brazil and Uruguay, Argentinean ichthyoarchaeological sites, situated on the coast of San Matías Gulf in Patagonia (6,800 to 890 yr BP), show evidence of local prehistoric fisheries. The abundance of otoliths indicates that the predominant bony fish target was the whitemouth croaker, *Micropogonias furnieri* [32,33,34,35], a coastal finfish species also present in Brazilian coastal shellmounds.

Fishery tools, such as projectile points, manufactured mainly with bones are frequently recovered from the Brazilian shellmounds [30]. Moreover, evidence that allows us to infer the use of wood and vegetable fiber for the construction of fish traps, beach seines and or gillnets used for massive captures of large fish schools is not preserved. Indirect evidence of boat constructions based on lithic tools and records in detailed rock paintings reveal fluvial skillful navigators in Northeastern of Brazil [36]. In addition, the frequency of skeletal anomalies found in human remains recovered from the shellmounds, including auditory meatus exostoses, osteoarthritis, osteoarthritis and other degenerative effects [37,38], is usually considered to be a marker of sailing and aquatic labor in cold waters [39,40,41], suggesting the engagement of those human populations in traditional fishery activity for subsistence. Stable isotope analyses of $\delta^{14}\text{C}$ and $\delta^{15}\text{N}$ in human skeletons from the shellmounds of Southern Brazil indicate a diet strongly dependent on marine resources [42]. Therefore, in agreement with previous research

[43], the abundance and diversity of fish remains from the shellmounds provides unequivocal proof of fisheries that were able to operate in open waters over sandy and rocky bottoms.

Shellmounds are not necessarily horizontally stratified due to sequential periods of occupation, and the areas selected for specific activities may vary from the center to the periphery of the shellmound [44,45]. These sandy shellmounds are usually dome-shaped, and the archaeological variation in vertical section is based on changes in sediment texture and color, settlement size, abundance and diversity of mollusks, intercalation of sterile sandy layers (i.e., without ichthyoarchaeological remains) and evidence of cultural activities (e.g., burials and stoves). However, in contrast with archaeological evidence from settlements, in some shellmounds, the layers could not be distinguished [46].

The Rio de Janeiro shellmounds [1] are located in a coastal landscape characterized by sandy plains with coastal lagoons, rocky islands, islets and rocky bays [47]. These coastal areas are strongly influenced by seasonal upwellings, occurring during the austral summer [48,49,50,51], which increase marine productivity and the potential resources for fisheries.

Therefore, the main goal of this study is to highlight the fish diversity and faunal assemblage of ancient fisheries based on the skeletal remains deposited in the shellmounds along the south-eastern Brazilian coast. Additionally, we correlate shark diversity and abundance in the shellmounds with the abilities of specific fisheries to exploit resources, which could lead to overexploitation. The groundfish catches were the result of multi-gear strategies in seasonal fisheries. Radiometric age, geochemical analyses and climate reconstruction of these ichthyoarchaeological remains [52,53,54,55] were used to supporting our findings.

Historical accounts of ancient fisheries

Ichthyoarchaeological evidence and historical accounts from the Caribbean and tropical South America suggest that early prehistoric target species corresponded to the most accessible and vulnerable animals, such as sharks, large groupers [56,57,58], turtles [59,60] and sea mammals [61,62,63,64].

Colonial accounts from 1587 [56] on the semi-sedentary indigenous community that lived in Brazil during the early days of Portuguese colonization refer to the paleo-Indian fisheries' expertise and their use of rich marine food supplies; the accounts additionally demonstrate a traditional knowledge of fish diversity, reproductive aggregation of fish, fishery areas and environmental relationships. Regarding fish diversity, a list of the 43 most important species of fish was reported accurately (Table 1) [56]. These fish records are in agreement with the prehistoric data presented here. Most of the historic narrative regarding fish captures concerns reproductive periods in coastal and estuarine areas, where bony fishes form compact aggregations along the littoral zone during the intertidal phase. The account '*curiosi rerum naturae*' [56] refers to high marine fish abundance, especially during the summer on the coast of Salvador in the state of Bahia (northeast Brazil), when female bony fishes have large gonads. Additionally, the narrative talks about the 'docile and very easily caught giant grouper on the beaches', the large tarpon size, 'longer than an Indian is tall', and the 'thousands of mullet caught during a single day fishing', revealing a picture of the abundance of fishery resources [56,65]. All these historical narratives are in agreement with the fact that Brazilian neo-Indians were skilled fishermen, using arrows, marksmanship, and fishery lines with hooks; they built fish traps with wood and rocks in an intertidal beach and small nets for cooperative fisheries. These undeniable skills were possibly inherited from ancestral paleo-Indians who perfected the art of fishing [66].

Shellmounds are the best testimonial resource for understanding the paleo-Indian fishery activities. However, most of the original context of Brazilian shellmounds was destroyed due to the use of mollusk shells to produce lime and fertilizers for paving of roads and streets,

Table 1. Brazilian colonial fish records [56].

Indigenous names	Probable species	Fishery tool and remarks
aragoagoay	<i>Pristis</i> sp.	hooks and spear
uperu, panapaná, socorí	shark	hooks and spear
beijupirá	<i>Rachycentron canadum</i>	hooks
tapyrsiçá	<i>Seriola lalandi</i>	hooks
camuropi	<i>Megalops atlanticus</i>	hooks, very large
piraquirolá	<i>Selene</i> sp.	hooks
carapitanga	<i>Lutjanus</i> sp.	hooks
canapú	<i>Epinephelus itajara</i>	hooks, tides stones and sticks tramp, very large, easy capture
cupá	<i>Cynoscion</i> sp.	hooks
guaripicú	<i>Scomberomorus</i> sp.	trolling lines
guiará	<i>Chaetodipterus faber</i>	hooks and beach seine
guris and urutús	Ariidae	hooks
caramurú	<i>Echidna</i> sp.	hands
jabubirá	Dasyatidae or Myliobatidae	hooks and beach seine
tacupapirema	<i>Micropogonias furnieri</i> or <i>Cynoscion acoupa</i>	hooks
bonitos	Carangidae	hooks
dourada	<i>Coryphaena hippurus</i>	hooks
caraoatá	<i>Thunnus</i> sp.	hooks
garoupas	<i>Epinephelus</i> sp.	hooks, very large
camurís	<i>Centropomus</i> sp.	hooks
abróteas	<i>Urophycis brasiliensis</i>	hooks
ubaranas	<i>Elops saurus</i>	hooks
goaivicoára	<i>Conodon nobilis</i>	hooks
sororocas	<i>Scomberomorus maculatus</i>	hooks
timaçu	<i>Strongylura</i> sp.	used for bait
miracoia	<i>Stellifer</i> sp. or <i>Bairdiella</i> sp.	hooks
maracugua	<i>Balistes</i> sp.	hooks
paratís	<i>Mugil curema</i>	sticks tramp and net during high tide
zabucaí	<i>Selene</i> sp.	beach seine
tareira	<i>Caranx hippos</i>	beach seine
coirimás	<i>Mugil liza</i>	beach seine
arabori	<i>Brevoortia aurea</i>	beach seine
carapebas	<i>Eucinostomus</i> sp.	beach seine
jaguaracá	<i>Holocentrus adscensionis</i>	hooks, medicinal
piraçaque	<i>Conger</i> sp.	hooks, medicinal
bodiaens	<i>Scarus</i> sp.	hooks, medicinal
atucupá	<i>Cynoscion</i> sp.	hooks, medicinal
goayibicoati	Gobiidae	hooks, medicinal
uramaçã	Paralichthyidae	hooks, medicinal
baiacú	<i>Lagocephalus</i> sp.	fishes that producing poisoning
piraquirolá	<i>Chilomycterus antillarum</i>	fishes that producing poisoning
aimoré	Gobiidae	fishes that producing poisoning

The first ichthyofaunal list (indigenous name) documented from the Brazilian coast.

doi:10.1371/journal.pone.0154476.t001

construction of forts, colonial houses, churches, among others applications. This occurred starting in 1549 in the area of Salvador, Bahia and other colonial settlements along the Brazilian coast and continued until the 1960s, when archaeological shellmounds became protected under Brazilian federal law.

Geographic setting

From south to north, the landscape of the Rio de Janeiro coast is characterized by the presence of a crystalline shield (Serra do Mar relief), with plenty of high (approximately 1,200 m) mountain scarps parallel to the Atlantic Ocean in the vicinity of Angra dos Reis [67]. The area is characterized by short rivers flowing to the coast and fluvio-marine plains in an embayment (i.e., Ribeira Bay), a jagged coastline, with small peninsulas and several rocky islets [68]. The beaches and the sandy stretches are not developed, and the shellmounds are predominantly located over the rocky islets in an area dominated by mangrove flood plains. Small submarine channels approximately 6 to 10 m deep characterize the coastal bathymetry, and near Ilha Grande bay, the depth reaches approximately 30 m. Ilha Grande is a massive structural island with fairly rugged relief; it is isolated from the mainland by a channel approximately two kilometers wide.

The Itaipú-Camboinhas region is located on the oceanfront of Niterói and has sandy beaches dominated by dunes and sandy bars, which separate the sea from Piratininga and Itaipu coastal lagoons. The semicircular sandy beach has depths of approximately 3 to 16 m that extend almost 1,000 m offshore. The landscape has a mountainous relief aligned in the SW-NE direction [47]. The Camboinhas shellmound is located over a sand dune, near the coastal line and the tidal channel of Itaipu Lagoon.

The Saquarema region is located in a landscape characterized by a crystalline rock relief, which separates the two major drainage basins that feed the Saquarema lagoon complex [47,68]. The sandy shore is shallow but exposed to high-energy coastal wind and waves. The area has the highest concentration of shellmounds in Rio de Janeiro; they are distributed along the sandy coastal plain of the inland sandbanks, facing the lagoon (e.g., Beirada, Manitiba, Ponte do Girau and Saquarema shellmounds).

In Arraial do Cabo, the structural NE-SW trend is characterized by a metamorphic basement (i.e., Pontal de Atalaia), rising up to 172 m high, and the adjacent Cabo Frio Island, an igneous alkaline rock (syenite, trachyte and breccia) with altitudes of approximately 380 m [69]. The Usiminas shellmound is located in the Cabo Frio Island, 50 m above sea level, facing the coastal plain, where the water depth is approximately 5 m. In contrast, the oceanfront cliffs on the opposite coast reach 50 m in depth near the coastline. Additionally, the Ilha do Cabo Frio shellmound is located on a small sandy beach characterized by an active dune that faces towards the landscape, associated with the outcrop layers that overlap the Cabo Frio beach rock [65,70]. Part of the lower layer of this shellmound is located below sea level [29].

Materials and Methods

Selected shellmounds from the southeastern coast of Rio de Janeiro State, Brazil have three main features: (1), the potential marine influence of the Cabo Frio upwelling system (i.e., seasonal oceanographic mixing of South Atlantic Central Water, Subtropical Shelf Water and the Brazilian Current, increasing biological productivity); (2), the marine environment (i.e., shallow waters, coastal lagoons and a rocky bottom); and (3), coastal geomorphology (i.e., sandy coastal plains, rocky islands and rocky bays).

The shellmounds included in this study are the following: the Usiminas shellmound [71], on a rocky settlement, and the Ilha do Cabo Frio shellmound [72], on a sandy beach on Cabo

Frio Island (23°00' 18" S, 42°00' 20" W); Saquarema [72], Beirada [46], Manitoba [73] and Ponte do Girau [74] shellmounds, on a sandy coastal plain with coastal lagoons in the Saquarema lagoon complex (22°55' 66" S, 42°29' 00" W); Camboinhas shellmound [75], on a sandy coastal plain with coastal lagoons in the oceanic region of Niterói (22°57' 54" S, 44°02' 53" W); Algodão, Major, Bigode, Caieira and Peri shellmounds [76], on rocky islets and coastal rocky bays in the Ribeira Bay, Angra dos Reis (22°55' 48" S, 44°20' 48" W); and Acaia shellmound (personal communication of an unpublished manuscript: Tenório, M.C. 'Os sambaquieiros e a gruta do Acaia: Reconstituição do processo de formação de um sítio'), on a rocky island in the oceanfront of Ilha Grande (Fig 1).

We studied all the specimens deposited in the ichthyoarchaeological collection of Museu Nacional, Universidade Federal do Rio de Janeiro (MN-UFRJ), Rio de Janeiro, Brazil, and their use for this research was authorized by the collection managers, who are coauthors of this study (MCT and TL).

We have organized a referential species collection based on 679 diagnostic osteological and dental characters (details in S2 Appendix). Quantitative analyses of fish specimens and species recovered from each shellmound are necessary for accurate data interpretation. However, different archaeological methods were used in these shellmounds during excavation conducted by UFRJ archaeologists between 1981 and 2005. The methods included sieving techniques, removing material and curatorial processing. These and others questions regarding the repository, catalog and samples labels will require more detailed assessment, which is beyond of the scope of this study. Consequently, the fish diversity analyzed here is based on a qualitative study and the frequencies of species by shellmound.

All relevant specific characteristics of examined species were identified based on comparative anatomy, using 39 shark and ray specimens (among teeth, vertebrae and spines) housed at the Universidade Estadual do Rio de Janeiro (UERJ) and using at least 115 otoliths and 21 dry finfish individual skeletons housed at the Universidade Federal Fluminense (UFF) (details in S3 Appendix). All structures are in a good state of conservation and relevant specific osteological characteristics were also identified based on extensive bibliographic review.

Photographs of the otoliths were taken with a Leica M205A multifocal stereomicroscope. Photographs of bones, sharks and rays teeth were taken using a digital microscope and digital camera. A complete atlas of the most common diagnostic teeth, otoliths and bones recovered and observed from shellmound collections is included in the plates of the present paper.

Cluster analysis was performed under the Paleontological Statistics Software (PAST, version 2.17c) on Q-mode (i.e., grouping variables) to analyze shellmound similarity patterns. This exploratory technique identifies the relationships and patterns among multiple variables across samples and has been applied in a wide range of scientific fields, such as marine and fisheries ecology [77,78,79,80]. The analysis was based on the presence and absence of fish assemblages in the shellmounds, archaeological site ages, littoral geomorphology and prehistoric fisheries. The unweighted pair-group average (UPGMA) algorithm was used with the Bray-Curtis similarity-association matrix of [81].

Estimates of shark-body total length (TL) were based on 660 isolated vertebrae, using a unique linear regression equation for each species: *Carcharias taurus* TL = 36.786 + 10.753 CR [82], *Sphyrna lewini* TL = 4.51 + 23.64 CR [83], *Carcharhinus brevipinna* CD = 0.0159 PCL - 0.1285, PCL = 0.799 TL - 9.07 [84] and *Carcharodon carcharias* FL = 21 + 11.8 CR, FL = 0.94 TL - 5.74 [85], where TL is the total length, FL is the fork length, PCL the precaudal length, CD the vertebral centrum diameter and CR the vertebral centrum radius.

The main teleostean fish target, based on the frequency observed in the shellmounds, was the whitemouth croaker, *Micropogonias furnieri* [86], which was present in all shellmounds with the exception of the Usiminas and Ilha do Cabo Frio shellmounds, both on Cabo Frio




Fig 1. Study area along the southeastern Brazilian coast showing shellmound locations.

doi:10.1371/journal.pone.0154476.g001

Island. The life history of *M. furnieri* in the Western Atlantic Ocean is well known [87,88], and they can be found in commercial [89,90] and local artisanal fisheries [77,91]. We used this species to interpret and compare changes between past and present coastal fisheries on the southeastern Brazilian coast. We tested for differences between the medians of length frequency distributions (Student's t-test, PAST software v. 3.7) after checking for normality and homogeneity of variances.

A total of 5,532 archaeological whitemouth croaker, *Micropogonias furnieri*, otoliths were measured using digital callipers. These specimens were distributed among the shellmounds as follows: Ponte do Girau (376 specimens), Beirada (2,541 specimens), Manitiba (1,372 specimens), Algodão (1,148 specimens) and Camboinhas (95 specimens). Fish total lengths (TL) were calculated based on the archaeological otolith total length (OL) using the following equation, which we derived from a regression analysis: $TL = 24.34 + 22.57 OL$ ($r = 0.988$, $n = 93$). To compare the estimates of body size length data between shellmound samples, we performed a nonparametric variance analysis of medians (Kruskal-Wallis test) and an *a posteriori* test of the shellmound context, analyzing the localities and related environmental factors using the free software R [92].

We chose not to use the South American Western Atlantic fish records [88]; instead, specific references of modern fish diversity and abundance in the littoral areas close to the shellmounds

of Rio de Janeiro were obtained from Arraial do Cabo [58,93,94], Itaipu [77,95] and Ribeira Bay in Angra dos Reis [96], and these were used in comparisons with fishery records.

Data from the Brazilian fishery survey of whitemouth croaker in the Itaipu modern fisheries (i.e., artisanal and semi-industrial) are based on records compiled between 2000 and 2004 [77] from the Angra dos Reis Bay from semi-industrial fisheries captures reported between 2013 and 2014. At Guanabara Bay, a fishery study was conducted using a bottom trawl between 2005 and 2007 [91].

The samples for radiochronology were prepared and analyzed at the Radiocarbon Laboratory of the Universidade Federal Fluminense (LAC-UFF). The otolith samples were chemically treated with HCl and converted to CO₂ by hydrolysis with H₃PO₄. Graphitized samples were placed in the 40-sample wheel of the SNICS ion source and measured in an NEC 250 kV Single Stage Accelerator System (SSAMS) [97]. Typical currents were 50 $\mu\text{A}^{12}\text{C}^{-1}$ (measured at the low energy Faraday cup). Graphite standard and calcite blanks yielded average $^{14}\text{C}/^{13}\text{C}$ ratios of 6×10^{-13} and 7×10^{-13} , respectively. The average machine background was approximately 50 kHz for the unprocessed graphite, while the average precision ranged from 0.3 to 0.5%. Data analyses were carried out on LACAMS software developed at the Physics Institute of Universidade Federal Fluminense [98]. Calibration of otolith radiocarbon dates was performed with OxCal software v 4.2.3 from the Oxford University [99], using the Marine13 curve [100] in the 2-sigma range with an offset for local marine reservoir corrections (ΔR ^{14}C years) according to the following shellmound localities: Saquarema [53], Manitiba [55] and Southeastern Brazil [101]. Cross-comparisons were conducted in the Beta Analytic Inc. laboratory, following standard protocols.

We exclusively used the fish otolith radiocarbon dates to determine ages for chronological interpretation of the fish assemblages of most shellmounds, except for those from the Cabo Frio Island, where otoliths of whitemouth croaker could not be recovered; additionally, a few otoliths from Ilha Grande could not be dated. Consequently, we refer only to ages that were not derived from otolith samples for the Usiminas, Ilha do Cabo Frio and Acaia shellmounds. The age ranges of those sites are based on charcoal and shell data ([29,102]; personal communication).

Results

Shellmound radiocarbon dating based on otoliths and the identified fish assemblages are shown in Table 2. Manitiba serves as an example of a multilayer shellmound (i.e., seven archaeostratigraphic layers in 2.2 m of depth) and demonstrates the difficulty in distinguishing radiocarbon age differences between successive layers; here, the calibration curve associated with the error bar and the reservoir effect cause the age range to overlap (Fig 2). This inhibits inferring a chronological sequence for fisheries in shallow shellmounds. We, therefore, use individual shellmound dating as a marker of settlement period.

A total of 97 fish species have been identified from the Rio de Janeiro shellmounds, representing 37% of the total (265 spp.) modern species recorded from the Rio de Janeiro coast [e.g., Arraial do Cabo (135 spp.), Itaipu (165 spp.) and Angra dos Reis (139 spp.)]. Table 2 shows the occurrence in archaeological sites of the great white shark, *Carcharodon carcharias*, the sand tiger shark, *Carcharias taurus*, the porbeagle shark, *Lamna nasus*, the lemon shark, *Negaprion brevirostris*, the tiger shark, *Galeocerdo cuvier*, and a large diversity of other sharks (such as Carcharhinidae, Sphyrnidae, and other coastal species) in the rocky islet shellmounds of Angra dos Reis (Ribeira Bay: Algodão, Bigode, Caieira, Major and Peri sites), Ilha Grande (Acaia) and Cabo Frio Island (Usiminas and Ilha do Cabo Frio). The data also include the occurrence of other rarer shark species found in the shellmounds located over sandy coasts and coastal

Table 2. Ichthyoarchoaeological records from the Rio de Janeiro shellmounds (Saquarema, Cabo Frio, Niterói, Ilha Grande and Angra dos Reis).

Region	SAQUAREMA				NITEROI	ANGRA DOS REIS				ILHA GRANDE		ILHA DO CABO FRIO		Structures	Figures	
	Shellmounds		Geomorphology			Camboinhas	Algodão	Bigode	Caleira	Peri	Major	Acaia	Usiminas			Ilha do Cabo Frio
	Beirada	Saquarema	Manitiba	Girau												
Radiocarbon age ranges	3035 to 5595 cal BP	2100 to 4200 cal BP	3695 to 4515 cal BP	3730 to 4525 cal BP	4160 to 4960 cal BP	2345 to 4414 cal BP	3223 to 3525 cal BP	1875 to 2175 cal BP	890 to 1140 cal BP	675 to 900 cal BP	2760 to 2930 cal BP*	1265 to 1765 cal BP*	2710 to 3290 cal BP*			
	Chondrichthyes															
Odontaspidae	<i>Carcharias taurus</i>															
Alopiidae	<i>Alopias superciliosus</i>															
Lamnidae	<i>Carcharodon carcharias</i>															
Carcharhinidae	<i>Isurus oxyrinchus</i>															
	<i>Lamna nasus</i>															
	<i>Carcharhinus acronotus</i>															
	<i>Carcharhinus altimus</i>															
	<i>Carcharhinus brevipinna</i>															
	<i>Carcharhinus leucas</i>															
	<i>Carcharhinus limbatus</i>															
	<i>Carcharhinus obscurus</i>															
	<i>Carcharhinus plumbeus</i>															
	<i>Carcharhinus porosus</i>															
	<i>Carcharhinus priscus</i>															
	<i>Carcharhinus</i> sp.															
	<i>Galeocerdo cuvier</i>															
	<i>Negaprion brevirostris</i>															
<i>Rhizoprionodon landi</i>																
<i>Rhizoprionodon porosus</i>																

(Continued)

Table 2. (Continued)

Region	SAQUAREMA				NITEROI		ANGRA DOS REIS				ILHA GRANDE		ILHA DO CABO FRIO		Structures	Figures
	Beirada	Saquarema	Manitiba	Girau	Cambolinas	Algodão	Bigode	Caleira	Peri	Major	Acaia	Usiminas	Ilha do Cabo Frio			
Shellmounds																
Geomorphology	Sandy coastal plain and coastal lagoons				Sandy beach	Rocky bottom and rocky islet				Rocky island						
	3035 to 5595 cal BP	2100 to 4200 cal BP	3695 to 4515 cal BP	3730 to 4525 cal BP	4160 to 4960 cal BP	2345 to 4414 cal BP	3223 to 3525 cal BP	1875 to 2175 cal BP	890 to 1140 cal BP	675 to 900 cal BP	2760 to 2930 cal BP*	1265 to 1765 cal BP*	2710 to 3290 cal BP*			
Radiocarbon age ranges																

(Continued)

Table 2. (Continued)

Region	SAQUAREMA			NITEROI		ANGRA DOS REIS				ILHA GRANDE		ILHA DO CABO FRIO		Structures	Figures
	Shellmounds	Beirada	Saquarema	Manitiba	Girau	Cambolhas	Algodão	Bigode	Caleira	Peri	Major	Acalá	Usiminas		
Geomorphology	Sandy coastal plain and coastal lagoons			Sandy beach			Rocky bottom and rocky islet							Rocky Island	
	3035 to 5595 cal BP	2100 to 4200 cal BP	3695 to 4515 cal BP	3730	4160 to 4960 cal BP		2345 to 4414 cal BP	3223 to 3525 cal BP	1875 to 2175 cal BP	890 to 1140 cal BP	675 to 900 cal BP	2760 to 2930 cal BP*	1265 to 1765 cal BP*	2710 to 3290 cal BP*	
	<i>Epinephelus marginatus</i>														
	<i>Epinephelus</i> sp.														
	<i>Mycterperca</i> sp.														
Coryphaenidae	<i>Coryphaena hippurus</i>														
Carangidae	<i>Caranx hippos</i>														
	<i>Caranx</i> sp.														
	<i>Oligopites saurus</i>														
	<i>Selar crumenophthalmus</i>														
	<i>Selene vomer</i>														
Luftianidae	<i>Luftianus synagris</i>														
	<i>Luftianus</i> sp.														
	<i>Ocyurus chrysurus</i>														
	gen., sp. ind.														
Gerreidae	<i>Diapterus rhombus</i>														
	<i>Gerres cinereus</i>														
Haemulidae	gen., sp. ind.														
	<i>Anisostrenus virgicus</i>														
	<i>Anisostrenus</i> sp.														
	<i>Haemulon aurolineatum</i>														
	<i>Haemulon sciurus</i>														
	<i>Haemulon steindachneri</i>														
	<i>Haemulon</i> sp.														
	<i>Orthopristis ruber</i>														
Sparidae	<i>Archosargus</i>														

(Continued)

PLOS ONE | DOI:10.1371/journal.pone.0154476 June 29, 2016

Region	SAQUAREMA				NITEROI		ANGRA DOS REIS				ILHA GRANDE	ILHA DO CABO FRIO		Figures		
Shellmounds	Beirada	Saquarema	Manitiba	Girau	Camboinhas		Algodão	Bigode	Caleira	Peri	Major	Acalá	Usiminas	Ilha do Cabo Frio	Structures	Figures
	Sandy coastal plain and coastal lagoons				Sandy beach		Rocky bottom and rocky islet							Rocky Island	Frio Sandy beach	
Geomorphology																
Radiocarbon age ranges	3035 to 5595 cal BP	2100 to 4200 cal BP	3695 to 4515 cal BP	3730 to 4525 cal BP	4160 to 4960 cal BP		2345 to 4414 cal BP	3223 to 3525 cal BP	1875 to 2175 cal BP	890 to 1140 cal BP	675 to 900 cal BP	2760 to 2930 cal BP*	1265 to 1765 cal BP*	2710 to 3290 cal BP*		
	<i>Archosargus</i> sp.															
	<i>Diplodus</i> sp.															
Scleractinia	<i>Pagrus pagrus</i>															
	<i>Bairdiella ronchus</i>															
	<i>Cynoscion acoupa</i>															
	<i>Cynoscion jamalcensis</i>															
	<i>Cynoscion microlepidotus</i>															
	<i>Larinus breviceps</i>															
	<i>Micropogonias furnieri</i>															
	<i>Pareques acuminatus</i>															
	<i>Pogonias cromis</i>															
	<i>Umbina coroides</i>															
Mugilidae	<i>Mugiliza</i>															
	<i>Mugil</i> sp.															
Labridae	<i>Bodianus rufus</i>															
Scoridae	<i>Scarus</i> sp.															
Trichiuridae	<i>Sparisoma</i> sp.															
Scombridae	<i>Trichiurus lepturus</i>															
	<i>Katsuwonus pelamis</i>															
Istiophoridae	<i>Scomberomorus</i> sp.															
	<i>Istiophorus</i>															

(Continued)

Table 2. (Continued)

Region	SAQUAREMA			NITEROI		ANGRA DOS REIS			ILHA GRANDE		ILHA DO CABO FRIO		Figures				
	Shellmounds	Geomorphology		Beirada	Saquarema	Manitiba	Girau	Cambolhas	Algodão	Bigode	Caleira	Peri		Major	Acaia	Usiminas	Ilha do Cabo Frio
Radiocarbon age ranges																	
Ephippidae	<i>Chaetodipterus faber</i>	•															
Sphyaenidae	<i>Sphyaena barracuda</i> <i>Sphyaena guachancho</i> <i>Sphyaena</i> sp.	•															
Hyporhamphidae	<i>Hyporhamphus unifasciatus</i> <i>Lagocephalus laevis</i>	•															
Tetraodontidae	<i>Chilomycterus spinosus</i> <i>Diodon</i> sp.	•															

Ages are based on radiocarbon analyses of otoliths (*M. turneri*). Ages with asterisks indicate that otolith radiocarbon ages were not available and charcoal- and shell-derived ages were used instead ([29,102]; personal communication).

doi:10.1371/journal.pone.0154476.t002




Fig 2. Modeled age based on whitemouth croaker otolith samples from multiple archaeological layers of Manitoba in Saguarema using OxCal v 4.2.3 [99]. This figure shows ages obtained by radiochronology (vertical lines). The error bar, represented by the horizontal line, shows a range of ages that spans

approximately 370 years across the seven archaeostratigraphic sections, hindering a precise dating of the layers.

doi:10.1371/journal.pone.0154476.g002

lagoons. These remains (i.e., shark teeth and vertebrae) have been recovered in 100% of the analyzed shellmound samples, especially from those associated with rocky islands and islets. The remains of the spotted eagle ray, *Aetobatus narinari*, were the most frequent from the ray group (batoids). The frequency of bony fish remains suggest that some species of groundfish, inhabitants of sandy and muddy bottom or rocky spots, such as croakers (Sciaenidae), snappers (Lutjanidae), grunts (Haemulidae), groupers (Serranidae) and snooks (Centropomidae), were common target species. Sea catfish (Ariidae) and mullet (Mugilidae) species were frequent in localities with sandy and muddy bottoms associated with coastal lagoons. Rocky reef fishes, which are mostly durophagous (i.e., feeding on mollusks, echinoids and crabs), are represented by species of parrotfish (Scaridae), pufferfish (Tetraodontidae) and porcupinefish (Diodontidae), and their remains were particularly frequent in localities associated with rocky islands. Pelagic fish, including bluefish (Pomatomidae), sailfish (Istiophoridae), dolphinfish (Coryphaenidae), jacks (Carangidae), mackerel and tuna (Scombridae), were frequent in the shellmounds located in coastal and rocky islands oceanfront areas.

Based on the analyzed fish assemblages from shellmounds in Rio de Janeiro State, the near-shore fishery remains analyzed here were deposited 5,595 cal BP in the lagoon region of Saquarema and in the oceanfront region of Niterói (Fig 3, black circles). They were characterized by catches of coastal species associated with sandy bottoms and coastal lagoons. The fish could be accessible using beach seines during reproductive aggregation and spawning. These schools of fish include croaker, drum, catfish, mullet and snook. Later, approximately 4,414 cal BP, fishery activity records suggest a targeting of pelagic resources in protected rocky bays and around coastal rocky islets (Fig 3, black triangle). Rocky reef fishes were also a common target, and advances in artisanal fishery and multi-gear techniques remained successful until colonial times. At least since 3,290 cal BP, the fish assemblage recovery from the shellmounds located on the oceanic islands of Cabo Frio and Ilha Grande (Fig 3, black square) suggests a clear predominance of pelagic fisheries and a secondary use of rocky reef species.

The estimation of shark body size (TL) (Table 3) indicates that the spinner shark, *Carcharhinus brevipinna*, ranged from 44.3 to 263.1 cm; the sand tiger shark, *Carcharias taurus*, ranged from 79.3 to 192.2 cm; unidentified shark species of the genus *Carcharhinus* ranged from 38.8 to 115.1 cm; the hammerhead shark, *Sphyrna* sp., ranged from 10.1 to 40.0 cm; and the great white shark, *Carcharodon carcharias*, ranged from 88.2 to 249.6 cm.

Estimation of the body size distribution of whitemouth croaker, *Micropogonias furnieri* [range: 17.8 to 84.8 cm TL in all shellmounds (mean length \pm SD: 43.3 \pm 9.9 cm)]. These ichthyoarchaeological data overlap the modern size distributions from modern fisheries in Itaipu and Angra dos Reis in Rio de Janeiro State (Fig 4). However, the body size distribution of whitemouth croaker catches from prehistoric fisheries shows a probabilistic tendency toward higher frequencies of large specimens, resulting in an estimated 28% reduction in body size based on modern catches.

The results of ANOVA analyses (Kruskal-Wallis) on the median body sizes of whitemouth croaker show that, among the localities (Chi-squared = 1,042.3; df = 4, p values < 0.05), Beirada and Ponte do Girau shellmounds have significant similarities, and both are significantly different from other localities based on an *a posteriori* test (p values < 0.05). The comparative values of median body sizes from various environments and coastal geomorphologies (Chi-squared = 965.4, df = 2, p values < 0.05) reveal significant differences based on a *a posteriori* tests (p values < 0.05) (Fig 5).




Fig 3. Cluster analysis of shellmound similarities. Age ranges (green letters) based on otolith data. Calibration was performed using Marine13 [100] in the 2-sigma range. The overall ΔR was found to be -56.5 to 120.5 [101]. The results at specific sites are as follows: ΔR Saquarema: -272 to -8 [53] and ΔR Manitoba: -224 to 60 [55]. Question marks indicate that radiocarbon dating of otoliths was not available; consequently, we plot shellmound age ranges (red numbers) based on charcoal and shell analyses ([29,102]; personal communication).

doi:10.1371/journal.pone.0154476.g003

The overlap of age ranges between the sequence of archaeological layers (Fig 2) dismisses the multilayer interpretation of the body size distribution in a given shellmound, and we use the Manitoba shellmound as the best example of this. However, the median body sizes of white-mouth croakers in different layers of the Manitoba shellmound show different values and an apparent tendency toward cyclicity of medians from the archaeological surface layer toward deeper layers. This could be interpreted as the result of seasonal oscillations in the intensity of the seasonal marine coastal upwelling (Fig 6).

Individual drilled shark teeth included *Carcharodon carcharias*, *Carcharhinus altimus*, *C. leucas*, *C. plumbeus*, *Galeocerdo cuvier* and *Sphyrna mokarran* (Fig 7). The biodiversity of fish

Table 3. Summary of shark species and body size data recorded from the Rio de Janeiro shellmounds.

Common name	Species	N	Shellmound localities	Body size range (cm)	Mean Size (cm)
Spinner shark	<i>Carcharhinus brevipinna</i>	500	ALG, CAM, USI	44.3–263.1	124.1 ± 55.1
Shark	<i>Carcharhinus</i> sp.	87	USI	38.8–115.1	64.6 ± 15.1
Great white shark	<i>Carcharodon carcharias</i>	12	ALG, CAM, USI	88.2–249.6	153.3 ± 58.2
Sand tiger shark	<i>Carcharias taurus</i>	12	ALG, CAM, USI	79.3–192.2	122.1 ± 31.2
Hammerhead	<i>Sphyrna</i> sp.	49	ALG, CAM, USI	10.1–40.0	20.2 ± 5.1

(N) Number of ichthyofaunal remains of vertebrae. Shellmound localities: (ALG) Algodão, (CAM) Camboinhas, (USI) Usiminas.

doi:10.1371/journal.pone.0154476.t003

fauna records was illustrated based on individual diagnostic structures (i.e., otoliths, teeth or bones remains) from the recovered specimens (Figs 8–15).

Discussion

We recognized 97 fish species from the Rio de Janeiro shellmounds based on detailed anatomic analysis of diagnostic structures. However, some taxonomic records (i.e., at least 44 taxa) that were previously cited in technical reports and unpublished theses about the Rio de Janeiro shellmounds could not be located in the MN-UFRJ repository for identification. These unexamined species are not under institutional catalogue records, and their descriptions or illustrations are not available. We, therefore, choose to exclude those ‘specimens’ from our analysis.




Fig 4. Frequency of body size classes of whitemouth croaker, *M. furnieri*. Ichthyofaunal remains (dashed line, N = 5,532) and modern fisheries (solid line, N = 3,914).

doi:10.1371/journal.pone.0154476.g004




Fig 5. ANOVA analyses (Kruskal-Wallis) of medians based on estimates of body size classes of ichthyoarchaeological otoliths of *Micropogonias furnieri*, using R software (R Core Team 2012). (1) *A posteriori* test of body size medians versus shellmound localities. (2) *A posteriori* test of body size medians versus paleoenvironments where the shellmounds are located.

doi:10.1371/journal.pone.0154476.g005




Fig 6. Comparative analyses of body size medians of whitemouth croaker from the multiple archaeological layers in the Manitoba shellmound in the Saquarema region. Note the differences among the layers, from the surface to deeper layers, and the trend of apparent cyclicity. The red line indicates the median of the total data set.

doi:10.1371/journal.pone.0154476.g006




Fig 7. Drilled shark teeth. (1a-b) *Carcharodon carcharias*, lower tooth, Major, MNUFRJ-ZA-146. (2a-b) *Carcharhinus altimus*, upper tooth, Acaia, MNUFRJ-ZA-868. (3a-b) *Carcharhinus leucas*, upper tooth, Ilha do Cabo Frio, MNUFRJ-ZA-869. (4a-b) *Carcharhinus plumbeus*, upper tooth, Caieira, MNUFRJ-ZA-97. (5a-b) *Galeocercus cuvier*, indet. position tooth, Acaia, MNUFRJ-ZA-870. (6a-b) *Sphyrna mokarran*, upper tooth, Algodão, MNUFRJ-ZA-54. Scale bar 1 cm.

doi:10.1371/journal.pone.0154476.g007

Population structures from shellmounds suggest that the estimated body length of coastal species (e.g., carcharhinids) follows a common pattern of juvenile and adult body length. Moreover, very small teeth and vertebrae of lamnids and sphyrnids collected from the shellmounds of Angra dos Reis and Cabo Frio Island indicate the possible presence of pregnant females, neonates and juveniles in a protected coastal area. Similar evidence is provided by large and very small vertebrae of sawfishes, *Pristis* spp. (Fig 9).

The *Carcharhinus brevipinna* size estimation is in agreement with the common length of contemporary specimens (i.e., 250 cm TL [71]), and the species is classified as a threatened species [103]. The estimated sizes of *Carcharias taurus* are two times smaller than those of their living counterpart (i.e., 250 cm TL [71]), which is classified as a vulnerable species [104]. Juvenile sizes of *Carcharhinus* species coincide with the most common length of the living




Fig 8. Shark teeth from Rio de Janeiro shellmounds. (1a-b) *Carcharias taurus*, upper tooth, Usiminas, MNUFRJ-ZA-565. (2a-b) *Alopias superciliosus*, upper tooth, Saquarema, MNUFRJ-ZA-Col.L.Kneip 28151. (3a-b) *Carcharodon carcharias*, upper tooth, Algodão, MNUFRJ-ZA-02. (4a-b) *Isurus oxyrinchus*, upper tooth, Algodão, MNUFRJ-ZA-21. (5a-b) *Lamna nasus*, lower tooth, Algodão, MNUFRJ-ZA-16. (6a-b) *Carcharhinus acronotus*, upper tooth, Algodão, MNUFRJ-ZA-01. (7a-b) *Sphyrna mokarran*, lower tooth, Major, MNUFRJ-ZA-149. (8a-b) *Carcharhinus limbatus*, upper tooth, Algodão, MNUFRJ-ZA-05. (9a-b) *Carcharhinus plumbeus*, upper tooth, Algodão, MNUFRJ-ZA-44. (10a-b) *Galeocerdo cuvier*, tooth, Algodão, MNUFRJ-ZA-17. (11a-b) *Negaprion brevirostris*, upper tooth, Algodão, MNUFRJ-ZA-25. (12a-b) *Rhizoprionodon lalandii*, lower tooth, Bigode, MNUFRJ-ZA-87, and (13a-b) upper tooth, Algodão, MNUFRJ-ZA-70. (14a-b) *Rhizoprionodon porosus*, upper tooth, Bigode, MNUFRJ-ZA-88. (15a-b) *Sphyrna mokarran*, upper tooth, Caieira II, MNUFRJ-ZA-99. (16a-b) *Sphyrna zygaena*, lower tooth, Major, MNUFRJ-ZA-161. Scale bar: 1 cm. Views: labial (1b, 2a-5a, 6b, 7a, 8-10b, 11-13a, 14-15b, and 16a), lingual (1a, 2-5b, 6a, 7b, 8-10a, 11-13b, 14-15a, and 16b).

doi:10.1371/journal.pone.0154476.g008

counterpart. *Sphyrna* spp. size is nine times smaller than the common length estimation of the living counterpart and could be represented here by neonate specimens [in comparison with the adults body size (i.e., 370 cm TL [88])]. *Sphyrna* species are currently classified as an endangered species [105]. The size of *Carcharodon carcharias* is three times smaller than the common body length of their living counterpart (i.e., 541 cm TL for males and 594 cm for females



Fig 9. Ray teeth, vertebrae and dermal denticle from Rio de Janeiro shellmounds. (1a-b) *Pristis* sp., vertebrae, Usiminas, MNUFRJ-ZA-597 and, (2a-b) Algodão, MNUFRJ-ZA-624. (3a-c) *Rhinoptera* sp., tooth, Algodão, MNUFRJ-ZA-498, and (4a-c) Acaia, MNUFRJ-ZA-708. (5a-b) Ray indet., vertebrae, Saquarema, MNUFRJ-ZA-Col.L.Kneip-08125. (6a-b) *Dasyatis centroura*, caudal spine, Saquarema, MNUFRJ-ZA-Col.L.Kneip-087-1. (7a-b) *Actobatus narinari*, lower plate fragment, Usiminas, MNUFRJ-ZA-424. (8–10) *Dasyatidae* indet., dermal denticle, Usiminas, MNUFRJ-ZA-407. Scale bar: 1 cm.

doi:10.1371/journal.pone.0154476.g009

[88]), which is classified as a vulnerable species [106]. The presence of small vertebrae (and some teeth) is suggestive of neonates of Lamnidae, Carcharhinidae, Sphyrnidae and Pristidae species and provides irrefutable support for the presence of pregnant females and juveniles in these nursery areas.

There are three criteria to identify an area as a nursery [107]: (1), an area with a high frequency of sharks; (2), an area to which shark species have a tendency to remain or return for extended periods; (3), an area or habitat that is repeatedly used across years. Sharks' philopatry [108] and the evidence of residence and the site fidelity suggest that oceanic species preferentially return to their exact birthplaces (i.e., natal philopatry) or birth regions (i.e., regional philopatry) for either parturition or mating even though they make long distance movements that would allow them to breed elsewhere [109]. Modern philopatric are represented by Holocene species in the Brazilian shellmounds, and include *Carcharias taurus*, *Carcharodon carcharias*, *Carcharhinus leucas*, *C. limbatus*, *Galeocerdo cuvier* and *Negaprion brevirostris* [109].

Special remarks are made regarding the presence of cosmopolitan sharks with long-distance oceanic migration, such as *G. cuvier* and *C. carcharias*. The tiger shark, *G. cuvier*, spends the majority of its time in the upper 50 m of water and is recorded to migrate approximately 1,100 to 1,800 km [110,111,112]. The white shark, *C. carcharias*, during its 'patrolling' behavior, mostly swam in depths between 5 and 50 m and during migration, swam almost exclusively at the surface [113]. The long distance oceanic migration of this shark is record to be approximately 4,000 km between the Pacific coast of California and Mexico to the Hawaiian islands [113,114]. Both tiger and white sharks were represented in the shellmounds by perforated teeth, some of them recovered as pendant necklaces associated with human burials and bones [115]. Moreover, modern shark nursery areas were recognized in Brazil [116,117].



Fig 10. Shark and ray vertebrae from Rio de Janeiro shellmounds. (1a-c) *Carcharias taurus*, vertebrae, Usiminas, MNUFRJ-ZA-570. (2a-c) *Carcharodon carcharias*, vertebrae, Algodão, MNUFRJ-ZA-625, and (3a-b) Beirada, MNUFRJ-ZA-576. (4a-c) *Carcharias taurus*, vertebrae, Algodão, MNUFRJ-ZA-621. (5a-c) *Carcharhinus brevipinna*, anterior vertebrae, Usiminas, MNUFRJ-ZA-567, (6a-c) anterior vertebrae, Beirada, MNUFRJ-ZA-575, and (7a-c) posterior vertebrae, Usiminas, MNUFRJ-ZA-568. (8a-c) *Carcharhinus* sp., vertebrae, Usiminas, MNUFRJ-ZA-569. (9a-c) *Rhizoprionodon* sp., vertebrae, Usiminas, MNUFRJ-ZA-571. (10a-c) *Carcharhinus* sp., vertebrae, Algodão, MNUFRJ-ZA-620, (11a-c) Acaia, MNUFRJ-ZA-715, and (12a-c) Algodão, MNUFRJ-ZA-623. (13a-c) *Sphyrna* sp., vertebrae, Usiminas, MNUFRJ-ZA-572, and (14a-c) Camboinhas, MNUFRJ-ZA-853. Scale bar: 1 cm.

doi:10.1371/journal.pone.0154476.g010

Estimated teleostean size class ranges in selected shellmounds, based on 5,532 otoliths of *Micropogonias furnieri*, showed a long history of fishery exploitation (ca 5,595 cal BP until today). The frequency distribution of whitemouth croaker, *M. furnieri*, body sizes based on otoliths from the Ponte do Girau, Algodão, Beirada, Manitiba and Camboinhas shellmounds, and the body size (TL) estimates from modern fisheries (i.e., artisanal and semi-industrial) in the Itaipu and Angra dos Reis regions, Rio de Janeiro State (Fig 4), suggest two modal



Fig 11. Teleostean otoliths from the Rio de Janeiro shellmounds. (1) *Albula nemoptera*, Algodão, MNUFRJ-ZA-190. (2a-c) *Bagre marinus*, Algodão, MNUFRJ-ZA-196. (3a-c) *Genidens genidens*, Camboinhas, MNUFRJ-ZA-845. (4) *Hemiramphus* sp., Caieira II, MNUFRJ-ZA-316, and (5) Caieira II, MNUFRJ-ZA-317. (6) *Centropomus ensiferus*, Algodão, MNUFRJ-ZA-206. (7) *Centropomus undecimalis*, Algodão, MNUFRJ-ZA-205. (8) *Epinephelus* sp., Major, MNUFRJ-ZA-269. (9) *Epinephelus marginatus*, Usiminas, MNUFRJ-ZA-521. (10) *Lutjanus synagris*, Algodão, MNUFRJ-ZA-228. (11) *Lutjanus* sp., Major, MNUFRJ-ZA-279. (12) *Ocyurus chrysurus*, Usiminas, MNUFRJ-ZA-519. (13) *Diapterus rhombeus*, Caieira II, MNUFRJ-ZA-306.

(14) *Diplodus* sp., Algodão, MNUFRJ-ZA-215, and (15) Algodão, MNUFRJ-ZA-217. (16) *Gerres cinereus*, Algodão, MNUFRJ-ZA-223. (17) *Anisostremus virginicus*, Algodão, MNUFRJ-ZA-192. (18) *Haemulon aurolineatum*, Usiminas, MNUFRJ-ZA-517. (19) *Haemulon sciurus*, Algodão, MNUFRJ-ZA-184. (20) *Haemulon steindachneri*, Algodão, MNUFRJ-ZA-220. (21) *Archosargus rhomboidalis*, Algodão, MNUFRJ-ZA-194. (22) *Archosargus* sp., Algodão, MNUFRJ-ZA-193. (23) *Bairdiella ronchus*, Algodão, MNUFRJ-ZA-203. (24) *Cynoscion jamaicensis*, Acaiá, MNUFRJ-ZA-876. (25) *Cynoscion microlepidotus*, Algodão, MNUFRJ-ZA-213. (26) *Cynoscion acoupa*, Manitiba, MNUFRJ-ZA-559. (27) *Larimus breviceps*, Algodão, MNUFRJ-ZA-226. (28) *Micropogonias furnieri*, Algodão, MNUFRJ-ZA-232. (29) *Pareques acuminatus*, Manitiba, MNUFRJ-ZA-875.

doi:10.1371/journal.pone.0154476.g011

distributions of body size frequencies, with overlap between prehistoric and modern fisheries. In comparison to the size estimated based on otoliths of *M. furnieri* from the shellmounds, a probabilistic tendency of a reduction in body size of 28% in modern catches may be attributed to overexploitation.

Despite *M. furnieri* being a species with medium resilience to fishing pressure (i.e., criteria based on values of selected life-history parameters: high fertility and high body growth parameter, K [118]), the regional assessments of current stocks indicate overexploitation [89,90]. Results indicate that body size medians between shellmounds and related environment/coastal geomorphology were significantly different (Fig 5). These differences could be interpreted as a consequence of seasonal variations of fishery areas and the intrinsic life history of the species.

The prehistoric fisheries and time series records along archaeostratigraphic sections vary from a single to seven layers based on the chronological sequences of settlements studied here, while the calibrated radiocarbon models, based on fish otoliths and shell samples by layer, show close or overlapping age probability distributions of individual shellmounds [52,53,55]. Consequently, the shallow profile of individual shellmounds studied here from Rio de Janeiro State could be treated arbitrarily as a single unit for the purpose of a fisheries analysis.

The results, in terms of richness, point toward an early stage of overexploitation of medium-to large-sized, vulnerable fish species, characterized by late maturity, slow growth, low reproductive rates, longevity, spawning aggregations and often ovoviviparous and/or viviparous life histories [119,120,121]. Groupers also exhibit slow growth, low reproductive rates and increased longevity, and reproductive adults leave shallow water habitats and move to deep waters after spawning [122,123]. Our results suggest that prehistoric fishing pressure on coastal areas was sufficient to cause the initial phase of population declines [58,124,125].

Prehistoric fishery methods were able to catch medium to large sharks, skipjack tunas, sailfish and groupers and could have included seined or floating gillnets, spears and long-lines. However, more resilient demersal species did not decline as drastically. Therefore, it is plausible that seasonal fisheries that used beach seines during high upwelling productivity contributed to massive catches of groundfish schools. Other small species could be caught by hooks or traps.

Sawfishes were present during prehistoric times in southeastern Brazil [126]; however, nowadays, these species are a nearly extinct taxon [127], and the last record of the great white shark was in the mid 80's [128]. *Carcharias taurus* is a common shark caught in summer in small numbers in artisanal gillnet fisheries on the Rio de Janeiro coast [128]. Nothing is known about the porbeagle shark, *Lamna nasus*, in the Brazilian region, a rare shark caught in longlines from the 60's to the 90's [129]. This species is distributed from southern Brazil and Uruguay to Argentina [130].

Diverse evidence of archaeological artifacts of gorges, bones and shell fishhooks (including an early fishhook, dated to 42,000 yr BP [131]) and fishing lines (from native fiber plants or human hair) used for inshore or pelagic fisheries was recorded worldwide [15,132,133]. The presence of projectile points in archaeological sites in California, USA (~ 12,200 to 11,200 yr BP) associated with marine and aquatic faunal remains is common [134]. Moreover, the only




Fig 12. Teleostean otoliths from the Rio de Janeiro shellmounds. (1) *Pogonias cromis*, Ponte do Girau, MNUFRJ-ZA-562. (2) *Umbrina coroides*, Ilha do Cabo Frio, MNUFRJ-ZA-874. (3) *Mugil liza*, Algodão, MNUFRJ-ZA-233. (4) *Sphyrna guachancho*, Algodão, MNUFRJ-ZA-236. (5) *Hyporhamphus unifasciatus*, Algodão, MNUFRJ-ZA-2230.

doi:10.1371/journal.pone.0154476.g012

known evidence of fishery tools from Brazilian archaeological sites are manufactured bony spear-tips [30], without clear evidence of possible techniques for massive catches or refined manufacture of hooks used for pelagic or large demersal fishes (S4 Appendix).

Shark teeth are culturally significant in the worldview of indigenous mythology, and numerous archaeological burials of human skeletons in Brazilian shellmounds were decorated with necklaces made with drilled shark teeth [115]. Drilled shark teeth are also present in Argentinian and Uruguayan middens [135]. Individually, drilled shark teeth are here represented by *Carcharodon carcharias*, *Galeocerdo cuvier*, *Carcharhinus altimus*, *C. leucas*, *C. plumbeus* and *Sphyrna mokarran* (Fig 7). However, another possibility is the use of shark teeth for tool manufacturing, e.g., affixed to pieces of wood with vegetal fibers, similar to Polynesian artifacts [127], or as points of arrows.

The analysis of the ichthyofauna shows significant differences between expected fish assemblages (i.e., based on modern occurrences in the coastal area) and the record of observed species in shellmounds (Table 2, Figs 7–15). A high diversity of fish species in the shellmounds reveals the fishery and coastal navigation skills of ancient fishermen and the high importance of coastal fisheries for those prehistoric communities. Such prehistoric fishery activities on vulnerable species and special nursery areas could correspond to the beginning of fish stock depletion along the southeastern Brazilian coast. Overexploitation of such coastal fisheries became unequivocally intense during colonial times resulting in the near collapse of natural fish populations, especially affecting large-sized species such as sharks and giant groupers.

However, in terms of natural resources, we assume that eight biases might have affected the ichthyarchaeological samples under study: (1), selective targeting of certain species (i.e., unpalatable taste, poisonous fishes, small sizes, etc.); (2), the inaccessibility of certain available species (e.g., limitations of employed fishing techniques or presence of adverse marine environmental conditions); (3), minimal potential preservation of some species (i.e., some fishes such



Fig 13. Teleostean skeletal remains from the Rio de Janeiro shellmounds. (1a-b) *Aspistor* sp., hyoid bar, Manitiba, MNUFRJ-ZA-537. (2a-b) *Bagre marinus*, frontal, Algodão, MNUFRJ-ZA-500. (3a-c) *Cathorops* sp., dentary, Manitiba, MNUFRJ-ZA-542. (4) *Genidens barbatus*, supraoccipital, Manitiba, MNUFRJ-ZA-538. (5–7) *Genidens* sp., nucal plate, Manitiba, MNUFRJ-ZA-554, and (8a-b) hyoid bar, Manitiba, MNUFRJ-ZA-534. (9a-b) *Sargocentrum* sp., dentary, Acaia, MNUFRJ-ZA-675. (10) *Centropomus parallelus*, supraoccipital, Usiminas, MNUFRJ-ZA-387. (11a-b) *Centropomus undecimalis*, premaxillary, Usiminas, MNUFRJ-ZA-439. (12a-c) *Epinephelus* sp., premaxillary, Usiminas, MNUFRJ-ZA-444, and (13a-c) dentary, Ilha do Cabo Frio, MNUFRJ-ZA-871. (14a-c) *Mycteroperca* sp., dentary, Usiminas, MNUFRJ-ZA-392. (15a-c) *Caranx* sp., premaxillary, Camboinhas, MNUFRJ-ZA-851, and (16a-b) dentary, Acaia, MNUFRJ-ZA-744. (17a-b) *Selene vomer*, premaxillary, Acaia, MNUFRJ-ZA-663, and (18a-b) dentary, Acaia, MNUFRJ-ZA-667. Scale bar 1 cm.

doi:10.1371/journal.pone.0154476.g013




Fig 14. Teleostean skeletal remains from the Rio de Janeiro shellmounds. (1a-b) *Oligoplites saurus*, premaxillary, Usiminas, MNUFRJ-ZA-438. (2a-c) *Lutjanus* sp., dentary, Algodão, MNUFRJ-ZA-858. (3a-b) Gerreidae indet., premaxillary, Algodão, MNUFRJ-ZA-485. (4a-c) *Anisostremus* sp., premaxillary, Algodão, MNUFRJ-ZA-859, and (5a-b) dentary, Algodão, MNUFRJ-ZA-692. (6a-b) *Diplodus* sp., premaxillary, Acaiá, MNUFRJ-ZA-682. (7a-b) *Archosargus* sp., premaxillary, Ilha do Cabo Frio, MNUFRJ-ZA-873. (8a-b) *Sparisoma* sp., upper pharyngeal tooth plate, Acaiá, MNUFRJ-ZA-730. (9) *Pogonias cromis*, pharyngeal tooth, Manitiba, MNUFRJ-ZA-552. (10a-b) *Mugil* sp., opercle, Manitiba, MNUFRJ-ZA-550. (11a-b) *Bodianus rufus*, premaxillary, Acaiá, MNUFRJ-ZA-828, and (12a-b) dentary, Ilha do Cabo Frio, MNUFRJ-ZA-872. (13) *Sparisoma* sp., premaxillary, Usiminas, MNUFRJ-ZA-436. (14a-b) *Scarus* sp., maxillary, Algodão, MNUFRJ-ZA-993. Scale bar 1 cm.

doi:10.1371/journal.pone.0154476.g014




Fig 15. Teleostean skeletal remains from the Rio de Janeiro shellmounds. (1a-b) *Scarus* sp., angulo-articular, Algodão, MNUFRJ-ZA-496, and (2a-c) pharyngeal tooth, Acaiá, MNUFRJ-ZA-724. (3a-c) *Sparisoma* sp., dentary, Acaiá, MNUFRJ-ZA-720, and (4a-b) angulo-articular, Acaiá, MNUFRJ-ZA-666. (5a-b) *Scarus* sp., lower pharyngeal tooth plate, Acaiá, MNUFRJ-ZA-674. (6a-c) *Trichiurus lepturus*, dentary, Acaiá, MNUFRJ-ZA-746. (7a-b) *Katsuwonus pelamis*, maxillary, Acaiá, MNUFRJ-ZA-705, and (8a-c) vertebrae, Acaiá, MNUFRJ-ZA-710. (9) *Scomberomus* sp., hypural complex, Acaiá, MNUFRJ-ZA-701. (10a-b) *Istiophorus albicans*, hypural,

Usiminas, MNUFRJ-ZA-471. (11) *Sphyraena barracuda*, dentary, Usiminas, MNUFRJ-ZA-395. (12a-b) *Lagocephalus laevigatus*, premaxillary, Acaia, MNUFRJ-ZA-679. (13a-c) *Sparisoma* sp., lower pharyngeal tooth plate, Usiminas, MNUFRJ-ZA-428. (14) *Diodon* sp., tooth plate, Usiminas, MNUFRJ-ZA-429. (15a-b) *Chilomycterus spinosus*, dentary, Algodão, MNUFRJ-ZA-487. Scale bar 1 cm.

doi:10.1371/journal.pone.0154476.g015

as sardines could be consumed entirely); (4), non-uniform employment of archaeological sieving techniques (i.e., the use of large mesh sizes and loss of small stingray teeth such as those of *Dasyatis*, *Gymnura*, or *Mobula* species and otoliths, like those of syngnathids, cynoglossids, achirids, atherinids, blenniids and gobiids); (5), archaeological priority (i.e., ichthyoarchaeological remains such as bones, teeth and otoliths could be considered of secondary importance during field activities, except when they exhibit holes, cutting or intentional abrasion); (6), early overexploitation and fish stock depletion (i.e., large and diverse shark species and rocky reef fish are well represented only in the Angra dos Reis shellmounds); (7), the presence of non-diagnostic or broken bones or eroded otoliths (i.e., unclassified species); and (8), missing specimens.

Conclusions

Prehistoric fishery activity along the Rio de Janeiro coast under the influence of coastal marine upwellings was characterized by massive catches of demersal finfish that inhabit sandy and coastal marine lagoons (e.g., *Micropogonias furnieri*), rocky reef fishes caught near islands and islets (e.g., *Epinephelus morio*), and pelagic fishes caught near rocky cliffs and islands (e.g., *Istiophorus albicans*). Shark fisheries could have been located in nursery areas of protected rocky cliff bays in Arraial do Cabo, Cabo Frio Island, Angra dos Reis and Ilha Grande.

Prehistoric records of high elasmobranch diversity in the Ribeira Bay provide clear evidence for the exploitation of natural populations of sharks and rays since pre-colonial times, especially of bigger species such as the porbeagle shark, *Lamna nasus*, the sand tiger shark, *Carcharias taurus*, the great white shark, *Carcharodon carcharias*, and sawfishes, *Pristis* sp. All these are vulnerable species that could have been rare, especially when taking into consideration that today they are rare or present in reduced numbers in the Ribeira Bay or in the adjacent Angra dos Reis region.

Hence, the results produced here should be addressed as a baseline reference of the ichthyodiversity during the prehistoric times, promoting further debate on the relationships established with past fishing activities, as well as changes in local and regional oceanographic systems.

Supporting Information

S1 Appendix. Distribution of shellmounds along the South American coast. (DOCX)

S2 Appendix. Ichthyoarchaeological material. Ichthyological collection, Zooarchaeology, Museu Nacional, Universidade Federal do Rio de Janeiro (UFRJ)—curators: Maria Cristina Tenório, Tânia Lima. (DOCX)

S3 Appendix. Modern material. A. Ichthyological collection, Departamento de Biologia Animal e Vegetal—Instituto de Biologia, Universidade do Estado do Rio de Janeiro (UERJ)—curators: Ulisses Leite Gomes, Maisa da Cruz Lima, Cristina Paragó, Alexandra Pinto Quintans. B. Ichthyological collection, Otoliths, Departamento de Biologia Marinha—Instituto de Biologia,

Universidade Federal Fluminense (UFF)—curator: Orangel Aguilera. C. Ichthyological collection, Dry skeletons, Departamento de Biologia Marinha—Instituto de Biologia, Universidade Federal Fluminense (UFF)—curators: Orangel Aguilera.
(DOC)

S4 Appendix. The unknown prehistoric fishing. Artwork by Eduardo Agelvis.
(DOCX)

Acknowledgments

The authors would like to thank Angela Camardella and Winner Soares for assistance in the shellmounds repository of MN-UFRJ. Filomena Crancio helped us during our fieldtrip to the Saquarema region and the visit to the shellmounds. Fernando Tuna helped us during the acquisition of otolith measurements and the preparation of the osteological collection. We thank David Canabarro for assistance during the field trip to the Cabo Frio Island and for geomorphological explanations. Thanks to Ana Luiza Albuquerque and Andre Belem for constructive discussions about the coastal upwelling in southeastern Brazil. We thank Edson Silva and Agui-naldo Nepomuceno for UFF laboratorial facilities. Thanks also to Werner Schwarzhans for reviewing the otolith identification and Marcelo Ricardo Souza for helping with statistics in the R software. Orangel A. Rodrigues helped us with image editing, Filipe Leite Rufato with the macrophotography of Acaia samples and Thodoris Argyriou with the first English revision of the draft manuscript. The authors wish to especially thank Eduardo Agelvis for the remarkable artwork. Special thanks Aaron O'Dea, Anastasia Yutseva and anonymous reviewers who has greatly improved the manuscript.

Author Contributions

Conceived and designed the experiments: MSL TCPB RAT CMN OA. Performed the experiments: MSL TCPB RAT CMN ARGJ JDCB MH KM OA. Analyzed the data: MSL TCPB RAT CMN ARGJ JDCB MH KM OA. Contributed reagents/materials/analysis tools: LR MCT TL RS MH KM CC OA. Wrote the paper: MSL TCPB OAS. Revised and edited the manuscript: MSL TCPB RAT CMN ARGJ JDCB TL OA.

References

1. Wagner G, Hilbert K, Bandeira D, Tenório MC, Okumura MM. Sambaquis (shell mounds) of the Brazilian coast. *Quat Int.* 2011; 239: 51–60.
2. Lima TA. Dos mariscos aos peixes: um estudo zooarqueológico de mudança de subsistência na pré-história do Rio de Janeiro. M.Sc. Thesis, Universidade de São Paulo. 1991. Available: <http://www.worldcat.org/title/dos-mariscos-aos-peixes-um-estudo-zoo-arqueologico-de-mudanca-de-subsistencia-na-pre-historia-do-rio-de-janeiro/oclc/46754734>.
3. Dos Santos CMC. A Ictiofauna do sambaqui de Cambinhas, Niterói, RJ. Documento de trabalho: *Sér Arqueol.* 1995; 3: 65–72.
4. Magalhaes RMM, Curvelo MA, Mello EMB. O sambaqui de Manitiba I e outros sambaquis de Saquarema, RJ. In: A fauna na alimentação. Documentos de Trabalho, *Sér Arqueol Mus Nac, UFRJ.* 2001; 5: 55–69.
5. Kneip LM. O sambaqui de Manitiba I e outros sambaquis de Saquarema, RJ. *Sér Arqueol Mus Nac. UFRJ.* 2001; 5–91.
6. Gonzalez MMB. Use of *Pristis* spp. (Elasmobranchii: Pristidae) by hunter-gatherers on the coast of São Paulo, Brazil. *Neotrop. Ichthyol.* 2005a; 3: 421–426.
7. Gonzalez MMB. Tubarões e raia na pré-história do litoral de São Paulo. PhD thesis, Universidade de São Paulo, Brasil. 2005b. Available: <http://www.teses.usp.br/teses/disponiveis/71/71131/tde-29092006-114551/pt-br.php>

8. Gonzalez MMB. Utilização de ferrões de raia pelos grupos de pescadores-coletores do litoral de São Paulo. *Cad. CEON Cult. Mater.* 2014; 24: 99–122.
9. Guimarães MB. A ocupação pré-colonial da região dos Lagos, RJ: Sistema de assentamento e relações intersociais entre grupos sambaquianos e grupos ceramistas Tupinambá e de tradição Una. PhD Thesis. Universidade de São Paulo. 2007. Available: <http://www.teses.usp.br/teses/disponiveis/71/71131/tde-05092007-111015/pt-br.php>
10. Klokler D, Villagrán XS, Giannini PCF, Peixoto S, DeBlasis P. Juntos na costa: zooarqueologia e geoarqueologia de sambaquis do litoral sul catarinense. *Rev. Mus. Arqueol. Etnol.* 2010; 20: 53–75.
11. Hilbert LM. Análise ictioarqueológica dos sítios: Sambaqui do Recreio, Itapeva e Dorva, municípios de Torres e Três Cachoeiras, Rio Grande do Sul, Brasil. M.Sc. Thesis, Pontifícia Universidade Católica do Rio Grande do Sul. 2011. Available: <http://tede2.pucrs.br/tede2/bitstream/tede/204/1/430797.pdf>.
12. Fossile T. Peixes na alimentação de povos pré-coloniais: Estudo ictioarqueológico do sambaqui Cubatão I. M.Sc. Thesis, Universidade da Região de Joinville. 2013. Available: https://www.academia.edu/7820800/PEIXES_NA_ALIMENTA%C3%87%C3%83O_DE_POVOS_PR%C3%89-COLONIAIS_ESTUDO_ICTIOARQUEOL%C3%93GICO_DO_SAMBAQUI_CUBAT%C3%83O_I.
13. Baisre JA. Setting a baseline for Caribbean fisheries. *JICA.* 2010; 5: 120–147. doi: [10.1080/15564891003663943](https://doi.org/10.1080/15564891003663943).
14. Wing SR, Wing ES. Prehistoric fisheries in the Caribbean. *Coral Reefs.* 2001; 20: 1–8. doi: [10.1007/s003380100142](https://doi.org/10.1007/s003380100142).
15. Erlandson JM, Rick TC, Braje TJ, Steinberg A, Vellanoweth RL. Human impacts on ancient shellfish: a 10,000 year record from San Miguel Island, California. *J Archaeol Sci.* 2008; 35: 2144–2152.
16. Fitzpatrick SM. Viewing the sea from the reefs. Comment and forum synthesis on Julio Baisre's "Setting a baseline for Caribbean fisheries". *JICA.* 2010; 5: 173–178. doi: [10.1080/15564891003656624](https://doi.org/10.1080/15564891003656624).
17. McClenachan L, Hardt M, Jackson J, Cooke R. Mounting evidence for historical overfishing and long-term degradation of Caribbean marine ecosystems: comment on Julio Baisre's "Setting a baseline for Caribbean fisheries". *JICA.* 2010; 5: 165–169. doi: [10.1080/15564891003656178](https://doi.org/10.1080/15564891003656178).
18. Jones S. Considerations for advancing a dialogue on pre-modern marine exploitation in the Caribbean and beyond: comment on Julio Baisre's "Setting a baseline for Caribbean fisheries". *JICA.* 2010; 5: 159–161. doi: [10.1080/15564891003655998](https://doi.org/10.1080/15564891003655998).
19. Cramer KL. History of human occupation and environmental changes in western and central Caribbean Panama. *Bull Mar Sci.* 2013; 89(4): 955–982. doi: [10.5343/bms.2012.1028](https://doi.org/10.5343/bms.2012.1028).
20. Polis GA, Anderson WB, Holt RD. Toward an integration of landscape and food web ecology: the dynamics of spatially subsidized food webs. *Annu Rev Ecol Syst.* 1997; 28: 289–316.
21. Reitz E. Fishing down the food web: A case study from St. Augustine, Florida, USA. *Am Antiq.* 2004; 69: 63–83.
22. Erlandson JM, Rick TC. Archaeology meets marine ecology: the antiquity of maritime cultures and human impacts on marine fisheries and ecosystems. *Ann Rev Mar Sci.* 2010; 2: 165–185.
23. Rick TC, Kirch PV, Erlandson JM, Fitzpatrick SM. Archeology, deep history, and the human transformation of island ecosystems. *Anthropocene.* 2013.
24. Prous A. *Arqueologia brasileira.* 1st ed. Brasil: Editora da Universidade de Brasília, Brasília Press. 1992.
25. Gaspar MD. Análises das datações radiocarbônicas dos sítios de pescadores, coletores e caçadores. *Bol Mus. Paraense Emilio Goeldi.* 1996; 8: 81–91.
26. Lima TA. Em busca dos frutos do mar: os pescadores-coletores do litoral centro-sul do Brasil. *Rev USP.* 1999–2000; 44: 270–332.
27. Lima TA, Macário KD, Anjos RM, Gomes PRS, Coimbra RS, Elamore E. AMS dating of early shellmounds of the southeastern Brazilian coast. *Braz J Phys.* 2003; 33(2): 276–279.
28. Lima TA, Macário KD, Anjos RM, Gomes PRS, Coimbra MM, Elmore D. The earliest shellmounds of the central-south Brazilian coast. *Nucl Instr Meth Phys Res.* 2004; 691–694.
29. Tenório MC, Afonso MC, Pinto D. Arqueologia do Arraial do Cabo com foco nos sítios da Ilha do Cabo Frio. *Rev Mus Arqueol Etnol.* 2010; 20: 127–145.
30. Guimarães MB. Fishing strategies among prehistoric populations at Saquarema lagoonal complex, Rio de Janeiro, Brazil. *An Acad Bras Ciênc.* 2013; 85(1): 415–429. PMID: [23460437](https://pubmed.ncbi.nlm.nih.gov/23460437/)
31. Macário KD, Souza RCCL, Trindade DC, Decco J, Lima TA, Aguilera AO, et al. Chronological model of a Brazilian Holocene shellmound (sambaqui da Tarioba), Rio de Janeiro, Brazil Radiocarbon. 2014; 56: 489–499.

32. Scartascini FL, Charo M, Volpedo A. Caracterización de las estrategias de obtención de recursos icticos a partir del análisis de otolitos. El caso de la costa norte del Golfo San Matías. In: Salemm M, Santiago F, Álvarez M, Piana E, Vázquez M, Mansur E, editors. Arqueología de Patagonia: una mirada desde el último confin, Tomo II. Utopías. 2009. pp. 845–852.
33. Favier CM, Scartascini FL. Intensive fishery scenarios on the North Patagonian coast (Río Negro, Argentina) during the Mid-Holocene. *Quat Int*. 2012; 256: 62–70.
34. Scartascini FL, Volpedo AV. White croaker (*Micropogonias furnieri*) paleodistribution in the South-western Atlantic Ocean. An archaeological perspective. *J Archaeol Sci*. 2013; 40: 1059–1066.
35. Scartascini FL, Sáez M, Volpedo AV. Otoliths as a proxy for seasonality: The case of *Micropogonias furnieri* from the northern coast of San Matías Gulf, Río Negro, Patagonia, Argentina. *Quat Int*. 2014;.
36. Calippo FR. O surgimento da navegação entre os povos dos sambaquis: argumentos, hipóteses e evidências. *Rev Mus Arqueol Etnol*. 2011; 21: 31–49.
37. Okumura MMM, Boyadjian CHC, Eggers S. Análise da exostose do meato auditivo externo como um marcador de atividade aquática em restos esqueléticos humanos da costa e do interior do Brasil. *Rev Mus Arqueol Etnol*. 2005–2006; 15(16): 181–197.
38. Okumura M. Populações Sambaquianas Costeiras: saúde e afinidades biológicas dentro de um contexto geográfico e temporal. *Rev Temp Acad Arq Pré-Histórica*. 2013; 11: 38–53.
39. Frayer DW. Auditory exostoses and evidence for fishing at Vlasac. *Curr Anthropol*. 1988; 29: 346–349.
40. Hurst W, Bailey M, Hurst B. Prevalence of external auditory canal exostoses in Australian surfboard riders. *J Laryngol Otol*. 2004; 118: 348–351. PMID: [15165308](#)
41. Litu MLA, Al mamun A, Alam Z, Haque R, Hanif A, Hossain A. Exostoses of the external auditory canal. *Bangladesh J Otorhinolaryngol*. 2012; 18: 91–95.
42. Villagran XS, Klokler D, Peixoto S, DeBlasis P, Giannini PCF. Building coastal landscapes: Zooarchaeology and Geoarchaeology of Brazilian shellmounds. *JICA*. 2011; 6: 211–234. doi: [10.1080/15564894.2011.586087](#)
43. Wagner GP, Silva LA. Prehistoric maritime domain and Brazilian shellmounds. *Archaeol. Disc*. 2014; 2 (1): 1–5. doi: [10.4236/ad.2014.21001](#)
44. Barbosa-Guimarães MS. Análise espacial intra-sítio: Um estudo de caso no sítio Ilha da Boa Vista, Cabo Frio, RJ. *BRMNA*. 2006; 64(4): 271–286.
45. Gaspar MD, Klokler D, Bianchini GF. Arqueologia estratégica: abordagens para o estudo da totalidade e construção de sítios monticulares. *Boletim do Museu Paraense Emílio Goeldi. Ciências Humanas*. 2013; 8(3): 517–533.
46. Kneip LM, Pallestirini L, Crancio F, Machado LMC. As estruturas e suas inter-relações em sítios de pescadores-coletores pré-históricos do litoral de Saquarema, RJ. *B IAB*. 1991; 5: 1–42.
47. Turcq B, Martin L, Flexor JM, Suguio K, Pierre C, Tasayco-Ortega L. Origin and evolution of the Quaternary coastal plain between Guaratiba and Cabo Frio, State of Rio de Janeiro, Brazil. In: Knoppers BA, Bidone ED, Abrão JJ, editors. Environmental geochemistry of coastal lagoon systems of Rio de Janeiro. UFF/FINEP. 1999; 6: 25–46.
48. Souto DD, Lessa DVO, Albuquerque ALS, Sifeddine A, Turcq BJ, Barbosa CF. Marine sediments from southeastern Brazilian continental shelf: A 1,200 year record of upwelling productivity. *Palaeogeogr Palaeoclimatol Palaeoecol*. 2011; 299: 49–55.
49. Coelho-Souza SA, López MS, Guimarães JRD, Coutinho R, Candella RN. Biophysical interactions in the Cabo Frio upwelling system, Southeastern Brazil. *Braz J Oceanogr*. 2012; 60: 353–365.
50. Belem AL, Castela RM, Albuquerque ALS. Controls of subsurface temperature variability in a western boundary upwelling system. *Geophys Res Lett*. 2013; 40: 1362–1366.
51. Cordeiro LGM, Belem AL, Bouloubassi I, Rangel B, Sifeddine A, Capilla R, et al. Reconstruction of Southwestern Atlantic Sea Surface Temperatures during the last Century: Cabo Frio Continental Shelf (Brazil). *Palaeogeogr Palaeoclimatol Palaeoecol*. 2014.
52. Macario KD, Souza RCCL, Aguilera OA, Carvalho C, Oliveira FM, Alves EQ, et al. Marine reservoir effect on the Southeastern coast of Brazil: results from the Tarioba shellmound paired samples. *J Environ Radioact*. 2015; 143: 14–19. doi: [10.1016/j.jenvrad.2015.02.002](#) PMID: [25703433](#)
53. Alves E, Macario K, Souza R, Aguilera O, Goulart AC, Scheel-Ybert R, et al. Marine reservoir corrections on the southeastern coast of Brazil: paired samples from the Saquarema shell mound. *Radiocarbon*. 2015; 57: 1–9.
54. Aguilera OA, Angelica R, Macario K, Crapez M, Belem A, Monteiro-Neto C, et al. Fish bone diagenesis in Southeastern Brazilian shell mounds: Paleoenvironmental approach. *Quat Int*. 2015.

55. Carvalho C, Macario K, Oliveira MI, Oliveira F, Chanca I, Alves E, et al. Potential use of archaeological snail shells for the calculation of local marine reservoir effect. *Radiocarbon*, 2015; 57(3): 1–9.
56. Soares de Sousa G. Tratado descritivo do Brasil em 1587. Tipografia Universal Laemmert, Rio de Janeiro, 1851; 1: 518. Available: <http://www.brasiliana.com.br/brasiliana/colecao/obras/9/Tratado-descritivo-do-Brasil-em-1587>.
57. Jackson JBC, Kirby MX, Berger WH, Bjorndal KA, Botsford LW, Bourque BJ, et al. Historical overfishing and the recent collapse of coastal ecosystems. *Science*. 2001; 293–629. doi: [10.1126/science.1059199](https://doi.org/10.1126/science.1059199)
58. Bender MG, Machado GR, Silva PJA, Floeter SR, Monteiro-Netto C, Luiz OJ, et al. Local ecological knowledge and scientific data reveal, overexploitation by multigear artisanal fisheries in the South-western Atlantic. *Plos One*. 2014; 9 10: 110–332.
59. McClenachan L, Jackson JBC, Newman MJH. Conservation implications of historic sea turtle nesting beach loss. *Front. Ecol. Environ*. 2006; 4: 290–296.
60. Antczak A, Buitrago J, Mackowiak de Antczak MM, Guada HJ. A Contribution to the history of marine turtles exploitation in Venezuela. 59th Gulf Caribb. Fish. 2007; 63–73.
61. McClenachan L, Coope A. Extinction rate, historical population structure and ecological role of the Caribbean monk seal. *Proc Math Phys Eng Sci*. 2008; 275: 1351–1358.
62. Castilho PV. Utilization of cetaceans in shell mounds from the southern coast of Brazil. *Quat Int*. 2008; 180: 107–114.
63. Castilho PV, Simões-Lopes PC. Zooarqueologia dos mamíferos aquáticos e semi-aquáticos da ilha de Santa Catarina, sul do Brasil. *Rev Bras Zool*. 2001; 18: 719–727.
64. Castilho P, Simões-Lopes PC. Sea mammals in archaeological sites on the southern coast of Brazil. *Rev Mus Arqueol Etnol*. 2008; 18: 101–113.
65. Savi DC, Tenório MC, Calippo FR, Toledo FAL, Gonzalez MMB, Afonso MC. Beachrock e o sambaqui da Ilha do Cabo Frio. *ABEQUA*, 2005; 10: 29–32.
66. Plew MG. Explorations in American archaeology. Essays in honor of Wesley R. Hurt. Universidade Press of America. 1 st ed. Inc. Boston Press; 1998.
67. Eirado LG, Heilbron M, Almaeida JCH. Os terrenos tectônicos da faixa ribeirana na serra da Bocaina e na baía da Ilha Grande, Sudeste do Brasil. *Rev Bras Geoc*, 2006; 36(3), 426–436.
68. Dias GTM, Kjerfve B. Barrier and beach ridge systems of the Rio de Janeiro coast. In: Dillenburg SR, Hesp P, editors. *Geology and Geomorphology of Holocene Coastal Barriers of Brazil Lectures Notes in Earth Sciences* 107. Springer-Verlag Berlin Heidelberg. 2009. pp. 225–252.
69. Motoki A, Sichel SE, Savi DC, Aires JR. Mecanismo de intrusão dos corpos tabulares de colocação sub-horizontal discordantes, Município de Arraial do Cabo, RJ. *UNESP Geociências*, 2008; 27 (2): 207–218.
70. Castro JWA, Suguio K, Cunha am, Guedes, Tâmega FTS, Rodriguez RR. Rochas de Praia (Beachrocks) da Ilha do Cabo Frio, Arraial do Cabo: Registro Geológico Ímpar da Transição Pleistoceno—Holoceno no Estado do Rio de Janeiro. *Anu. Inst. Geociênc—UFRJ*, 2012; 35(1): 236–241.
71. Tenório MC, Afonso MA, Savi D, Pinto DC, Gonzalez MMB, Amenomori SN, et al. O sítio ou os sítios da Ilha do Cabo Frio: primeiros resultados. In: *An XIII Congr SAB*, 2005, Campo Grande- MS.
72. Kneip LM. O sambaqui de Saquarema, RJ: estrutura e estratigrafia. *Rio de Janeiro, Sér Arqueol Mus Nac UFRJ*, 1995; 3:39–52.
73. Kneip LM. O sambaqui de Manitiba I e outros sambaquis de Saquarema, RJ. *Sér Arqueol Mus Nac UFRJ*. 2001; 1:5–91.
74. Beltrão M. Pré-História do Estado do Rio de Janeiro. *Forense Universitária*, Rio de Janeiro, 2001; 1996 p.
75. Kneip LM, Pallestrini L, Cunha FL. Pesquisas arqueológicas no litoral de Itaipu, Niterói, RJ. *Rio de Janeiro*, 1981.
76. Lima TA. Dos mariscos aos peixes: um estudo zooarqueológico de mudança de subsistência na pré-história do Rio de Janeiro. M.Sc. Thesis, Universidade de São Paulo. 1991. Available: <http://www.worldcat.org/title/dos-mariscos-aos-peixes-um-estudo-zoo-arqueologico-de-mudanca-de-subsistencia-na-pre-historia-do-rio-de-janeiro/oclc/46754734>.
77. Tubino RA, Monteiro-Neto C, Moraes LES, Paes ET. Artisanal fisheries production in the coastal zone of Itaipu, Niterói, RJ, Brazil. *Braz J Oceanogr*, 2007; 55(3): 187–197.
78. Gottschalk S, Kahlert M. Shifts in taxonomical and guild composition of littoral diatom assemblages along environmental gradients. *Hydrobiologia*, 2012; 694(1): 41–56.

79. Rábago-Quiroz CH, López-Martínez J, Valdez-Holguín JE, Nevárez-Martínez M. O, Acevedo-Cervantes A. Fish assemblages in the bycatch of bottom shrimp trawls on the west side of the Gulf of California, Mexico. *Mar Biol*, 2012; 8(9): 865–876.
80. Hughes RM, Herlihy AT, Sifneos JC. Predicting aquatic vertebrate assemblages from environmental variables at three multistate geographic extents of the western USA. *Ecol Indic*, 2015; 57: 546–556.
81. Hammer O, Harper DAT, Ryan PD. PAST: Paleontological statistics software package for education and data analysis. *Palaeontol Electronica*, 2001; 4(1): 9.
82. Goldman KJ, Branstetter S, Musick JA. A re-examination of the age and growth of sand tiger sharks, *Carcharias taurus*, in the western North Atlantic: the importance of ageing protocols and use of multiple back-calculation techniques. *Environ Biol Fishes*, 2006; 77: 241–252.
83. Kotas JE, Mastrochirico V, Petrere M Jr.. Age and growth of the Scalloped Hammerhead shark, *Sphyrna lewini* (Griffith and Smith, 1834), from the southern Brazilian coast. *Braz J Biol*, 2011; 71: 755–761. PMID: 21881801
84. Allen BR, Wintner SP. Age and growth of the spinner shark *Carcharhinus brevipinna* (Muller and Henle, 1839) off the KwaZulu-Natal coast, South Africa. *Afr J Mar Sci*, 2002; 24: 1–8.
85. Natanson LJ. Preliminary investigations into the age and growth of the shortfin mako, *Isurus oxyrinchus*, white shark, *Carcharodon carcharias*, and thresher shark, *Alopias vulpinus*, in the Western North Atlantic Ocean. *Col. Vol. Sci. Pap. ICCAT*, 2002; 54(4): 1280–1293.
86. Desmarest AG. Première décade ichthyologique, ou description complète de dix espèces de poissons 1823 nouvelles, ou imparfaitement connues, habitant la mer qui baigne les côtes de l'île Cuba. *Mém Soc Linn Paris*, 1983; 2: 163–210.
87. Menezes NA, Buckup PA, Figueiredo JL, Moura RL. Catálogo das espécies de peixes marinhos do Brasil. São Paulo: Museu de Zoologia, 2003, 159 p.
88. Froese R, Pauly D, editors. FishBase. 2012; version 06–2012. Available: <http://www.fishbase.org>
89. Carneiro MH, Castro PMG, Tutui SLS, Bastos GCC. *Micropogonias furnieri* (Desmarest, 1823): Estoque sudeste. In: Cergole MC, Ávila-da-Silva AO, Rossi-Wongtschowski CL. Análise das principais pescarias comerciais da região Sudeste-Sul do Brasil: Dinâmica populacional das espécies em exploração. São Paulo, Instituto Oceanográfico-USP; 2005, pp 94–100. Available: http://www.mma.gov.br/estruturas/revizee/arquivos/revizee_analise_prin_pes.pdf
90. Haimovici M, Ignácio JM. *Micropogonias furnieri* (Desmarest, 1823): Estoque sul. In: Cergole MC, Ávila-da-Silva AO, Rossi-Wongtschowski CL. Análise das principais pescarias comerciais da região Sudeste-Sul do Brasil: Dinâmica populacional das espécies em exploração. São Paulo, Instituto Oceanográfico-USP; 2005. pp. 101–107.
91. Mulato IP, Corrêa B, Vianna M. Distribuição espaço-temporal de *Micropogonias furnieri* (Perciformes, Sciaenidae) em um estuário tropical no sudeste do Brasil. *Bol. Inst. Pesca*, 2015; 41(1): 1–18.
92. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria, 2012. ISBN 3-900051-07-0, Available: <http://www.R-project.org>.
93. Netto EB, Gaelzer LR. Associações de peixes bentônicos e demersais na região do Cabo Frio, RJ, Brasil. *Nerítica*, 1991; 6(1–2): 139–156.
94. Silva PJA. Onze anos de produção pesqueira na região de Arraial do Cabo-RJ. M.Sc. Thesis, Universidade Federal Fluminense, Brasil. 2004.
95. Monteiro-Neto C, Tubino RA, Moraes LES, Neto JPM, Gustavo V, Esteves GV, et al. Associações de peixes na região costeira de Itaipu, Niterói, RJ. *Iheringia, Sér Zool*, 2008; 98(1): 50–59.
96. Andreata JV, Meurer BC, Baptista MGS, Manzano FV, Teixeira DE, Longo MM, et al. Composição da assembleia da Baía da Ribeira, Angra dos Reis, Rio de Janeiro, Brasil. *Rev Bras Zool*, 2002; 19(4): 1139–1146.
97. Macario KD, Gomes PRS, Anjos RM, Carvalho C, Linares R, Alves EQ, et al. The Brazilian AMS radiocarbon laboratory (LAC-UFF) and the intercomparison of results with CENA and UGAMS. *Radiocarbon*, 2013; 55: 325–330.
98. Castro MD. Espectrometria de massa com aceleradores. Software para análise de dados e efeito reservatório em Cuba. PhD Thesis, Universidade Federal Fluminense, Instituto de Física, Brasil. 2015.
99. Bronk Ramsey C, Lee S. Recent and planned developments of the program OxCal. *Radiocarbon*, 2013; 55(2–3): 720–730.
100. Reimer RW, Richards DA, Scott EM, Southon JR, Staff RA, Turney CSM, et al. IntCal13 and marine13 radiocarbon age calibration curves 0–50,000 years cal BP. *Radiocarbon*, 2013; 55: 1869–1887.
101. Alves E, Macario K, Souza R, Pimenta A, Douka K, Oliveira F, et al. Radiocarbon reservoir corrections on the Brazilian coast from pebbles marine shells. *Quat Geochronol*, 2015b; 29:30–35.

102. Tenório MC, Pinto DC, Afonso MC. Dinamica de ocupação, contatos e trocas no litoral do Rio de Janeiro no período de 4000 a 2000 anos antes do presente. *Arq Mus Nac*, 2008; 66(2): 311–321.
103. Burgess GH. *Carcharhinus brevipinna*. The IUCN red list of threatened species. 2009; version 2014.3. Available: <http://www.iucnredlist.org>
104. Pollard D, Smith A. *Carcharias taurus*. The IUCN red list of threatened species. 2009; version 2014.3. Available: <http://www.iucnredlist.org>
105. Denham J, Stevens J, Simpfendorfer CA, Heupel MR, Cliff G, Morgan A, et al. *Sphyrna mokarran*. The IUCN red list of threatened species. 2007; version 2014.3. Available: <http://www.iucnredlist.org>
106. Fergusson I, Compagno LJV, Marks M. *Carcharodon carcharias*. The IUCN red list of threatened species. 2009; version 2014.3. Available: <http://www.iucnredlist.org>
107. Heupel MR, Carlson JK, Simpfendorfer CA. Shark nursery areas: concepts, definition, characterization and assumptions. *Mar Ecol Prog Ser*, 2007; 337: 287–297.
108. Hueter RE, Heupel MR, Heist EJ, Keeney DB. Evidence of Philopatry in Sharks and Implications for the Management of Shark Fisheries. *J Northwest Atl Fish Sci*, 2004; 35: 239–247.
109. Chapman DD, Feldheim KA, Papastamatiou YP, Hueter RE. There and back again: a review of residency and return migrations in sharks, with implications for population structure and management. *Ann Rev Mar Sci*, 2015; 7:547–704. doi: [10.1146/annurev-marine-010814-015730](https://doi.org/10.1146/annurev-marine-010814-015730) PMID: [25251267](https://pubmed.ncbi.nlm.nih.gov/25251267/)
110. Holmes BJ, Pepperell JG, Griffiths SP, Jaine FRA, Tibbetts IR, Bennett MB. Tiger shark (*Galeocerdo cuvier*) movement patterns and habitat use determined by satellite tagging in eastern Australian waters. *Mar Biol*, 2014; 161: 2645–2658.
111. Vaudo JJ, Wetherbee BM, Harvey G, Nemeth RS, Aming C, Burnie N, et al. Intraspecific variation in vertical habitat use by tiger sharks (*Galeocerdo cuvier*) in the western North Atlantic. *Ecol Evol*, 2014; 4(10): 1768–1786. doi: [10.1002/ece3.1053](https://doi.org/10.1002/ece3.1053) PMID: [24963376](https://pubmed.ncbi.nlm.nih.gov/24963376/)
112. Werry JM, Planes S, Berumen ML, Lee KA, Braun CD, Clua E. Reef-fidelity and migration of tiger sharks, *Galeocerdo cuvier*, across the Coral Sea. *PlosOne*, 2014; 9(1): 1–18. doi: [10.1371/journal.pone.0083249](https://doi.org/10.1371/journal.pone.0083249)
113. Jorgensen SJ, Arnoldi NS, Estess EE, Chapple TK, Rückert M, Anderson SD, et al. Eating or Meeting? Cluster Analysis Reveals Intricacies of White Shark (*Carcharodon carcharias*) Migration and Off-shore Behavior. *PLoS ONE*, 2012; 7(10): 47819. doi: [10.1371/journal.pone.0047819](https://doi.org/10.1371/journal.pone.0047819)
114. Domeier ML, Nasby-Lucas N. Migration patterns of white sharks *Carcharodon carcharias* tagged at Guadalupe Island, Mexico, and identification of an eastern Pacific shared offshore foraging area. *Mar Ecol Prog Ser*, 2008; 370: 221–237.
115. Gonzalez MMB. Use of *Pristis* spp. (Elasmobranchii: Pristidae) by hunter-gatherers on the coast of São Paulo, Brazil. *Neotrop. Ichthyol.* 2005; 3: 421–426.
116. Bornatowski H. A parturition and nursery area for *Carcharhinus limbatus* (elasmobranchii, carcharhinidae) off the coast of Paraná, Brazil. *Braz J Oceanogr*, 2008; 56(4), 317–319.
117. Yokota L, Lessa RP. A nursery area for sharks and rays in Northeastern Brazil. *Environ Biol Fish*, 2006; 75: 349–360. doi: [10.1007/s10641-006-0038-9](https://doi.org/10.1007/s10641-006-0038-9)
118. Musick JA. Criteria to define extinction risk in marine fishes. The American Fisheries Society initiative. *Fisheries*, 1999; 24(12): 6–14.
119. Ferretti F, Worm B, Britten GL, Heithaus MR, Lotze HK. Patterns and ecosystem consequences of shark declines in the ocean. *Ecol Lett*, 2010; 13: 1055–1071. doi: [10.1111/j.1461-0248.2010.01489.x](https://doi.org/10.1111/j.1461-0248.2010.01489.x) PMID: [20528897](https://pubmed.ncbi.nlm.nih.gov/20528897/)
120. Ward-Paige CA, Mora C, Lotze HK, Pattengill-Semmens C, McClenachan L, Arias-Castro E, et al. Large-Scale absence of sharks on reefs in the Greater-Caribbean: A footprint of human pressures. *PLoS One*, 2010; 5(8): 11968. doi: [10.1371/journal.pone.0011968](https://doi.org/10.1371/journal.pone.0011968)
121. Worm B, Davis B, Kettner L, Ward-Paige CA, Chapman D, Heithaus MR, et al. Global catches, exploitation rates, and rebuilding options for sharks. *Mar Policy*, 2013; 40: 194–204.
122. Sadovy Y, Eklund AM. Synopsis of biological data on the Nassau grouper, *Epinephelus striatus* (Bloch, 1792), and the Jewfish, *E. itajara* (Lichtenstein, 1822). NOAA Tech Rep NMFS, 1999; 146: 1–65.
123. Starr RM, Sala E, Ballesteros E, Zabala M. Spatial dynamics of the Nassau grouper *Epinephelus striatus* in a Caribbean atoll. *Mar Ecol Prog Ser*, 2007; 343: 239–249.
124. Morris AV, Roberts CM, Hawkins JP. The threatened status of groupers (Epinephelinae). *Biodivers Conserv*, 2000; 9: 919–2000.
125. Aguilar-Perera A. Disappearance of a Nassau grouper spawning aggregation off the southern Mexican Caribbean coast. *Mar Ecol Prog Ser*, 2006; 327: 289–296.

126. Gonzalez MMB. Tubarões e raias na pré-história do litoral de São Paulo. PhD thesis, Universidade de São Paulo, Brasil. 2005. Available: <http://www.teses.usp.br/teses/disponiveis/7/7171131/tde-29092006-114551/pt-br.php>
127. Drew J, Philipp C, Westneat MW. Shark tooth weapons from the 19th century reflect shifting baselines in central Pacific predator assemblies. PLoS One, 2013; 8(4): 59855. doi: [10.1371/journal.pone.0059855](https://doi.org/10.1371/journal.pone.0059855)
128. Gadig OBF, Rosa RS. Occurrence of the white shark along the Brazilian coast. In: Kimley AP, Ainley DG, editors. Great white sharks: The biology of *Carcharodon carcharias*. Academic Press. 1998. 517 pp.
129. Mazzoleni RC, Schwingel PR. Elasmobranch species landed in Itajaí harbor, Southern Brazil. Notas Téc. FACIMAR, 1999; 3:111–118.
130. Pons M, Domingo A. Standardized CPUE of porbeagle shark (*Lamna nasus*) caught by the uruguayan pelagic longline fleet (1982–2008). Collect. Vol. Sci. Pap. ICCAT, 2010; 65(6): 2098–2108.
131. Salls R. To catch a fish: some limitations on prehistoric fishing in southern California with special reference to native plant fiber fishing line. J Ethnobiol, 1989; 9(2): 173–199.
132. Walker KJ. The material culture of Precolumbian fishing: artifacts and fish remains from coastal South-west Florida. Southeast Archaeol, 2000; 19(1): 24–45
133. O'Connor S, Ono R, Clarkson C. Pelagic fishing at 42,000 years before the present and the maritime skills of modern humans. Science, 2011; 334: 111–121.
134. Erlandson JM, Rick TC, Braje TJ, Casperson M, Culleton B, Fulfrost B et al. Paleocoastal seafaring, maritime technologies and coastal foraging on California's Channel Islands. Science, 2011; 331: 1181–1185. doi: [10.1126/science.1201477](https://doi.org/10.1126/science.1201477) PMID: [21385713](https://pubmed.ncbi.nlm.nih.gov/21385713/)
135. Cione AL, Bonomo M. Great white shark teeth used as pendants and possible tools by Early-Middle Holocene terrestrial mammal hunter-gatherers in the Easter Pampas (Southern South America). Int J Osteoarchaeol, 2003; 13: 222–231. doi: [10.1002/oa.678](https://doi.org/10.1002/oa.678)